

ROLE OF SOME RELATIVELY NOVEL ALTERNATIVES IN INCREASING INSECTICIDES TOXICITY AGAINST SOME SUCKING PESTS

Nassef, A. M.

Plant Protection Institute A.R.C., Egypt

ABSTRACT

Laboratory experiments were carried out to evaluate the bioactivity of certain traditional insecticides (i.e. profenofos, chlorpyrifos-methyl, monocrotophos and carbosulfan) against certain sucking pests. The tested alternative were also evaluated when combined in mixtures with conventional insecticides at 1: 10 ratio based on the LC₅₀'s and LC₉₀'s value Profenofos showed the highest toxicity against adults of whitefly, *B. tabaci* (Genn.) and two-spotted spider mite, *T. urticae* (Koch.) (LC₅₀'s 52 & 8 ppm and LC₉₀ 150 & 25 respectively), wherever carbosulfan showed the highest toxicity against aphids (LC₅₀ 75 ppm and LC₉₀ 160 ppm).

Chlorpyrifos-methyl showed moderate toxicity against aphids and mites wherever monocrotophos showed the least. For the joint action pesticides/alternatives at 1: 10, the tested mineral oil Kz, and vegetative oil (black cumin) improved the toxicity of all tested pesticides against aphids and mites. It was obvious that non of the tested alternatives (i.e. mineral oil Kz, natural oil black cumin or liquid detergents) showed synergistic action to profenofos, monocrotophos, and carbosulfan against adults of *Bemisia tabaci* (Genn.), meanwhile, chlorpyrifos-methyl mixtures with Kz oil and vegetative oil black cumin induced a high increase in toxicity 87.6% and 86.5%, also the mixtures of chlorpyrifos-methyl with nastabon increase the toxicity with 28% and 7.96% for LC₅₀ and LC₉₀ values, respectively.

INTRODUCTION

Sucking pests e.g. whitefly, *Bemisia tabaci* (Genn.), green peach aphids, *Myzus persicae* (Sulz.), and two spotted spider mites, *Tetranychus urticae* (Koch) are deleterious one stacking many crops and vegetables.

The economic importance of these pests is attributed to their high reproductive potential and transmit plant pathogenic viruses (Harris and Maramorsch, 1977). Also extracting large quantities of phloem sap which can result in greater than 5% yield reduction (Loiyd, 1922).

Chemical control measures are usually effective, potentially cover the control of most pests and diseases and may be used effectively under a wide variety of condition.

On the other hand the wide use of insecticides currently registered for control resulted many problem such as chemical resistance and hazards to mammals (Prabhaker *et al.*, 1985).

The honeydew excreted by these homopteran insects serves as a medium for sooty mold fungi (Perkins, 1983 and 1987). For all that, the aim of the present study is to find out relatively novel alternatives reduce dosage of insecticides through mixing process, relatively, low cost material compared to most insecticides and low human health hazards.

MATERIALS AND METHODS

I. Pesticides tested and their alternatives:

a. Pesticides:

- Selecron (profenofos) 72% EC.
0-4-bromo-2-chlorophenyl o-ethyl s-propyl phosphorothioate.
- Reldan (chloropyrifos-methyl) 50% EC.
o,o-dimethyl o-3, 5,6-trichloro-2-pyridyl phosphorothioate.
- Novacron (monocrotofos) 40% SCW.
dimethyl (E)-1 methyl-2-(methyl carbamyl) vinyl phosphate.
- Marshal (carbosulfan) 25 WP.
2,3 dihydro-2, 2-dimethyl benzofuran-7 yl (dibutylaminothio) methyl carbamate.

b. Mineral oil:

Kz oil 95% EC provided by Kafr El-Zayat Comp. of pesticides, Egypt.

c. Vegetative oil:

A natural oil was isolated from seed of black cumin, *Nigella sativum*.

d. Liquid detergent:

Nastabon and Oki.

II. Test arthropods:

a. Whitefly, *Bemisia tabaci* (Genn.)

Laboratory strain reared as described by Azab *et al.*, 1971 with minor modification.

The toxicity of the tested pesticides and pesticides/alternatives was investigated. Serial concentration of toxicants were prepared using water. Triton X100 was added when testing the natural oil as an emulsifier. The bioassay technique described by El-Helaly *et al.* (1976) was adopted with some modification.

Data obtained were corrected by using Abbott's formula (1925) percentages of increase in toxicity were calculated according to (Sun 1950).

b. Aphids, *Myzus persicae* (Sulz.):

The green peach aphid, *Myzus persicae* (Sulz.) (Homoptera: Aphididae) was reared in the laboratory according to the method adopted by Norman and Sutton (1967).

The rapid-dip test (FAO Methods No. 17 1980) was applied to evaluate the toxicity of the tested compounds against apterous adults of aphids. Mortality percentages were calculated and corrected according to Abbott's formula (1925) for pesticides/alternatives mixtures, increase in toxicity was calculated.

c. Mites *Tetranychus urticae* (Koch.):

Original strain of *T. urticae* was provided and identified by Acrology Dept. Plant Protection, Institute, Sakha Agriculture Research Station.

The leaf-disc dip methods (Siegler, 1947) was used to evaluate the efficiency of pesticides or pesticides/alternatives mixtures at 1: 10 ratio against *Tetranychus urticae* by means of bionocular, mortality counts were recorded and corrected by Abbott's formula 1925. Increase in toxicity was calculated by the equation of Sun (1950).

RESULTS AND DISCUSSION

I. Insecticidal activity against aphids:

Results presented in Table (1), showed that carbosulfan had the highest toxicity followed by profenofos LC₅₀'s, 75 and 88 ppm and LC₉₀'s 160 and 220 ppm respectively. Monocrotophos showed week toxicity (460 and 960 ppm), wherever, chlorpyrifos-methyl showed moderate toxicity. This is logical because carbosulfan acts as an aphicide. (Halawa *et al.*, 1992; Nassef, 1998) reported that furathiocarb showed the highest toxicity followed by pirimiphos methyl (LC₅₀'s 32, 150 ppm, respectively). The results were agreed with our results.

Table (1): The effect of some insecticides and its mixture with some oils and detergents to aphid *Myzus persicae* (Sulz.).

Pesticides and it mixtures	LC ₅₀	LC ₉₀	% increase in toxicities at	
			LC ₅₀	LC ₉₀
Profenofos	88	220	-	-
+ (1) Kz oil	9	13	89.7	94.09
+ (2) Black cumin oil	7	13	92.0	94.09
+ (3) Nastabon	32	60	63.6	72.73
+ (4) Oki	20	48	77.3	78.18
Chlorpyrifos-methyl	220	600	-	-
+ (1) Kz oil	100	380	54.5	36.67
+ (2) Black cumin oil	108	290	50.9	51.67
+ (3) Nastabon	480	920	-118.2	-53.33
+ (4) Oki	1000	3800	-354.5	-533.33
Monocrotophos	460	900	-	-
+ (1) Kz oil	220	400	52.17	55.56
+ (2) Black cumin oil	160	500	65.22	44.44
+ (3) Nastabon	400	1000	13.04	-11.11
+ (4) Oki	1300	4000	-182.6	-344.4
Carbosulfan	75	160	-	-
+ (1) Kz oil	20	42.5	73.33	73.44
+ (2) Black cumin oil	23	76	69.33	52.50
+ (3) Nastabon	170	500	-126.67	-212.50
+ (4) Oki	95	260	-26.67	-62.50

*% Increasing toxicity =

$$\frac{LC_{50} \text{ or } LC_{90} \text{ of insecticides alone} - LC_{50} \text{ or } LC_{90} \text{ of insecticides in mixture}}{LC_{50} \text{ or } LC_{90} \text{ of insecticides alone}} \times 100$$

For the joint action of pesticides/alternatives combination (1: 10) results are recorded in Table (1) evident that the tested mineral oil (Kz oil) and the vegetative oil (Black cumin) improved the toxicity of all tested pesticides, the effect was more pronounced with profenofos % increase in toxicity 89.7 and 94.09, 92.0 and 94.09 at LC₅₀'s and LC₉₀'s respectively. Such results might be of great importance in IPM programmes where it is desirable to control simultaneously more than one pest (i.e. aphid and mites for instance). Improving the toxicity of insecticides against aphid by mixing with mineral oils were reported by many authors (Omar *et al.*, 1987; Mourad *et al.*, 1994; Hydar *et al.*, 1996 and Nassef, 1998). The two detergents Oki and Nastabon showed

antagonistic action when added to chlorpyrifos-methyl, monocrotophos and carbosulfan. On the other hand, profenofos mixed with Nastabon and Oki at 1: 10 ratio, induced toxicity to aphids more than to that produced by carbosulfan, the recommended aphicide that most toxic compound (LC₅₀'s 32 and 75, respectively).

II. Insecticidal activity against adults of whitefly, *Bemisia tabaci* (Genn.):

Results of toxicity of pesticides against adults of *B. tabaci* (Genn.) are recorded in Table (2), based on L.C₅₀ values, it was cleared that, Profenofos had the highest toxicity followed by Monocrotophos and Carbosulfan with LC₅₀'s 52, 110, 110, ppm, respectively. Chlorpyrifos-methyl showed moderate toxicity with LC₅₀ 250 ppm. The results partly agreed with those of Ayoub (1994) who found that pirimiphos-methyl and IGR, diafenthiuron had relatively high values of LC₅₀'s (21.5 and 26.25 ppm, respectively).

Compared with those of deltamethrin or profenofos (5.2 and 7.1 ppm, respectively).

The joint action of conventional pesticides and their alternatives mixtures at 1: 10 ratio was evaluated against the adults of whitefly. It is obvious that none of the tested alternatives (i.e. mineral oil Kz, natural oil (black cumin) or liquid detergents) showed synergistic action to profenofos, monocrotophos and carbosulfan except the mineral oil Kz increase the toxicity of carbosulfan by 10% increase in toxicity. On the other hand, chloropyrifos-methyl mixtures with Kz oil and vegetative oil black cumin induced a high increase in toxicity (80-70% and 86.50%, respectively). Data showed that the use of chlorpyrifos-methyl mixtures with detergents nastabon increase the toxicity of 60th LC₅₀ and LC₉₀ values. Ahmed and Gardiner (1987); Grunber, 1967, reported that, in some case petroleum oils increased toxicity and persistence of contact insecticides and serve as a useful addition to various insect control agents. The mechanisms of this phenomenon has not yet been established. Petroleum oil may increase the uptake of the toxicant by the insect or reduce its evaporation, dissipation or both (Ishaaya *et al.*, 1986). In some insect species, petroleum oils inhibit respiration (Grünberg 1967). Addition of petroleum oil to fenopropathrin under high volume spray conditions resulted in higher mortality of *B. tabaci* adults (Ishaaya *et al.*, 1986), Nassef, 1998 reported that furathiocarb was potentiated by the natural oil or detergents Oki and strongly by mineral oil Kz synergistic ratio (SR) 1.15, 1.22, and 3.33, respectively.

III. Pesticidal activity against mites:

The toxicity and joint action of some pesticides and the mixtures of pesticides and some alternatives were tested against *Tetranychus urticae* (Koch.). The results of toxicity tests were recorded in Table (3). It is clear that, a high toxicity was exhibited by profenofos followed by carbosulfan and chlorpyrifos-methyl (LC₅₀'s 8, 52 and 59, ppm, respectively). The same trend obtained at (LC₉₀'s 25, 120 and 220 ppm). Shoukry *et al.* (1989) found that LC₅₀ of pirimifos-methyl to adult females of *T. urticae* using the same technique was 134.04 ppm. For the joint action of pesticides/alternatives 1: 10

mixtures, results revealed that the mineral oil Kz or vegetative oil remarkably potentiated the tested pesticide. The toxicity of all tested pesticides were greatly increased when mixed with Kz oil or vegetative oil (% increase: 91 & 84 and 83 & 74) and 40 & 86, respectively, at LC₅₀'s.

Table (2): The effect of some insecticide and its mixture with some oils and detergents to adults of whitefly *Bemisia tabaci* (Genn.).

Pesticides and it mixtures	LC ₅₀	LC ₉₀	% increase in toxicities at	
			LC ₅₀	LC ₉₀
Profenofos	52	150	-	-
+ (1) Kz oil	60	750	-15.38	-400
+ (2) Black cummin oil	170	810	-226.9	-433.3
+ (3) Nastabon	350	1450	-573.0	-866.6
+ (4) Oki	250	140	-380.7	6.6
Chlorpyrifos-methyl	250	520	-	-
+ (1) Kz oil	36	100	85.6	80.75
+ (2) Black cummin oil	31	70	87.6	86.5
+ (3) Nastabon	180	780	28.0	7.69
+ (4) Oki	340	1050	-36.0	-101.9
Monocrotophos	110	360	-	-
+ (1) Kz oil	120	1000	-9.1	-177.7
+ (2) Black cummin oil	6	800	45.5	-122.2
+ (3) Nastabon	200	1300	-81.8	-261.1
+ (4) Oki	88	1000	20.0	-177.7
Carbosulfan	110	600	-	-
+ (1) Kz oil	125	540	-13.6	10
+ (2) Black cummin oil	130	4000	-18.2	-566.6
+ (3) Nastabon	1000	8000	-809.8	-123.3
+ (4) Oki	320	1200	-190.8	-100

* % Increasing toxicity =

$$\frac{LC_{50} \text{ or } LC_{90} \text{ of insecticides alone} - LC_{50} \text{ or } LC_{90} \text{ of insecticides in mixture}}{LC_{50} \text{ or } LC_{90} \text{ of insecticides alone}} \times 100$$

The same trend obtained at LC₉₀'s. The mechanisms by which oils increased the toxicity and persistence of contact insecticide was discussed above-mentioned. Mineral and vegetative oils were found to increase the toxicity of pesticides (Ishaaya *et al.*, 1986; Kansouh *et al.*, 1988; Mourad *et al.*, 1994; Nassef, 1998; Gamieh *et al.*, 2000 and Magouz, 2003).

In conclusion, mineral oils (especially Kz), vegetative oil (extracted from black cummin seeds) and detergents especially Nastabon are advantageous as protectants singly or combined with pesticides against some sucking pests.

Table (3): The effect of some insecticide and its mixture with some oils and detergents to mite *Tetranychus urticae* (Koch).

Pesticides and it mixtures	LC ₅₀	LC ₉₀	% increase in toxicities at	
			LC ₅₀	LC ₉₀
Profenofos	8	25	-	-
+ (1) Kz oil	0.72	3.3	91.0	86.8
+ (2) Black cumin oil	1.25	2.5	84.4	90.0
+ (3) Nastabon	2.6	15	67.5	40.0
+ (4) Oki	19	75	-137.5	-200
Chlorpyrifos-methyl	59	220	-	-
+ (1) Kz oil	10	58	83.05	73.63
+ (2) Black cumin oil	10	100	83.05	54.5
+ (3) Nastabon	35	250	46.6	-13.6
+ (4) Oki	660	2500	-1018.6	-1036.3
Monocrotophos	260	560	-	-
+ (1) Kz oil	42	110	83.8	80.35
+ (2) Black cumin oil	66	130	74.6	76.7
+ (3) Nastabon	200	800	23.0	-42.8
+ (4) Oki	12	200	95.4	64.3
Carbosulfan	52	120	-	-
+ (1) Kz oil	31.0	180	40.4	-50
+ (2) Black cumin oil	7.0	54	86.5	55
+ (3) Nastabon	110	3000	-111.5	-2400
+ (4) Oki	2500	< 10000	-4707.6	-

* % Increasing toxicity =

$$\frac{LC_{50} \text{ or } LC_{90} \text{ of insecticides alone} - LC_{50} \text{ or } LC_{90} \text{ of insecticides in mixtured}}{LC_{50} \text{ or } LC_{90} \text{ of insecticides alone}} \times 100$$

REFERENCES

- Abbotts, W.S. (1925). A method of computing the effectiveness of an insecticide. *J. Econ. Entomol.*, (18): 265-267.
- Ahmed, H. and B.G. Gardiner (1967). Effect of mineral oil solvent on the toxicity and speed of action of malathion, *Nature (London)* 214: 1338-1339.
- Ayoub, S.M.A. (1994). Field and laboratory studies on the susceptibility of whitefly *Bemisia tabaci* (Genn.) to insecticides. M.Sc. Thesis. Faculty of Agric. Cairo University.
- Azab, A.K.; M.M. Megahed and D.H. Mersawi (1971). On the biology of *Bemisia tabaci* (Genn.). *Bull. Soc. ent. Egypt*, LV. 1971 [305].
- El-Helaly, M.S.; F.H. El-Gayar and A.Y. El-Shajli (1976). Development of a standard bioassay technique for the adult of *Bemisia tabaci* (Genn.) (Homoptera: Aleyrodidae). *J. Agric. Sci.*, 87: 543-548.
- FAO (1980). Tentative method for the peach-potato aphid. Recommended methods for the detection and measurement of resistance of agricultural pests to pesticides. *FAO Plant Protection Bull*, 18: 16-18.

- Gamieh, G.N.; Sohair E. Saadoon; A.M. Nassef and A.Y. Youns (2000). Efficacy of mineral oils, acaricides and their mixtures against *Tetranychus cucurbitacearum* (Sayed.). Zagazig. J. Agric. Res., 27(2): 591-601.
- Grünberg, A. (1967). Mode of action of mineral oils upon armoured scales (Diaspididae) in conjunction with low-volume application Isr.J. Entomol., 2: 179-186.
- Halawa, L.A.; R.M. Salem; A.A. Korkor and M.Z.F. Awad (1992). Field evaluation of certain pesticides against some sucking insect pests attacking cotton plants. J. Agric. Sci. Mansoura Univ., 17(10): 3366-3371.
- Harris, K.F. and K. Maramorosch (1977). Aphids as virus vectors. Acad. Press, New York. San Francisco and London, 559 pp.
- Haydar, M.F.; M.Z. Dawood and S.M. Ibrahim (1996). The efficiency of mineral oils, some insecticides and their mixtures against *Aphis gossypii* Glover. J. Agric. Sci. Mansoura Univ., 21(6): 2343-2346.
- Ishaaya, I.; M. Austerweil and H. Frankel (1986). Effect of the petroleum oil virol on toxicity and chemical residue of fenprothrin applied against adult of *Bemisia tabaci*. (Homoptera: Aleyrodidae) as high and low volume sprays. J. Econ. Entomol., 79: 596-599.
- Kansouh, A.S.H.; A.L.B. Shawkat and K.G. Ahmed (1988). Control of aphids transmitting mosaic disease to squash plants in Nineveh, Iraq. J. Coll. Agric., King Saud Univ., 1: 191-196.
- Loiyd, L.L. (1922). The control of the greenhouse whitefly (*Asterochiton vaporariorum*) with notes on its biology. Ann. Appl. Biol., 9: 1-34.
- Magouz, R.I. (2003). Integrated Spider Mite Management. Ph.D. Thesis Fac. Agric. Kafr El-Sheikh, Tanta, Univ.
- Mourad, E.I.; M.E. Keddis; M.S. Osman; F.A. Ayad and M. El-Guindy (1994). Suitability of some natural and mineral oils as pesticidal synergists against *Aphis gossypii* Glover Egypt. J. Agric. Res., 72(3): 739-744.
- Nassef, A.M. (1998). Toxicological studies on some sucking pests and their natural enemies. Ph.D. Thesis. Fac. Agric Kafr El-Sheikh, Tanta Univ.
- Norman, P.A. and R.A. Soutton (1967). Host plants for laboratory rearing of melon aphid. J. Econ. Ent., 60(5): 1205-1207.
- Omar, H.I.H.; M.F. Hayder and Y.A. Issa (1987). Ovicidal action of certain pesticides on whitefly *Bemisia tabaci* (Genn.) Agric. Res. Rev., 65(1): 69-75.
- Perkins, H.H. (1983). Identification and processing of honeydew-contaminated cottons. Text. Res. J. pp. 508-12.
- Perkins, H.H. (1987). Sticky Cotton Proc. West. Cotton Prod. Conf. pp. 53-55.
- Prabhaker, N.; D.L. Coudriet and D.E. Meyerdirk (1985). Insect resistance in the sweet-potato whitefly, *Bemisia tabaci* (Homoptera: Aleyrodidae). J. Econ. Entomol., 78: 748-752.
- Shoukry, A.; Y.M. Ahmed; M.A. Mostafa and A.M.M. El-Adawy (1989). Susceptibility of different developmental stages of spider mite, *T. urticae* to certain pesticides. 3rd Conf. of Pests of Dis. of Veg. of Fruits in Egypt and Arab Count. Ismailia, Egypt.

Nassef, A. M.

Siegler, E.H. (1947). Leaf disc technique for laboratory tests of acaricides. J. Econ. Entomol., 40: 617-618.

Sun, Y.P. (1950). Toxicity index, an improved method of comparing the relative toxicity of insecticides. J. Econ. Entomol., 43: 45-53.

دور بعض البدائل الحديثة نسبيا في زيادة سمية المبيدات المستخدمة في مكافحة بعض الآفات الثاقبة الماصة

على معدوح ناصف

معهد بحوث وقاية النباتات - مركز البحوث الزراعية - الدقى

من خلال التجارب المعملية تم تقييم النشاط الابادى لثلاثة من المبيدات الفوسفورية التقليدية ومبيد المن الكربماتى ودراسة التأثير المشترك لبعض بدائل المبيدات من الزيوت المعدنية زيت KZ والزيت المستخلص من حبة البركة واتشان من المنظفات الصناعية السائلة مثل التستابون والـ OKI ضد بعض الآفات الثاقبة الماصة وهى الذبابة البيضاء والمن والعنكبوت الاحمر العادى.

وقد أظهرت النتائج بناء على قيم الجرعات النصفية القاتلة والجرعات القاتلة لـ ٩٠% من الافراد ان اكثر المركبات سمية للذبابة البيضاء والاكاروس هو مبيد البروفينوفوس حيث بلغت التركيزات القاتلة لـ ٥٠% هي ٥٢ ، ٨٠ جزء فى المليون ، بينما بلغت التركيزات القاتلة لـ ٩٠% من الافراد ١٥٠ ، ٢٥ جزء فى المليون بينما أظهرت مبيد الكاربوسلفان سمية عالية ضد المن حيث بلغت قيم الجرعات النصفية القاتلة ٧٥ جزء فى المليون والجرعات القاتلة لـ ٩٠% ١٦٠ جزء فى المليون.

ايضا اظهر مبيد الكلوربيريفوس مثيل سمية متوسطة ضد المن والعنكبوت الاحمر فى حين اظهر المونوكروتفوس سمية ضعيفة ضدها.

أما بالنسبة للتأثير المشترك للمبيدات وبدائلها عند خلطها بنسبة ١ : ١٠ وجد ان الزيت المعدنى KZ والزيت النباتى black cumin أعلى سمية من كل المبيدات المختبرة ضد المن والعنكبوت فى حين لم تظهر أى تنشيط للمبيدات المختبرة ضد الذبابة البيضاء بل كان هناك تضاد فى السمية باستثناء مبيد الكلوربيريفوس مثيل الذى أظهر زيادة فى السمية ٨٧,٦ ، ٨٦,٥ عند قيم الـ LC50 ، LC90 عند خلطه بالزيت النباتى وكذلك ٨٥,٦ ، ٨٠,٧ عند خلطه بالزيت المعدنى. أيضا عند خلطه بالتستابون أدى الى زيادة فى السمية عند قيم LC90, LC50 تقدر بـ ٢٨ ، ٧,٦ زيادة مثل هذه النتائج الواعدة تشجع على استخدام هذه الخلطات فى برامج IPM حين يراد مكافحة أكثر من آفة فى وقت واحد.