

(K)197V
✓

STUDIES ON THE SERPENTINE LEAFMINER *Liriomyza trifolii* (BURGESS) AND ITS HYMENOPTEROUS PARASITOIDS IN NORTH SINAI, EGYPT

Eid, F.M.

Biological Control Research Department, Plant Protection Research Institute, ARC, Egypt.

ABSTRACT

The percentage of infestation caused by the larvae of *Liriomyza trifolii* (Burgess) varied from 23.21 % and 52.08 % in the first season and from 30.83 % and 50.0 % in the second season at El-Arish locality. On the other hand, the percentage of infestation at Rafah locality ranged between (19.48 % and 49.25 %) and (18.88 and 47.68 %) in the first season and the second season, respectively.

The average percentage of parasitism caused by the hymenopterous parasitoids at El-Arish locality were 89.16 % and 87.43 % during the first season 2001 / 2002 and the second season 2002 / 2003, respectively.

At Rafah locality, the average rates of parasitism were 84.58 % and 83.53 % during the first season and the second season, respectively.

INTRODUCTION

The leafminer of both genera *Liriomyza* and *phytomyza* (Diptera: Agromyzidae) are cosmopolitan and are economically important pests of many agricultural crops (Spencer, 1973). In Egypt, *L. trifolii* (Burgess) was recorded as serious pest attacking broad bean, peas, Lentil (Hammad, 1955; Assem, 1966; Hafez et al., 1970; Dimetry, 1971; Attia, 1989 and Eid, 1998). At the present time *L. trifolii* has become an important pest attacking broad bean (Mesbah and Sheriff, 1994 and Awadalla, 1998). Cowpea (Awadalla and Fathy, 1998) and Tomato (Sharaf El -Din 1994). Four parasitoids were encountered by Hafez et al (1974) as parasitoids of *Liriomyza* spp. these parasitoids were *Diglyphus* sp., *Hemiptarsenus zilahisebossi* (Eulophidae) as a larval parasitoids; *Opius* sp. (Barconidae), *Halipectera* sp. (Pteromalidae) as pupal parasitoids. *Diglyphus* sp. proved to be the most common and efficient parasitoid on *L. trifolii*. (Prieto and De ulloa, 1982; Parrella et al, 1983). Both *Diglyphus* sp. and *opius* sp. are widely spread all over the country except at Aswan Governorate, whereas *Halipectera* sp. and *H. zilahisebossi* are restricted only to Giza and Qalubia Governrates and at a very low rate. (Hafez et al 1974).

The present study was conducted to evaluate the percentage of infestation caused by the larvae of *Liriomyza trifolii* as well as the hymenopterous parasitoids on the immature stages of the insect pest.

MATERIALS AND METHODS

This study had been carried out during two seasons, (2001-2002) and (2002-2003) in El-Arish and Rafah localities in North Sinai Governorate. Sampling was carried out in fixed areas at weekly intervals. The number of larvae and pupae varied in each search according to their

Availability. Each infested leaf with larvae and pupae was kept in a glass vial (10x4 cm). The vial was covered with muslin kept in position by

means of a rubber band until the parasitoid emergence. All emerging parasitoids were collected and identified.

Percentage of infestation by faba bean leafminer, *L. trifolii* in the field.

Percentage of infestation by Faba bean leafminer, *L. trifolii* was carried out in Faba bean fields in El-Arish and Rafah in North Sinai Governorate. Ten plants were chosen from the field and the number of infested leaflets related to the whole number of leaflets in the plant was estimated. The average percentage of infestation was then calculated.

Percentage of parasitism:

The percentage of parasitism by all species obtained from collected samples of faba bean was calculated. A study of fluctuation of percentages of parasitism of larvae and pupae of *L.trifolii* was made.

RESULTS AND DISCUSSIONS

The hymenopterous parasitoids that emerged in the laboratory from the leafminer *L. trifolii*.

Samples of *L. trifolii* larvae and pupae collected from faba bean fields during the two seasons of investigation (2001-2002) and (2002-2003) gave rise to the following species of parasitoids (Table 1).

Table (1): The hymenopterous parasitoids emerged from *L. trifolii* immature stages in the laboratory.

| Parasitoids | Family | Host stage |
|----------------------------|--------------|--------------------|
| <i>Halticoptera</i> sp. | Pteromalidae | Larval -Pupa stage |
| <i>Diglyphus isaea</i> | Eulophidae | Larval stage |
| <i>D. crossinervis</i> | Eulophidae | Larval stage |
| <i>Chrysocharis</i> sp. | Eulophidae | Larval -Pupa stage |
| <i>Neochrysocharis</i> sp. | Eulophidae | Larval stage |

Percentage of infestation

Ten plants were chosen from the field and the number of infested leaflets related to the whole number of leaflets in the plant was obtained. Data are presented in Tables 2, 3, 4 & 5

El-Arish locality

The percentage of infestation ranged between 23.21 % in the second week of January 2002 and 52.08 % in the second week of March 2002 during the first season 2001 / 2002, (Table 2). The average percentage of infestation was 29.9 %. On the other hand, the percentage of infestation in the second season 2002 / 2003 at El-Arish locality ranged between 30.83 % and 50.0 % in the second week of February 2003, respectively (Table 3). The average percentage of infestation in the second season 2002 / 2003 at El-Arish locality was 40.27 %.

Table (2): Percentages of Infestation by *L. trifolii* in faba bean fields (10 plants examined per sample) in El-Arish, North Sinai during (2001/2002) season.

| Sampling date | No. of leaves in the sample | | Total no. of leaves in the sample | % Infestation |
|-----------------------------------|-----------------------------|-------------|-----------------------------------|---------------|
| | Infested | Un Infested | | |
| 26/12/2001 | 17 | 80 | 97 | 39.5 |
| 2/1/2002 | 32 | 57 | 89 | 35.96 |
| 9/1 | 26 | 86 | 112 | 23.21 |
| 16/1 | 53 | 84 | 137 | 38.69 |
| 23/1 | 53 | 102 | 155 | 34.19 |
| 30/1 | 81 | 78 | 159 | 50.94 |
| 6/2 | 80 | 109 | 189 | 42.23 |
| 13/2 | 95 | 110 | 205 | 46.34 |
| 20/2 | 87 | 128 | 215 | 40.47 |
| 27/2 | 60 | 150 | 210 | 28.57 |
| 6/3 | 80 | 145 | 225 | 35.56 |
| 13/3 | 125 | 115 | 240 | 52.08 |
| 20/3 | 60 | 131 | 191 | 31.41 |
| 27/3 | 58 | 139 | 224 | 37.95 |
| Total | 732 | 1514 | 2448 | |
| Average percentage of infestation | | | | 29.90 |

Table (3): Percentages of infestation by *L. trifolii* in faba bean fields (10 plants examined per sample) in El-Arish, North Sinai during (2002/2003) season.

| Sampling date | No. Of leaves in the sample | | Total no. Of leaves in the sample | % Infestation |
|-----------------------------------|-----------------------------|-------------|-----------------------------------|---------------|
| | Infested | Un infested | | |
| 11/1/2003 | 24 | 45 | 69 | 34.78 |
| 18/1 | 31 | 59 | 90 | 34.44 |
| 25/1 | 38 | 79 | 117 | 32.48 |
| 1/2 | 55 | 69 | 124 | 44.36 |
| 8/2 | 41 | 92 | 133 | 30.83 |
| 18/2 | 70 | 94 | 164 | 42.68 |
| 22/2 | 83 | 83 | 166 | 50.00 |
| 1/3 | 88 | 107 | 193 | 44.56 |
| 8/3 | 116 | 124 | 240 | 48.33 |
| 15/3 | 94 | 139 | 233 | 40.34 |
| 23/3 | 77 | 133 | 210 | 36.67 |
| 29/3 | 133 | 135 | 268 | 46.50 |
| 4/4 | 87 | 142 | 229 | 37.99 |
| 12/4 | 60 | 125 | 185 | 32.43 |
| Total | 975 | 1426 | 2421 | |
| Average percentage of infestation | | | | 40.27 |

Eld, F.M.

Table (4): Percentages of infestation by *L. trifolii* in faba bean fields (10 plants examined per sample) in Rafah, North Sinai during (2001/2002 season).

| Sampling date | No. of leaves in the sample | | Total no. of leaves in the sample | % Infestation |
|-----------------------------------|-----------------------------|-------------|-----------------------------------|---------------|
| | Infested | Un Infested | | |
| 26/12/2001 | 15 | 62 | 77 | 19.48 |
| 2/1/2002 | 44 | 64 | 108 | 40.74 |
| 9/1 | 38 | 74 | 112 | 33.93 |
| 16/1 | 49 | 76 | 125 | 39.2 |
| 23/1 | 67 | 123 | 190 | 35.26 |
| 30/1 | 61 | 106 | 167 | 36.53 |
| 6/2 | 68 | 102 | 170 | 40 |
| 13/2 | 94 | 105 | 199 | 47.24 |
| 20/2 | 88 | 120 | 208 | 42.32 |
| 27/2 | 75 | 126 | 201 | 37.31 |
| 6/3 | 74 | 160 | 234 | 31.62 |
| 13/3 | 111 | 124 | 235 | 47.23 |
| 20/3 | 84 | 114 | 198 | 42.42 |
| 27/3 | 98 | 114 | 198 | 49.25 |
| Total | 966 | 1470 | 2422 | |
| Average percentage of infestation | | | | 39.88 |

Table (5) Percentages of infestation by *L. trifolii* in faba bean fields (10 plants examined per sample) in Rafah, North Sinai during (2002/2003) season.

| Sampling date | No. of leaves in the sample | | Total no. of leaves in the sample | % Infestation |
|-----------------------------------|-----------------------------|-------------|-----------------------------------|---------------|
| | Infested | Un Infested | | |
| 15/1/2003 | 15 | 58 | 73 | 16.88 |
| 22/1 | 39 | 54 | 93 | 41.94 |
| 29/1 | 24 | 96 | 120 | 20 |
| 5/2 | 45 | 74 | 119 | 37.82 |
| 12/2 | 59 | 130 | 189 | 31.22 |
| 19/2 | 56 | 89 | 145 | 38.62 |
| 26/2 | 58 | 111 | 169 | 34.32 |
| 5/3 | 83 | 125 | 208 | 39.9 |
| 12/3 | 98 | 144 | 242 | 40.5 |
| 19/3 | 70 | 147 | 217 | 32.26 |
| 26/3 | 81 | 143 | 224 | 36.16 |
| 2/4 | 113 | 124 | 237 | 47.68 |
| 9/4 | 77 | 139 | 216 | 35.65 |
| 16/4 | 100 | 126 | 226 | 44.24 |
| Total | 918 | 1560 | 2478 | |
| Average percentage of infestation | | | | 37.05 |

Rafah locality:

The percentage of infestation ranged between 19.48 and 49.25 % in the last week of December 2001 and the end of March 2002, respectively (Table4). The average percentage of infestation during the first season 2001 / 2002 at Rafah locality was 29.88 %.

On the other hand, the minimum percentage of infestation in the second season 2002 / 2003 was 16.88 at the beginning of this study, 2003, while the maximum percentage of infestation was 47.68 % in the first week of April 2003 (Table5). The average percentage of infestation at Rafah locality during the second season 2002 / 2003 recorded 37.05 %.

Percentage of Parasitism:

Number of larvae were chosen at evaluate of the percentage of parasitism. Ecto- Parasitized larvae were easily manipulated. Other larvae remained from the chosen sample were dissected under stereomicroscope to determine the parasitized larvae. Data are shown in Tables (6, 7 & 8)

Table (6): Percentages of parasitism in samples of *L. trifolii* larvae collected from faba bean fields in El-Arish, North Sinal during (2001/2002) season

| Sampling date | No. of larvae in the sample | | Total no. of larvae in the sample | % Parasitism |
|----------------------------------|-----------------------------|------------------------|-----------------------------------|--------------|
| | Parasitized larvae | Non-Parasitized larvae | | |
| 26/12/2001 | 13 | 3 | 16 | 68.25 |
| 2/1/2002 | 11 | 2 | 13 | 70.62 |
| 9/1 | 12 | 5 | 17 | 79.59 |
| 16/1 | 15 | 0 | 15 | 100 |
| 23/1 | 13 | 0 | 13 | 100 |
| 30/1 | 3 | 5 | 8 | 100 |
| 6/2 | 7 | 0 | 7 | 60.5 |
| 13/2 | 24 | 0 | 24 | 80.98 |
| 20/2 | 9 | 0 | 9 | 100 |
| 27/2 | 8 | 0 | 8 | 100 |
| 6/3 | 11 | 1 | 12 | 91.67 |
| 13/3 | 9 | 0 | 9 | 100 |
| 20/3 | 8 | 0 | 8 | 100 |
| 27/3 | 5 | 2 | 7 | 71.42 |
| Total | 148 | 18 | 166 | |
| Average percentage of parasitism | | | | 89.16 |

Table (7): Percentages of parasitism in samples of *L. trifolii* larvae collected from faba bean fields in El-Arish, North Sinal during (2002/2003) season

| Sampling date | No. of larvae in the sample | | Total no. of larvae in the sample | % Parasitism |
|----------------------------------|-----------------------------|------------------------|-----------------------------------|--------------|
| | Parasitized larvae | Non-Parasitized larvae | | |
| 11/1/2003 | 13 | 3 | 16 | 81.25 |
| 18/1 | 11 | 2 | 13 | 84.62 |
| 25/1 | 12 | 5 | 17 | 70.59 |
| 1/2 | 15 | 0 | 15 | 100 |
| 8/2 | 13 | 0 | 13 | 100 |
| 18/2 | 3 | 5 | 8 | 37.50 |
| 22/2 | 10 | 0 | 10 | 100 |
| 1/3 | 24 | 0 | 24 | 100 |
| 8/3 | 9 | 0 | 9 | 100 |
| 15/3 | 8 | 0 | 8 | 100 |
| 23/3 | 8 | 4 | 12 | 91.67 |
| 29/3 | 9 | 0 | 9 | 100 |
| 4/4 | 5 | 2 | 7 | 71.42 |
| 12/4 | 6 | 0 | 6 | 100 |
| Total | 146 | 21 | 167 | |
| Average percentage of parasitism | | | | 87.43 |

El-Arish locality:

The percentage of parasitism caused by hymenopterous parasitoids varied from 60.5 % and 100 % during the first season 2001 / 2002 (Table 6).

On the other hand, the percentage of parasitism ranged between 37.50 % and 100 % during the second season 2002 / 2003 at El-Arish locality (Table 7).

In conclusion, data presented in table 6 and 7 indicated at that, the average percentage of parasitism at the larvae of *L. trifolii* caused by the hymenopterous parasitoids at El-Arish locality were 89.16 % and 87.43 % during the first season 2001 / 2002 and the second season 2002 / 2003, respectively.

Rafah locality:

In conclusion, data presented in table 8 and 9 indicated at that, the average percentage of parasitism at the larvae of *L. trifolii* caused by the hymenopterous parasitoids at Rafah locality were 84.58 % and 83.53 % during the first season 2001 / 2002 and the second season 2002 / 2003, respectively.

Table (8): Percentages of parasitism in samples of *L. trifolii* larvae collected from faba bean fields in Rafah, North Sinai during (2001/2002) season.

| Sampling date | No. of larvae in the sample | | Total no. of larvae in the sample | % Parasitism |
|----------------------------------|-----------------------------|------------------------|-----------------------------------|--------------|
| | Parasitized larvae | Non-Parasitized larvae | | |
| 26/12/2001 | 14 | 4 | 18 | 93.33 * |
| 2/1/2002 | 9 | 1 | 10 | 90 |
| 9/1 | 15 | 4 | 19 | 78.95 |
| 16/1 | 15 | 3 | 18 | 83.33 |
| 23/1 | 13 | 1 | 14 | 92.86 * |
| 30/1 | 8 | 1 | 9 | 88.89 |
| 6/2 | 9 | 1 | 10 | 90 * |
| 13/2 | 17 | 3 | 20 | 85 |
| 20/2 | 8 | 1 | 9 | 88.89 * |
| 27/2 | 8 | 2 | 10 | 80 |
| 6/3 | 12 | 2 | 14 | 85.71 |
| 13/3 | 17 | 2 | 19 | 89.47 |
| 20/3 | 15 | 0 | 15 | 100 * |
| 27/3 | 17 | 0 | 17 | 100 |
| Total | 170 | 32 | 201 | |
| Average percentage of parasitism | | | | 84.58 |

Table (9): Percentages of parasitism in sample of *L. trifolii* larvae collected from faba bean fields in Rafah, North Sinai during (2002/2003) season:

| Sampling date | No. of larvae in the sample | | Total no. of larvae in the sample | % Parasitism |
|----------------------------------|-----------------------------|------------------------|-----------------------------------|--------------|
| | Parasitized larvae | Non-Parasitized larvae | | |
| 15/1 | 13 | 2 | 15 | 86.67 * |
| 22/1 | 8 | 6 | 14 | 57.67 |
| 29/1 | 10 | 1 | 11 | 90.91 |
| 5/2 | 21 | 1 | 22 | 95.47 * |
| 12/2 | 8 | 4 | 12 | 66.67 |
| 19/2 | 6 | 2 | 8 | 60 |
| 26/2 | 8 | 1 | 9 | 88.89 * |
| 5/3 | 15 | 5 | 20 | 75 |
| 12/3 | 12 | 0 | 12 | 100 * |
| 19/3 | 9 | 0 | 9 | 100 |
| 26/3 | 6 | 6 | 12 | 100 |
| 2/4 | 9 | 0 | 9 | 92.783 |
| 9/4 | 7 | 0 | 7 | 100 * |
| 16/4 | 10 | 0 | 10 | 100 |
| Total | 142 | 28 | 170 | |
| Average percentage of parasitism | | | | 83.53 |

Field observation showed the predator *Chrysoperia carnea* Steph. *Thrips* sp. and some coccinellids feeding on the larvae of *L. trifolii* in faba bean fields in the two localities El-Arish and Rafah during the two successive seasons 2001/ 2002 and 2002/ 2003.

REFERENCES

- Assem, M.A. (1966). *Studies on vegetable leaf miners*. Ph.D. Thesis, Fac. of Agric., Cairo Univ.
- Attia, M.B. (1989). Studies on the ecology and control of the broad bean leafminer *Liriomyza trifolii* (Burgess) (Diptera: Agromyzidae) infesting bean fields in Egypt Third National Conf. of Pests & Dis. of Veg. & Fruits, Ismailia, and Vol. I: 278-288.
- Awadalla, s.s. (1998): Relationship between the serpentine leafminer, *Liriomyza trifolii* (Burgess) and its parasitoids on faba bean in Mansoura region. J. Agric. Sci. Mansoura Univ. 23 (9): 4019 - 4026.
- Awadalla, s.s. and H.M.Fathy (1998): Studies on the hymenopterous parasitoids of the serpentine leafminer, *Liriomyza trifolii* (Burgess) in Mansoura region. J. Agric. Sci. Mansoura Univ. 23 (12): 6257 - 6262.
- Dimetry, N.Z. (1971). Biological studies on leaf mining Diptera, *Liriomyza trifolii* (Burgess) attacking beans in Egypt (Diptera: Agromyzidae). Bull. Ent. Soc. Egypt, 55: 55-69.
- Eid, Fawzy, M.H. (1998) Studies on leafminers and their natural enemies in Egypt. Ph.D. Thesis, Fac. of Agric., Cairo Univ., 136 pp.
- Hafez, M.A.H.; M.N. El-Kifl and A.E.A. Wahab (1974). Studies on parasites of *Liriomyza congesta* (Burgess) in Egypt. Bull. Soc. ENT. Egypte, 59: 249-259.
- Hafez, M.A.H.; S. El-Ziady and N.Z. Dimetry (1970). Leaf mining Diptera of vegetables and crops in Egypt. Bull. Soc. ENT Egypte, 54: 389-414.
- Hammad, S.M. (1955). On some dipterous leaf miners from Egypt. Bull. Soc. ENT. Egypte, 39: 391-394.
- Mesbah, I. I. And M. R. Sherif (1994): Population studies on *Liriomyza congesta* (Backer) (Diptera: Agromyzidae) in faba bean fields at Kafr El- Sheikh region. Agric. Sci. Mansoura Univ. 19 (3): 1215 - 1222.
- Parrella, M.P.; G.D. Christie and K.L. Robb (1983). Compatibility of insect growth regulators and *Chrysocharis parksi* (Hymenoptera: Eulophidae) for the control of *Liriomyza trifolii* (Diptera: Agromyzidae) J. Econ. Entomol., 76: 949-951.
- Prieto, A.J. and P. Chaco de Ulloa (1982). Biology and ecology of the chrysanthemum leaf miner *Liriomyza trifolii* Burgess (Diptera: Agromyzidae) in the

Department of Valle del Cauca. Colombiana Revista de Entomologia, 6 (3/4): 77-84.

Sharaf El- Din, A.A.A. (1994): Influence of tomato and four selected weeds on some biological aspects of the serpentine leafminer, *Liriomyza trifolii* (Burgess) (Diptera: Agromyzidae). J. Agric. Sci. Mansoura Univ. 19 (1): 319 - 327.

Spencer, K.A. (1973). Agromyzidae (Diptera) of economic importance. Series Entomologica, Vol. 9, Dr. W. Junk, B.V- The Hague, Netherlands, 418pp.

دراسات علي صناعة أنفاق أوراق الفول و طفيلياتها في شمال سيناء

فوزى محمد عيد

قسم بحوث المقاومة الحيوية- معهد بحوث وقاية النبات- مركز البحوث الزراعية

تتعدت النسبة المئوية للإصابة بصنعة أنفاق أوراق الفول في مركز العريش من ٢٢,٢١ % إلي ٥٢,٠٨ % في الموسم الأول، ومن ٣٠,٨٣ % إلي ٥٠ % في الموسم الثاني . وعلى الجانب الأخر تراوحت نسب الإصابة في مركز رفح ما بين ١٩,٤٨ % إلي ٤٩,٢٥ % و ١٦,٨٨ % إلي ٤٧,٦٨ % في الموسم الأول والثاني علي الترتيب. وكان متوسط نسب التطفل بطفيليات الهيمينيتروس في مركز العريش ٨٩,١٦ % و ٨٧,٤٣ % في الموسم الأول عام ٢٠٠٢/٢٠٠١ والموسم الثاني عام ٢٠٠٣/٢٠٠٢ علي الترتيب . إما في مركز رفح كان متوسط نسب التطفل يتراوح ما بين ٨٤,٥٨ % و ٨٣,٥٣ % أثناء الموسم الأول والثاني علي الترتيب