

## **SUGAR BEET RESPONSE TO ZINC-APPLICATION UNDER DIFFERENT WATER REGIMES IN NORTHERN DELTA SOILS**

**Abd El-Wahab, S.A. \* and E.A.E. Nemeat Alla\*\***

**\* Soil, Water, and Environment Res. Inst., Agric. Res. Center Egypt**

**\*\* Sugar Crop Res. Inst., Agric. Res. Center Egypt.**

Field experiment was conducted at Sakha Agricultural Res. Station to study the effect of zinc application (2, 4 and 6 kg fed<sup>-1</sup>) under different amounts of irrigation water (3317.6, 2792.6, and 2267.6 m<sup>3</sup> fed<sup>-1</sup>).

The obtained results could be summarized as follows:

- Root yield and sugar yield (ton fed<sup>-1</sup>) were significantly increased as irrigation amounts and zinc application rates were increased.
- Sucrose percentage, impurities (Na, K, αN) in the roots decreased and purity increased with increasing irrigation water amounts, while the application rate of zinc had no significant effect on these characteristics up to 4 kg Zn fed<sup>-1</sup>.
- Root diameter was increased significantly as water amount increased and not affected by application of Zn.
- Root length was increased significantly as water amount decreased and zinc fertilization level increased.
- Root yield, sugar yield, root diameter, sucrose percentage, root length as well as the purity were significantly affected by the interaction between studied treatments irrigation regime and zinc fertilization. The highest values of 44.72, 8.8 ton fed<sup>-1</sup> and 12.31 cm root and sugar yields and root diameter, respectively were obtained by treatment (I<sub>1</sub> x Zn<sub>3</sub>) while the highest values (20.86%, 43.73 cm and 85.13%) for sucrose %, root length, and purity were recorded by treatments (I<sub>3</sub> x Zn<sub>2</sub>), (I<sub>3</sub> x Zn<sub>1</sub>) and (I<sub>1</sub> x Zn<sub>2</sub>), respectively.
- The values of water consumptive use and the moisture uptake from the surface layer were increased with increasing the amount of water applied to the soil and vice versa.
- The values of seasonal KC were 0.82, 0.75, and 0.68 (using modified Penman equation) for treatments I<sub>1</sub>, I<sub>2</sub>, and I<sub>3</sub>, respectively while its values were 0.87, 0.79 and 0.72 (using Radiation equation) for treatment I<sub>1</sub>, I<sub>2</sub>, I<sub>3</sub>, respectively. Using Modified Blaney equation the KC values were 1.3, 1.19 and 1.08 from treatments I<sub>1</sub>, I<sub>2</sub> and I<sub>3</sub>, respectively.
- Water utilization efficiency (W.Ut.E.) for root yield was calculated to be 12.78, 13.79 and 10.10 kg/m<sup>3</sup> for treatments I<sub>1</sub>, I<sub>2</sub> and I<sub>3</sub>, respectively while they were found to be 2.52, 2.72 and 2.07 kg/m<sup>3</sup> for sugar yield, respectively.
- The values of W.U.E. were 17.53, 17.36 and 11.40 kg/m<sup>3</sup> for root yield and 3.45, 3.43 and 2.34 kg /m<sup>2</sup> for sugar yield, with using treatments of I<sub>1</sub>, I<sub>2</sub> and I<sub>3</sub>, respectively.
- Treatment I<sub>3</sub> recorded the highest value of water application efficiency (88.55%). While the lowest value (72.89%) was recorded with I<sub>1</sub> treatment.

**استجابة بنجر السكر للتسميد بالزنك تحت كميات ري مختلفة في أراضي شمال الدلتا**

**\*صلاح على عبدالوهاب - \*\*السيد احمد السيد نعمت الله**

**\* معهد بحوث الأراضي والمياه والبيئة - مركز البحوث الزراعية - الجيزة**

**\*\* معهد المحاصيل السكرية - مركز البحوث الزراعية - الجيزة**

- أجريت تجربة حقلية بمحطة البحوث الزراعية بسخا عام ١٩٩٩/١٩٩٨م لدراسة تأثير كميات مختلفة من ماء الري: ٣، ٦، ٩، ١٢ م<sup>٣</sup>/لقدان، والتسميد بالزنك بمعدلات ٢، ٤، ٦ كيلو جرام زنك/لقدان. ويمكن تلخيص النتائج المتحصل عليها كالآتي:
- ١- زيادة محصول الجذور ومحصول السكر بالطن/فدان معنويا بزيادة كميات الماء المضافة ومعدل التسميد بالزنك.
  - ٢- زيادة نسبة السكر والمواد غير النقية في الجذور ونقص النقاوة بنقص كمية ماء الري المضافة بينما لم يتضح تأثير التسميد بالزنك على هذه المكونات حتى ٤ كجم زنك/لقدان.
  - ٣- زيادة قطر الجذور بالسم معنويا مع زيادة كميات ماء الري بينما لا يوجد تأثيرا معنويا للتسميد بالزنك.
  - ٤- زيادة طول جذر البنجر (بالسم) معنويا بنقص كمية ماء الري وزيادة معدلات التسميد بالزنك.
  - ٥- تأثر محصول الجذور ومحصول السكر وقطر الجذر ونسبة السكروز % وطول الجذر والنقاوة تأثيرا معنويا بالتفاعل بين كمية الماء ومعدل التسميد بالزنك فأعلى قيم (٤٤,٧٢ طن/فدان، ٨,٨ طن/فدان، ١٢,٣١ سم) لمحصول الجذر. ومحصول السكر وقطر الجذر أمكن الحصول عليها باستخدام المعاملة (I<sub>1</sub> X Zn<sub>3</sub>) بينما أعلى القيم (٢٠,٨٦ %، ٤٣,٧٣ سم، ٨٥,١٣ %) للسكروز وطول الجذر والنقاوة % أمكن الحصول عليها بالمعاملات (I<sub>3</sub> X Zn<sub>2</sub>) و (I<sub>1</sub> X Zn<sub>3</sub>) و (I<sub>3</sub> X Zn<sub>1</sub>) على الترتيب.
  - ٦- أتضح أن أعلى قيمة للأستهلاك المائي السنوي ٥٧,٥٨ سم حصل عليها عندما روى النبات بـ ٢٦٢٥ م<sup>٣</sup>/لقدان بينما أقل قيمة للأستهلاك المائي وجدت ٤٧,٨١ سم عند الري بـ ١٥٧٥ م<sup>٣</sup>/لقدان.
  - ٧- وجدت القيم العالية لأمتصاص الرطوبة من الطبقة السطحية كانت ٣٢,٨٤ %، ٣١,٨٣ %، ٣٣,٦١ % للمعاملات I<sub>1</sub>, I<sub>2</sub>, I<sub>3</sub> على الترتيب.
  - ٨- كانت قيم معامل المحصول السنوي ٠,٨٢، ٠,٧٥، ٠,٦٨ للمعاملات I<sub>1</sub>, I<sub>2</sub>, I<sub>3</sub> باستخدام معادلة Modified Penman بينما كانت قيم معامل المحصول السنوي هي ٠,٨٧، ٠,٧٩، ٠,٧٢ للمعاملات I<sub>1</sub>, I<sub>2</sub>, I<sub>3</sub> باستخدام معادلة الإشعاع Radiation وباستخدام معادلة Modified Planey-Cridd أمكن الحصول على قيم معامل المحصول ١,٣، ١,١٩، ١,٠٨ للمعاملات I<sub>1</sub>, I<sub>2</sub>, I<sub>3</sub> على الترتيب.
  - ٩- وجد أن قيمة الكفاءة الإستعمالية للمياه كانت ١٢,٧٨، ١٣,٧٩، ١٠,١٠ كجم/م<sup>٣</sup> لمحصول الجذور وكانت ٢,٥٢، ٢,٧٢، ٢,٠٧ كجم/م<sup>٣</sup> لمحصول السكر مقابل معاملات الماء I<sub>1</sub>, I<sub>2</sub>, I<sub>3</sub> على التوالي.
  - ١٠- وجد أن قيم كفاءة استخدام المياه (W.U.E.) كانت ١٧,٥٣، ١٧,٣٦، ١١,٤ كجم/م<sup>٣</sup> لمحصول الجذور وكانت ٣,٤٥، ٣,٤٣، ٢,٣ كجم/م<sup>٣</sup> لمحصول السكر مقابل معاملات الماء I<sub>1</sub>, I<sub>2</sub>, I<sub>3</sub> على الترتيب.
  - ١١- وفيما يتعلق بكفاءة الري التطبيقية (W.A.E.) وجد أن أعلى القيم (٨٨,٥٥ %) أمكن الحصول عليها بإضافة ٢٢٦٧,٦ م<sup>٣</sup>/فدان.

## INTRODUCTION

Sugar beet (*Beta vulgaris* L.) is considered one of the important economic crops in Egypt and was found to be the most suitable ancillary source of sugar compared to sugar cane. The total amount of sugar produced from sugar beet in 1999 season was 317470 tons and represents about 25% of the total amount (1242587 tons). Egyptian policy aims to increase the amount of sugar from beet to reach 500 000 tons by using proper varieties, irrigation and fertilizations Sobh (1985). Sugar beet is reasonably drought tolerant with yield roughly proportional to total water use (Carter *et al.*, 1980, Miler and Aarstad, 1976, Nicholson *et al.*, 1974, and Winter, 1980).

Water stress in almost cases decrease fresh root weight (Winter, 1980, Sobh, 1985, Gouda *et al.*, 1993, Abd El-Wahab *et al.*, 1996 and Emara, 1996). Foliar application of 0.6 kg Zn/ha increased roots and sugar yields (Stratieva *et al.*, 1990). Besheit *et al.* (1992) reported that soaking sugar beet seeds before sowing in 40 ppm of Zn gave the maximum fresh and dry weights of leaves and roots. Also, zinc application to soil increased dry matter accumulation and sugar content in roots (Sun *et al.*, 1994).

On the other hand, zinc uptake increased as application Zn rate increased but this increase was not affected on root yield (Sofronovskaya, 1998). Sucrose concentration in a fresh weight can be increased due to water stress (Loomis and Worker, 1963; Carter *et al.*, 1980; Abd El-Wahab *et al.*, 1996).

The effects of irrigation on purity not well understood, Bauer *et al.*, 1975; Reichman *et al.*, 1977; Sobh, 1985; Attia and Sultan, 1987 and Abdel-Wahab *et al.*, 1996.

Monthly and seasonal consumptive water use were increased as the irrigation water amounts increased (Doorenbos *et al.*, 1979; Sobh 1985 and Abdel Wahab *et al.*, 1996).

The objective of this research is to determine the influence of zinc fertilization and the amount of seasonal irrigation water on sugar beet crop and its water consumptive use in Northern Delta soils.

## MATERIALS AND METHODS

The effect of zinc fertilization and irrigation treatments on sugar beet during growing season 1998/1999 was tested in a split plot design three levels of seasonal irrigation water amounts used the main plots  $I_1 = 3317.6$ ,  $I_2 = 2792.6$ , and  $I_3 = 2267.6$  m<sup>3</sup>/fed. these amounts including planting irrigation and effective rain off. Three levels of zinc fertilizer ( $Zn_1 = 2$  kg fed<sup>-1</sup>,  $Zn_2 = 4$  kg Zn fed<sup>-1</sup>, and  $Zn_3 = 6$  kg Zn fed<sup>-1</sup>) were added to the sub plots by soil application before the second irrigation. Both N-fertilizer (80 kg N/fed.) as urea (46.5% N) and K fertilizer in the form of K-sulfate (48% K<sub>2</sub>O) at rate of 48 kg fed<sup>-1</sup> K<sub>2</sub>O were applied for all plots in two equal doses at the thinning time and before the second irrigation P-fertilizer as superphosphate (15.5% P<sub>2</sub>O<sub>5</sub>) was added during tillage operation with the rate of 15 kg P<sub>2</sub>O<sub>5</sub>/fed. Sugar beet (Cawamera) variety was sown on 15 November. Plot area was 20 m<sup>2</sup> (4 x 5 m).

The soil characteristics of the experimental site (were determined according to Black (1965) and presented in Table (1).

The climatological data of the studied area during the growing season are recorded in (Table 2). The plants were harvested on May 31 and the following parameters were recorded:

1. Sugar beet yield (root and sugar ton/fed.).
2. Yield component (root length and diameter).
3. Yield quality [sucrose %, impurities (K, Na,  $\alpha$ N) and purity] was determined in Delta Sugar Company Limited Laboratories at Kafr El-Sheikh Governorate.

Soil water relations were determined as follows:

- A) Water consumptive use (Cu) in each irrigation was calculated according to (Hansen *et al.*, 1979). as follows:

$$Cu = \sum \frac{Pw_2 - Pw_1 \times BDi \times Di}{100}$$

**Where:**

- Cu = Water consumptive use (cm) for the effective root zone (0-60 cm).  
i = Number of soil layers.  
 $Pw_2 - Pw_1$  = Percentage of soil moisture content 48 hours after irrigation and before irrigation for the specified soil layer.  
BDi = Bulk density for specified layer.  
Di = Depth of layer (15 cm).

- B) The amounts of water applied to each plot was measured using small siphon according to FAO (1974).**

**C) Water efficiencies:**

1. Water utilization efficiency (W.Ut.E) was calculated according to Doorenbos and Pruitt (1975) as follows:

$$W.Ut.E = \frac{\text{Root or sugar yield in kg fed}^{-1}}{\text{Total water applied in m}^3 \text{ fed}^{-1}}$$

2. Water use efficiency (W.U.E.):

$$W.U.E. = \frac{\text{Root or sugar yield in kg fed}^{-1}}{\text{Actual evapotranspiration m}^3 \text{ fed}^{-1}}$$

3. Water application efficiency (W.A.E.):

$$W.A.E. = \frac{\text{Total water stored in root zone}}{\text{Total water applied}} \times 100$$

**D) Crop coefficient (KC):**

It was calculated according to Doorenbos and Pruitt (1975) as follows:

$$KC = \frac{\text{Actual evapotranspiration ETA}}{\text{Potential evapotranspiration ETP}}$$

ETP was calculated by using, modified Penman, Radiation and Modified Blany-Criddle equations.

Statistical analysis was carried out according to Snedecor and Cochran (1967), and treatment means were compared by least significant differences (L.S.D.) at the levels of 1% and 5% probability.

**Table (1): Mechanical and chemical analysis of experimental soil (0-60 cm)\*.**

Particle size distribution			Texture class	E.C dSm <sup>-1</sup>	pH 1: 2.5	Available nutrients ppm			
Sand %	Silt %	Clay%				N	P	K	Zn
17.53	30.34	49.95	Clayey	1.32	7.86	58.83	9.84	416.0	1.08

\* Mechanical analysis was determined according to the International Method Piper, (1950). Available N, and K according to Black (1965) Available P according to Olsen *et al.* (1954) Available Zinc was determined according to Lindsay and Norvell (1978).

**Table (2): Meteorological data of Sakha Agriculture Research Station 1998/1999.**

Month	Temperature, °C		Relative humidity %		Wind. sp., km/hr	Rain, mm
	Max.	Min.	Max.	Min.		
Nov. 1998	26.0	12.0	75.4	42.5	73.0	-
Dec. 1998	21.30	8.2	72.20	41.9	109.9	3.6
Jan. 1999	19.33	8.8	69.83	46.69	60.90	7.8
Feb. 1999	17.73	6.73	67.46	46.00	98.11	11.8
March 1999	21.13	10.71	67.84	44.90	109.23	-
April 1999	25.43	9.22	73.57	45.77	124.30	5.1
May 1999	29.27	15.5	70.06	41.42	141.13	-

## RESULTS AND DISCUSSION

### 1) Root yield (ton fed<sup>-1</sup>):

Data in Table (3) indicated that root yield was significantly increased with increasing the amounts of irrigation water and the maximum yield (42.39 ton fed<sup>-1</sup>) was obtained when 3317.6 m<sup>3</sup> water per feddan was added to the soil. The magnitude of the increase was 85.1% as compared to the treatment received 2267.6 m<sup>3</sup> water per feddan that gave the lowest value of root yield (22.90 ton fed<sup>-1</sup>). This may be due to the direct effect of the availability of soil water in the effective root zone. These results are in agreement with those obtained by Sobh (1985) Gaber *et al.* (1986); Cucci and Caro (1986); Emara (1990, 1996) and Abd El-Wahab *et al.* (1996). With respect to zinc effect on the root yield. Data in Table (3) showed that zinc fertilization treatments gave significant synergistic effect on root yield. The maximum root yield 35.82 ton

**Table (3): Effect of irrigation and zinc fertilization on sugar beet characteristics.**

Treatment	Root quality						Root component		Root yield ton fed <sup>-1</sup>	Sugar yield ton fed <sup>-1</sup>
	Sucrose pol. %	Impurities (mg/100 g).				Purity %	Root diameter cm	Root length cm		
		K	Na	αN	$\frac{K+Na}{N}$					
A. irrigation m <sup>3</sup> fed <sup>-1</sup>										
I <sub>1</sub> =3317.6	19.69	6.38	1.52	2.21	3.54	84.55	12.16	32.57	42.39	8.31
I <sub>2</sub> =2792.6	19.74	7.05	1.88	2.72	3.34	82.03	12.14	41.40	38.50	7.60
I <sub>3</sub> =2267.6	20.50	6.66	1.69	2.73	3.08	82.54	9.71	43.47	22.90	4.70
L.S.D.	**	**	*	*	*	**	**	**	**	**
B. Zinc fer. kg fed <sup>-1</sup>										
Zn <sub>1</sub> =2	20.04	6.67	1.69	2.49	3.37	82.93	11.46	38.34	32.76	6.57
Zn <sub>2</sub> = 4	20.05	6.80	1.77	2.56	3.34	83.17	11.39	39.07	35.82	7.18
Zn <sub>3</sub> = 6	19.80	6.61	1.63	2.62	3.25	83.03	11.17	40.03	35.22	6.97
L.S.D.	**	**	**	n.s	n.s	n.s	n.s	**	**	**

fed<sup>-1</sup>, using 4 kg Zn fed<sup>-1</sup> while the lowest yield was recorded with 2 kg Zn fed<sup>-1</sup>. (32.76 ton fed<sup>-1</sup>) which achieved an increment of 9.4%.

These results are in harmony with those obtained by Stratieva *et al.* (1990); Besheit *et al.* (1992) and Sun *et al.* (1994).

## **2. Gross sugar yield (ton fed<sup>-1</sup>):**

Sugar yield per unit area is the main goal of sowing any sugar crop and it is the sum product of sugar extractable % and root yield per unit area. Data in Table (3) revealed that decreasing the amounts of irrigation water applied to the soil ( 3317.6 to 2267.6 m<sup>3</sup> water fed<sup>-1</sup>) resulted in 76% decrease in gross sugar yield may be due soil moisture stress as shown before. These results supported the results obtained by Carter *et al.* (1980); Roberts *et al.* (1980); Sobh (1985); Cucci and Caro (1986) and Abd El-Wahab (1996).

Dealing with the effect of zinc fertilization on gross sugar yield, data clarify that gross sugar yield was significantly increased with increasing zinc levels applied to the soil up to 4 kg Zn fed<sup>-1</sup> which gave the maximum yield 7.18 ton fed<sup>-1</sup> compared with 6.57 and 6.97 ton fed<sup>-1</sup> given by of 2 and 6 kg Zn fed<sup>-1</sup>, respectively. These results are in agreement with those obtained by Sun *et al.* (1994).

The impact of the interaction between irrigation and zinc levels on sugar beet yield (root and sugar yields) was clear. The optimum yield of root and sugar yields (44.14 and 8.59 ton fed<sup>-1</sup>), respectively were obtained with application of 3317.6 m<sup>3</sup> water fed<sup>-1</sup> and adding 4 kg Zn fed<sup>-1</sup>.

## **3. Sucrose percentage:**

Data in Table (3) demonstrate that there was a significant negative relation between sucrose % and the amount of water added to the soil consequently the maximum value 20.54% was given by 2267.6 m<sup>3</sup> water fed<sup>-1</sup> (I<sub>3</sub>). These results stand in the same line with those obtained by Carter *et al.* (1980), Roberts *et al.* (1980); Abd El-Wahab *et al.* (1996). In this regard Loomis and Hadock (1967) attributed the increase in sucrose percentage with increasing soil moisture stress to slower accumulations of dry matter and more rapid accumulation of sucrose.

As concerns zinc fertilization effect, data showed that sucrose % was not significantly affected by 2 or 4 kg zinc while it significantly affected by 6 kg Zn fed<sup>-1</sup>.

## **4. Impurities (Na, K and αN):**

Impurities in the roots were increased and the purity was decreased with decreasing irrigation water amounts applied to the soil (Table, 3). The highest values of the impurity and the lowest purity value were detected with 27926 m<sup>3</sup> water fed<sup>-1</sup> (I<sub>2</sub>). These results are in accordance with the findings of Winter (1980).

On the other hand, purity was not significantly affected by Zn, while there was a significant effect on alkalinity of sugar juice

The interaction between irrigation and zinc not significantly affected on αN and alkalinity while its effect was significant for Na and K contents and juice purity. The maximum purity value 85.13% was achieved by the combination between application of 3317.6 m<sup>3</sup> water (I<sub>1</sub>) and 4 kg Zn fed<sup>-1</sup>.

**5) Root diameter (cm):**

With respect to root diameter, data in Table (3) showed that decreasing the amounts of irrigation water added to the soil significantly decreased root diameter especially at the irrigation water level of 2267.6 m<sup>3</sup> water fed<sup>-1</sup> (I<sub>3</sub>) which gave the lowest diameter of 9.71 cm, the diameter 12.16 cm was recorded with 33176 m<sup>3</sup> water fed<sup>-1</sup> (I<sub>1</sub>). This may be due to the availability of soil water which had the direct effect on the growth of root. In this regard, Sorour (1995) attributed the reduction in root diameter at low soil moisture level to decreasing leaf area index which might decrease light interception and in turn decrease dry matter accumulation and root dimensions. These results are similar to those obtained by Attia and Sultan (1987); Emara (1990, 1996) and Abd El-Wahab (1996) For zinc effect where there was no significant effect on root diameter as zinc fertilizer applied to the soil.

The interaction between irrigation and zinc treatments recorded significant effect on root diameter and the best value of interaction was obtained by combination of 33176.6 m<sup>3</sup> water fed<sup>-1</sup> (I<sub>1</sub>) and 6 kg Zn fed<sup>-1</sup> (Zn<sub>3</sub>) (Table 4).

**Table (4): Sugar beet characteristics as affected by irrigation and zinc fertilization interaction.**

Treatment		Root quality						Root component		Root yield ton fed <sup>-1</sup>	Sugar yield ton fed <sup>-1</sup>
Water regime	Zn application	Sucrose pol. %	Impurities (mg/100 g).				Purity %	Root diameter cm	Root length cm		
			K	Na	αN	$\frac{K+Na}{N}$					
I <sub>1</sub>	Zn <sub>1</sub>	19.67	6.42	1.64	2.15	3.71	83.60	11.93	30.93	38.31	7.54
	Zn <sub>2</sub>	19.47	6.52	1.49	2.21	3.53	85.13	12.25	31.88	44.14	8.59
	Zn <sub>3</sub>	19.67	6.19	1.42	2.27	3.38	84.93	12.31	34.90	44.72	8.80
I <sub>2</sub>	Zn <sub>1</sub>	20.04	6.98	1.84	2.67	3.33	82.58	12.28	40.38	36.63	7.34
	Zn <sub>2</sub>	19.82	7.02	2.04	2.68	3.38	81.43	12.10	41.79	39.29	7.79
	Zn <sub>3</sub>	19.37	7.15	1.76	2.80	3.31	82.10	12.05	42.05	39.59	7.67
I <sub>3</sub>	Zn <sub>1</sub>	20.41	6.60	1.60	2.64	3.08	82.63	10.17	43.73	23.34	4.76
	Zn <sub>2</sub>	20.86	6.86	1.78	2.78	3.10	82.95	9.83	43.54	24.02	5.01
	Zn <sub>3</sub>	20.35	6.51	1.71	2.78	3.05	82.05	9.13	43.15	21.35	4.34
L.S.D. 0.01		**	**	**	n.s	n.s	**	**	**	**	**

**6. Root length (cm):**

Data in Table (3) revealed that decreasing irrigation water amounts significantly promoted increasing the root length to reach maximum value 43.47 cm at 2267.6 m<sup>3</sup> water fed<sup>-1</sup> level.

The results showed also that the roots grow longer under water stress than that excessive water status. The water stress enhanced deep rooting. These results are in line with those obtained by Winter (1980); Emara (1990-1996) and Abd El-Wahab *et al.* (1996).

With relation to the effect of zinc on root length, data clarify that application of zinc fertilizer to the soil gave significant pronounced positive effect on root length recording maximum value 40.03 cm when the zinc fertilizer applied at the rate of 6 kg Zn fed<sup>-1</sup>.

Also, the interaction between irrigation and zinc achieved significant synergistic effect and continued to increase the root length to reach the maximum value 43.54 cm by using 2267.6 m<sup>3</sup> water fed<sup>-1</sup> level and 4 kg Zn fed<sup>-1</sup> (Table 4).

**7. Consumptive water use (Cu):**

Data in Table (5) shows that both monthly and seasonal Cu values were increased with increasing the amounts of irrigation water applied to the soil. For monthly values the rate of Cu were gradually increased with crop development and reached its peaks in April and then reduced during ripening stage (May).

**Table (5): Effect of irrigation regimes on monthly or seasonal consumptive use and water uptake pattern for sugar beet plants**

Irrigation treatment	Layer depth, cm	Monthly and seasonal consumptive use cm							Total	W.U.P pattern
		Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May		
I <sub>1</sub>	0-15	0.736	2.073	2.062	1.847	2.700	5.451	4.038	18.907	32.84
	0-30	0.482	1.359	1.653	1.860	2.284	5.049	3.303	15.990	27.77
	0-45	0.213	0.601	1.026	1.459	1.591	4.371	3.124	12.385	21.51
	0-60	0.092	0.259	0.700	1.185	1.319	3.607	3.136	10.298	17.88
Total		1.523	4.292	5.441	6.351	7.894	18.478	13.601	57.58	
I <sub>2</sub>	0-15	0.566	1.596	1.858	2.006	2.298	4.506	4.00	16.830	31.83
	0-30	0.403	1.136	1.623	2.074	2.116	4.595	3.185	15.132	28.62
	0-45	0.208	0.586	0.833	1.063	2.090	4.051	3.269	12.100	22.88
	0-60	0.082	0.230	0.323	0.407	1.673	2.881	3.218	8.814	16.67
Total		1.259	3.548	4.637	5.550	8.177	16.033	13.672	52.88	
I <sub>3</sub>	0-15	0.549	1.548	1.802	1.945	2.229	4.00	4.00	16.07	33.61
	0-30	0.371	1.045	1.259	1.409	2.053	3.803	3.088	13.028	27.25
	0-45	0.132	0.372	0.617	0.862	2.027	3.683	3.024	10.717	22.42
	0-60	0.035	0.101	0.242	0.394	1.623	2.785	2.808	7.988	16.71
Total		1.087	3.066	3.92	4.61	7.932	14.271	12.92	47.81	

Sowing irrigation = 520.4 m<sup>3</sup> fed<sup>-1</sup>  
 Rain off = 172.2 m<sup>3</sup> fed<sup>-1</sup>

In relation to seasonal Cu of sugar beet plants the maximum value 57.58 cm which corresponding to 0.29 cm/day was obtained when 3317.6 m<sup>3</sup> water fed<sup>-1</sup> (or water regime I<sub>1</sub>) was applied, whereas the minimum one (47.81 cm) that corresponding to 0.24 cm/day was recorded at 2267.6 m<sup>3</sup> water fed<sup>-1</sup> level (or at water regime I<sub>3</sub>). These results are in harmony with those obtained by Doorenbos *et al.* (1979); Sobh (1985); Abd El-Wahab *et al.* (1996) who stated that the seasonal values may be differ due to climate, availability of soil water and length of the total growing period.

**8. Water uptake patterns of sugar beet:**

The prediction of the degree of root distribution among different depths of the effective root zone can be realized by using the parameter of water uptake patterns. Thus, data in Table (5) revealed that the uptake of soil water was decreased with the soil depth and the greater uptake values were 32.8, 31.8 and 33.62% for the surface layer of the water treatments I<sub>1</sub>, I<sub>2</sub> and



$I_3$ , respectively. While less water was extracted from the subsurface layers. The relatively higher water uptake from the upper layers compared to the deepest ones was attributed to the concentrated roots in the upper layer (Ibrahim *et al.*, 1988).

**9. Crop coefficient KC of sugar beet:**

Data in Table (6) shows that the seasonal crop coefficient values (KC) increased with increasing  $C_u$  value. The theoretical values of evapotranspiration ( $ET_0$ ) was calculated to be 705.23 mm from Modified Penman equation (with KC values of 0.82, 0.75 and 0.68), 664.64 mm from Radiation equation (with KC values 0.87, 0.79, and 0.72) and 442.93 mm from Modified Blaney-Criddle equation (with KC values of 1.30, 1.19 and 1.08) for water regimes  $I_1$ ,  $I_2$ ,  $I_3$ , respectively.

**Table (6): Effect of different water regimes on consumptive use indices of sugar beet**

Water treat.	$C_u$ $m^3$ fed <sup>-1</sup>	Added water $m^3$ fed <sup>-1</sup>	KC* $ET_0 = 705.23$ mm	KC** $ET_0 = 664.64$ mm	KC** $ET_0 = 442.93$ mm	Root yield kg fed <sup>-1</sup>	Sugar yield kg fed <sup>-1</sup>	W.Ut.E $kg/m^3$		W.U.E. $kg/m^3$		W.A.E. %
								Root	Sugar	Root	Sugar	
$I_1$	2418.4	3317.6	0.82	0.87	1.30	42390	8347	12.78	2.52	17.53	3.45	72.89
$I_2$	2218.0	2792.6	0.75	0.79	1.19	38500	7600	13.79	2.72	17.36	3.43	79.42
$I_3$	2008.0	2267.6	0.68	0.72	1.08	22900	4704	10.10	2.07	11.40	2.34	88.55

\* Calculated evapotranspiration values from Modified Penman Equation.

\*\* Calculated evapotranspiration values from Radiation Equation.

\*\*\* calculated evapotranspiration values from Modified Blaney-Criddle-equation.

The data also indicate that  $ET_0$  values calculated using Blaney-Criddle equation is lower than the actual consumptive use of water ( $C_u$ ). However, Modified Penman or Radiation equations agree well with the value of  $C_u$  determined under the present study.

**10. Water efficiencies:**

The optimum water management is achieved by obtaining the greatest consumptive use of water with increasing the grain yield per unit area. Thus with respect to the effect of different irrigation treatments on water use efficiency (W.U.E.), water utilization efficiency (W.Ut.E) and water application efficiency (W.A.E.) are shown in (Table 6). The data showed that the irrigation regime ( $I_2$ ) seemed to be superior in increasing the value of W.Ut.E for both root and sugar yields (13.79 and 2.72  $kg/m^3$ , respectively followed by  $I_1$  and  $I_3$ . The highest values of W.U.E. for root and sugar (17.53 and 3.45  $kg/m^3$ , respectively, were achieved with  $I_1$  (3317.6  $m^3$  fed<sup>-1</sup>) followed by  $I_2$  (2792.6  $m^3$ /fed<sup>-1</sup>) treatment

This trend may be due to the severe reduction in yield (root and sugar) as a result of soil moisture stress. In relation to water application efficiency (W.A.E.) data indicated that its values were increased as the amount of irrigation water decreased and reached to the maximum value (88.55%) with treatment ( $I_3$ ). It can be said that as the total irrigation water delivered increased, the application efficiency decreased and vice versa. These results are supported those obtained by Abd El-Wahab (1996).

However, I<sub>2</sub> treatment gave the suitable values of W.Ut.E and W.U.E. indicating that the water regime which utilize reasonable amount of water with suitable Zn application resulted in a great yield, W.Ut.E. and W.U.E.

It could be concluded that for sugar beet production the suitable water regime (3317.6 and 2792.6 m<sup>3</sup> fed<sup>-1</sup>) and zinc application (4 kg Zn fed<sup>-1</sup>) which should be adopted in clayey soils in Northern Delta.

## REFERENCES

- Abd El-Wahab, S.A.; A.A. Amer; M.I. El-Shahawy and M.M. Sobh (1996). Effect of different irrigation amount and potassium fertilizer rates on yield and quality of sugar beet and water efficiencies. Mansoura Univ. J. Agric. Sci. 21: 4687-4699.
- Attia, A.N. and M.S. Sultan (1987). Response of some sugar beet varieties (*Beta vulgaris*, L.) to irrigation intervals and harvesting dates. Conf. of Agric. Sci. on Food Deficiency Over Coming Through Autonomous Efforts in Egypt. vol. 5, 22-29. June. Fac. of Agric. Mansoura Univ.
- Bauer, A.T.; D.M. Heimbuch, Cassel and L. Zimmerman (1975). Production potential of sugar beet under irrigation in the West Oakes Irrigation District. Bulletin, North Dakota Agric. Exper. Station (No. 498). p. 4.
- Besheit, S.Y.; F.M. Abd El-Naeem; M. El-Houssiny and Z.R. Moustafa (1992). Effect of micronutrients on biochemical changes, yield, and quality of sugar beet. Egyptian J. Agric. Res. 70: 1227-1242.
- Black, C.A. (1965). Methods of Soil Analysis. Amer. Soc. of Agro. Inc. Madison, Wisconsin, U.S.A.
- Carter, J.N.; M.E. Jensen and D.J. Traveller (1980). Effect of mid-to late-season water stress on sugar beet growth and yield. Agron. J. 72: 806-815.
- Cucci, G. and A. de Caro (1986). Impact of irrigation on productivity of autumn-sown sugar beet in Italy. Institute de Agron. colivaziani erbacee dell de Bari, Italy. Irrigation drenaggio, 33: 29-32 [C.F. Irrigation and Drainage Abst. 13: 1332 (1987)].
- Doorenbos, J. and W.O. Pruitt (1975). Crop water requirements; Irrigation and drainage paper., 24, FAO, Rome.
- Doorenbos, J.; A.H. Kassam; C.L.M. Bentvelsen and V. Brancheid (1979). Yield response to water FAO irrigation and drainage paper, No. 33. Rome.
- Emara, T.K. (1990). Effect of irrigation intervals, growth regulators and NK fertilizers on yield and quality of sugar beet. M.Sc. Thesis Fac. of Agric., Mansoura Univ.
- Emara, T.K. (1996). Studies on the effect of some cultural practices on water parameter of sugar beet in North Delta Region. Ph.D. Thesis. Fac. of Agric. Mansoura Univ. Egypt.
- FAO (1974). Surface Irrigation FAO Agricultural Development. Paper NO. 95 FAO. Land and Water Development. No. 3. FAO. Rome, 1974.
- Gaber, A.A.; M.M. El-Banna and A.H. Nour (1986). Effect of irrigation intervals and nitrogen levels on yield and sugar beet. Alex. Sci. Exch. Vol. 7 No. 4.

- Gouda, M.; A.A. Shiha and M.M. Sobh (1993). Effect of water management and N-fertilization on sugar beet. Egypt. J. Appl. Sci., 8: 144-159.
- Hansen, V.W.; O.W. Israelsen and Q.E. Stringham (1979). Irrigation Principles and Practices. Edition, John Wiley and Sons., 1979.
- Ibrahim, M.A.M.; S.A. El-Gohary and M.A. Rezk (1988). Preliminary studies towards effective water management in Kafr El-Sheikh Governorate. Alex. Eng. J. Alex. Univ., 23: 29-36.
- Lindsay, W.L. and W.A. Norvell (1978). Development of a DTPA Test for Zn, Fe, Mn and Cu. Soil Sci. Soc. Amer. J. 42: 427.
- Loomis, R.S. and J.L. Haddock (1967). Sugar oil, and fiber-crops. 1-Sugar beet in R.M. Hagan *et al.* (ed.) irrigation of Agricultural Lands. Agronomy, 11: 640-648.
- Loomis, R.S. and G.F. Worker, Jr. (1963). Response of the sugar beet to low soil moisture at two levels of nitrogen nutrition-Agron. J. 55: 509-515.
- Miller, D.E. and J.S. Aarstad (1976). Yield and sugar content of of sugar beets as affected by deficit high-frequency irrigation. Agron. J. 68: 231-234.
- Nicholson, M.R.; T.K. Breab; R.E. Danielson and R.A. Young (1974). Yield and economic implications of sugar beet production as influenced by irrigation and nitrogen fertilizer Amr. Soc. Sugar Beet Tel.
- Olsen, S.R.; O.V. Cole; F.S. Watanabe and L.A. Dean (1954). Estimation of available phosphorus in soils by extraction with sodium bicarbonate. U.S.D. Agric. Circ. 939.
- Piper, C.S. (1950). Soil and Plant Analysis. Inter-Science Publisher, New York.
- Reichman, G.A.; E.J. Doering; L.C. Benze and R.E. Follett (1977). Effects of water-table depth and irrigation on sugar beet yield and quality. J. Am. Soc. Sugar Beet Technol. 19: 275-287.
- Roberts, S.; J.E. Middleton; A.W. Richards; W.H. Weaver and L.F. Hall (1980). Sugar beet production under center pivot irrigation with different rates of nitrogen. Bulletin, College of Agric. Res. Cent. Washington State Univ. No. 884: 5 pp.
- Safronovskaya, G.M. (1998). Effect of zinc fertilizers on the productivity and changes in Sod-podzolic loamy soil. Pochvovedenie-i-Hgrokimiya 30: 166-171.
- Snedecor, G.W. and W.G. Cochran (1967). Statistical Methods. Six Edition. Iowa State Univ. pp. 593.
- Sobh, M.M. (1985). Some factors affecting the consumptive use of water by sugar beet (*Beta vulgaris* L.). Ph.D. Thesis, Zagazig Univ. Egypt.
- Sorour, S.Gh.R. (1995). Response of sugar beet to irrigation regime and plant population. Egypt, J. Appl. Sci., 10: 648-668.
- Stratieva, S.; B. Sedlarska and D. Stoyanov (1990). Effect of zinc and boron on sugar beet grown on a leached smonitza chernozem soil. Pochvoznanie-i-Agrokimiya 25: 9-14.
- Sun, S.J.; F.S. Li; Y. Wan and G.C. Zheng (1994). Effects of zinc and potassium on dry matter accumulation of sugar beet in mid-late growing season. China-sugar beet, No. 4: 26-29.
- Winter, S.R. (1980). Suitability of sugar beet for irrigation in a semi-arid climate Agron. J. 72: 118-123.

استجابة بنجر السكر للتسميد بالزنك تحت كميات ري مختلفة في أراضى شمال الدلتا  
\*صلاح على عبدالوهاب - \*\*السيد احمد السيد نعمت الله  
\* معهد بحوث الأراضى والمياه والبيئة - مركز البحوث الزراعية - الجيزة  
\*\* معهد المحاصيل السكرية - مركز البحوث الزراعية - الجيزة

- أجريت تجربة حقلية بمحطة البحوث الزراعية بسخا عام ١٩٩٨/١٩٩٩م لدراسة تأثير كميات مختلفة من ماء الري: ٣٣١٧,٦ ، ٢٧٩٢,٦ ، ٢٢٦٧,٦ م<sup>٣</sup> للفدان ، والتسميد بالزنك بمعدلات ٢ ، ٤ ، ٦ كيلو جرام زنك/الفدان. ويمكن تلخيص النتائج المتحصل عليها كالاتى:
- ١- زيادة محصول الجذور ومحصول السكر بالطن/فدان معنويا بزيادة كميات الماء المضافة ومعدل التسميد بالزنك.
  - ٢- زيادة نسبة السكر والمواد غير النقية فى الجذور ونقص النقاوة بنقص كمية ماء الري المضافة بينما لم يتضح تأثير التسميد بالزنك على هذه المكونات حتى ٤كجم زنك/الفدان.
  - ٣- زيادة قطر الجذور بالسهم معنويا مع زيادة كميات ماء الري بينما لا يوجد تأثيرا معنويا للتسميد بالزنك.
  - ٤- زيادة طول جذر البنجر (بالسم) معنويا بنقص كمية ماء الري وزيادة معدلات التسميد بالزنك.
  - ٥- تأثير محصول الجذور ومحصول السكر وقطر الجذر ونسبة السكروز% وطول الجذر والنقاوة تأثيرا معنويا بالتفاعل بين كمية الماء ومعدل التسميد بالزنك فأعلى قيم (٤٤,٧٢ طن/فدان ، ٨,٨ طن/فدان ، ١٢,٣١ سم) لمحصول الجذر. ومحصول السكر وقطر الجذر أمكن الحصول عليها باستخدام المعاملة (I<sub>1</sub> X Zn<sub>3</sub>) بينما أعلى القيم (٢٠,٨٦% ، ٤٣,٧٣ سم ، ٨٥,١٣%) للسكروز وطول الجذر والنقاوة% أمكن الحصول عليها بالمعاملات (I<sub>3</sub> X Zn<sub>2</sub>) و (I<sub>3</sub> X Zn<sub>1</sub>) و (I<sub>1</sub> X Zn<sub>3</sub>) على الترتيب.
  - ٦- أتضح أن أعلى قيمة للأستهلاك المائى السنوى ٥٧,٥٨ سم حصل عليها عندما روى النبات ب ٢٦٢٥م<sup>٣</sup>/الفدان بينما أقل قيمة للأستهلاك المائى وجدت ٤٧,٨١ سم عند الري ب ١٥٧٥م<sup>٣</sup>/الفدان.
  - ٧- وجدت القيم العالية لأمتصاص الرطوبة من الطبقة السطحية كانت ٣٢,٨٤% ، ٣١,٨٣% ، ٣٣,٦١% للمعاملات I<sub>1</sub> , I<sub>2</sub> , I<sub>3</sub> على الترتيب.
  - ٨- كانت قيم معامل المحصول السنوى ٠,٨٢ ، ٠,٧٥ ، ٠,٦٨ للمعاملات I<sub>1</sub> , I<sub>2</sub> , I<sub>3</sub> باستخدام معادلة Modified Penman بينما كانت قيم معامل المحصول السنوى هى ٠,٨٧ ، ٠,٧٩ ، ٠,٧٢ للمعاملات I<sub>1</sub> , I<sub>2</sub> , I<sub>3</sub> باستخدام معادلة الإشعاع Radiation وباستخدام معادلة Modified Planey-Cridd أمكن الحصول على قيم معامل المحصول ١,٣ ، ١,١٩ ، ١,٠٨ للمعاملات I<sub>1</sub> , I<sub>2</sub> , I<sub>3</sub> على الترتيب.
  - ٩- وجد أن قيمة الكفاءة الإستعمالية للمياه كانت ١٢,٧٨ ، ١٣,٧٩ ، ١٠,١٠ كجم/م<sup>٣</sup> لمحصول الجذور وكانت ٢,٥٢ ، ٢,٧٢ ، ٢,٠٧ كجم/م<sup>٣</sup> لمحصول السكر مقابل معاملات الماء I<sub>1</sub> , I<sub>2</sub> , I<sub>3</sub> على التوالى.
  - ١٠- وجد أن قيم كفاءة استخدام المياه (W.U.E.) كانت ١٧,٥٣ ، ١٧,٣٦ ، ١١,٤ كجم/م<sup>٣</sup> لمحصول الجذور وكانت ٣,٤٥ ، ٣,٤٣ ، ٢,٣ كجم/م<sup>٣</sup> لمحصول السكر مقابل معاملات الماء I<sub>1</sub> , I<sub>2</sub> , I<sub>3</sub> على الترتيب.
  - ١١- وفيما يتعلق بكفاءة الري التطبيقية (W.A.E.) وجد أن أعلى القيم (٨٨,٥٥%) أمكن الحصول عليها بإضافة ٢٢٦٧,٦ م<sup>٣</sup>/فدان.