

EFFECT OF SOWING DATES AND NPK LEVELS ON ACTIVE SUBSTANCES IN SEEDS OF ISABGOL PLANTS (*Plantago ovata* Forsk, L.).

Massoud. Hekmat, Y.A.; H.A. Ahmed and Seham M.A. El-Gamal
Veget. and Flori. Dept., Fac. of Agric., Mansoura Univ.

ABSTRACT

Two field experiments were carried out in the Experimental Station and Laboratory of the Vegetables and Floriculture Dept., Fac. of Agric., Mansoura Univ., during the winter seasons of 1995/96 and 1996/97. The investigation aimed to study the effect of different sowing dates and NPK fertilization levels, as well as their interactions on active substances extracted from Isabgol plant (mucilage contents).

Four sowing dates and six NPK levels were studied. The main findings in this investigation can be summarized as follows:-

1. Plants grown in the first sowing date (Nov. 1st week) produced the highest content of mucilage as g / 5 g seeds or as percentage of mucilage per plant.
2. The addition of NPK fertilization significantly increased the mucilage seed content and percentage per plant. The highest content and percentage were of plants treated with the fourth level of NPK (178.0 g/3m²).
3. Concerning effect of interaction between sowing date and NPK levels on mucilage content, the highest content and percentage per plant were produced from plants grown in the first sowing date and treated with the fourth level of NPK.

INTRODUCTION

The genus *Plantago* comprises 200 species, one of them is Isabgol plant (*Plantago ovata* Forsk, L.). Such plant is characterized as an annual herb and produces a heavy yield of seeds that makes it the main source of such seeds.

Since early time, it was proved from several studies the importance of Isabgol seeds in different pharmaceutical purposes. The seed husks take a lot of attention in this respect as a source forming tasteless mucilaginous substances (Wallis, 1967; Russell, 1975 and Trease and Evans, 1978).

The main constituent of such substances in mucilage compound. Anderson (1987) reported the effective use of mucilaginous husk for lowering blood cholesterol level. As for seeds, Bloss (1982) stated their use as a laxative and Chiej (1988) found that seeds can be used as cosmetic products, e.g. face masks to makes the skin soft. More recent, Nordgaard *et al.* (1996) pointed to the role dietary fiber from Isabgol seeds as effective in treatment of colonic cancer.

In Egypt, Isabgol was found widely in the Nile Delta and along the Mediterranean coastal strip from El-Salam to Rafa. In addition, it was found in the Isthmic desert, i.e. El-Tih and regions of North Wadi Tumilat and Sinai proper, i.e. South of El-Tih desert (Tackholm, 1974).

Factors affecting the production of such medical substance did not receive the considerable attention. A very limit studies were carried out, such as those of Khater *et al.* (1994), who found that the highest yield of seeds mucilage was obtained from Isabgol plants supplied.

Therefore, the objectives of the present research were a trial in this direction. It was designed to test the effect of four sowing date and six NPK levels either solely or in combinations on certain characteristics represent mucilage content.

MATERIALS AND METHODS

Isabgol plant seeds (*Plantago ovata* Forsk [L]) Fam. Plantagonaceae, sourced from Muggenburg Firm, Hamburg, Germany were sown seed-bed of uniform fine tilth located in the Experimental Station of Agriculture College, Mansoura University, sowing dates tested in 1995/96 and 1996/97 seasons were four consequent dates, the first week of November, the third week of November, the first week of December and the third week of December. Also during the growth season, six NPK levels were applied at a ratio of 2:1:1.5 and its duplicates as indicated in Table (1).

Table 1. Fertilization treatments studied for NPK ratio and rates in 1995/96 and 1996/97 seasons.

NPK Treatments	NPK ratio	NPK rates (g/3m ²)			
		NH ₄ NO ₃	Ca ₃ (P ₂ O ₅) ₂	K ₂ SO ₄	Total
N ₁ P ₁ K ₁	0 : 0 : 0	0.0	0.0	0.0	0.0
N ₂ P ₂ K ₂	2 : 1 : 1.5	22.8	12.8	11.7	47.3
N ₃ P ₃ K ₃	4 : 2 : 3	45.4	49.6	23.4	118.4
N ₄ P ₄ K ₄	6 : 3 : 4.5	68.0	75.0	35.0	178.0
N ₅ P ₅ K ₅	8 : 4 : 6	90.0	100.0	46.8	236.8
N ₆ P ₆ K ₆	10 : 5 : 7.5	113.6	125.0	58.6	297.2

Each NPK level was divided into two equal doses, the first dose was applied one month after sowing, and the second one was two weeks later.

The obtained seedlings received the other horticultural practices as common for Isabgol plants.

In both seasons, the effect of such two factors (sowing dates and NPK level) on chemical constituents of Isabgol seeds per plant was studied.

Samples of seeds were taken at harvest time (at 130 to 140 days from sowing date) when all spikes were filled with mature seeds and turned from pink to red colour. In addition, the lower leaves began to dry and the upper ones remained in yellow colour. Samples were oven dried at 70°C to determine seed content of mucilaginous substance (g per 5 g seeds). Mucilage percentage also was estimated according to the method described by Anderson (1949) in plants of each sowing date in the two tested seasons.

Isabgol substance extraction:

Representative samples (5 g) each were randomly taken from seed yield under each of the tested treatment. Such seeds sample under investigation were separately percolated with 90% ethanol (5 x 250 ml) to get rid of the clouring matter. The marc, in each case, was air dried, then extracted with cold water (3 x 500 ml) under stirring for 6 hours at room temperature and left to stand for another 12 hours. The supernatant in each case was filtered by passing through folded muslin and the filtrate in each case was centrifuged to remove the seed debris. The mucilages were then precipitated by adding to the aqueous solution four volumes of 95% ethanol under stirring. The precipitated mucilages were then separated by

centrifugation, washed several times with ethanol, vigorously shaken with absolute acetone, centrifuged again, pressed between two filter paper and left to dry in a desiccator over anhydrous calcium chloride to obtain the mucilages as a buff to dirty-white starchy powders. In all cases, the obtained mucilages were starch free, protein free, and did not reduce Fehling's solution.

The marc after the exhaustive extraction with cold water, was repeatedly extracted with boiling water (3 x 250 ml) until complete extraction of the mucilage is affected (Laid and Percival, 1950). The extracted mucilages were precipitated and purified adopting the same procedure mentioned before. The percentages of isolated mucilages were calculated.

Table 2. Mechanical and chemical analysis of the soil under investigation during 1995/96 and 1996/97 seasons.

Soil properties	1995/96	1996/97
Mechanical analysis:		
Clay (%)	41.70	42.93
Silt (%)	27.65	26.83
Fine sand (%)	28.22	28.39
Coarse sand (%)	2.43	1.85
Air pore space (%)	21.00	25.00
Chemical analysis:		
Available nitrogen (ppm)	43.00	47.00
Available phosphorus (ppm)	9.00	11.00
Available potassium (ppm)	2.51	2.89
Organic matter (%)	0.32	0.36
E.C. (mmhos/cm)	0.26	0.21
E.C. (%)	0.11	0.12
pH	7.63	7.52
CaCO ₃	2.51	2.33
Soluble cations and anions (meq/100gm soil):		
Ca	0.30	1.00
Mg	0.30	0.50
HCO ₃	0.70	0.50
Cl	0.35	0.80
SO ₄	0.28	0.64

Experimental design and statistical analysis

Factorial experiment in a randomized complete blocks design with 4 replicates was used according to Cochran and Cox (1957). Collected data were subjected to the statistical analysis according to the technique of analysis of variance (ANOVA) of factorial experiment in RCB design. The treatment means were compared using the least significant difference (LSD) procedures as mentioned by Gomez and Gomez (1984).

RESULTS AND DISCUSSION

The present research was trial to throw some lights on the relationship between sowing dates and NPK level solely or in combination and the mucilaginous substance in Isabgol seeds, which had different important was in certain medical purposes.

The obtained results are presented in Table (3) and showed the effect of sowing dates on mucilage content in the Isabgol seeds. It could be seen that plants grown in the first sowing date (First week of Nov.) reflected the highest values of mucilage content than the other sowing dates (1.57 g/5g or 31.49%) and (1.65 g/5g or 32.91%) in the first and second seasons, respectively.

Plants of the second sowing date (third week of Nov.) were the next in this respect, since they tabulated the values of 1.57 g/5g seeds or 31.39% and 1.60 g/5g or 32.01% for both seasons of study, respectively. On the other hand, fourth sowing date (third week of Dec.) indicated the least effect with 1.52 g/5g seeds or 30.39% and 1.63 g/5g or 30.58% values for the same seasons. The differences between sowing dates were significant in both seasons.

The above mentioned results behaved the same manner as those obtained by Russel (1975) on Isabgol, who reported that Isabgol plant when sown in late October and early November became a good source of mucilage content, which has an important uses as pharmaceuticals.

Table 3. Effect of sowing dates on the mucilage content in seeds as g per 5 g or as percentage of *Plantago ovata* in 1995/96 and 1996/97 seasons.

Sowing dates	Seeds mucilage content (g/5g)		Seeds mucilage content (%)	
	1995/96	1996/97	1995/96	1996/97
Nov. 1 st week	1.57	1.65	31.49	32.91
Nov. 3 rd week	1.57	1.60	31.39	32.01
Dec. 1 st week	1.53	1.56	30.53	31.27
Dec. 3 rd week	1.52	1.53	30.39	30.58
LSD at 5%	--	--	1.01	1.12

Concerning the effect of NPK levels, the data in Table (4) indicated that the applied NPK at any level significantly increased the mucilage percentage in seeds compared to the control (No NPK applied). Among the tested levels, the highest mucilage content and percentage (1.71 g/5g seeds or 34.27% and 1.76 g/5g or 35.17% in both seasons, respectively) were of plants treated with the fourth level (178.0 g/3m²) followed by those under the third NPK level (118.4 g/3m²), which tabulated for the first and second seasons, respectively, the values 1.62 g/5g seeds or 32.38% and 1.66 g/5g or 33.21%. The lowest mucilage content and percentage were of control plants and those received the second NPK level (47.3 g/3m²).

These results may be due to the influence of NPK fertilization on photosynthesis and the products. Potassium is necessary for the synthesis of carbohydrates and proteins, while phosphorus is an important factor influencing the rate of photosynthesis and phosphates act as catalysts for the action of some enzymes, which involved in the hydrolytic changes of starch to sugar (one of mucilage constituents) in plants (Alex, 1969). So, increasing of P and K fertilization increased the mucilage content and percentage.

Table 4. Effect of NPK levels on the mucilage content in seeds as g per 5 g or as percentage of *Plantago ovata* in 1995/96 and 1996/97 seasons.

NPK levels (g/3m ²)	Seeds mucilage content (g/5g)		Seeds mucilage content (%)	
	1995/96	1996/97	1995/96	1996/97
N ₁ P ₁ K ₁	1.45	1.46	28.91	29.26
N ₂ P ₂ K ₂	1.46	1.51	29.26	30.12
N ₃ P ₃ K ₃	1.62	1.66	32.38	33.21
N ₄ P ₄ K ₄	1.71	1.76	34.27	35.17
N ₅ P ₅ K ₅	1.58	1.63	31.53	32.64
N ₆ P ₆ K ₆	1.54	1.59	30.85	31.91
LSD at 5%	--	--	1.24	1.37

Comparing the results in Tables (3 and 4), it could be noticed that the effect of NPK at 178.0 g/3m² (fourth level) was higher than that of the earliest sowing date (First week of Nov.). The former factor recorded for mucilage constituents the values 1.745 g/5g seeds or 34.72%, whereas, the corresponding values for the later one were 1.61 g/5g seeds or 31.7%. These values are the average of both studied seasons.

Such significant effect of NPK at the fourth level is greatly confirmed the results reported above for Alex (1969), Khater *et al.* (1994), Dessouky (1995), Harphool *et al.* (1997) and Patel *et al.* (1997). They all reported that mucilage of Isabgol was highest with increasing N and P fertilization.

The effect of interaction between sowing dates and NPK levels is presented in Table (4). From this table, it was cleared that all combinations used gave a significant increase in mucilage % in seeds in both seasons, except for those which combined the lately sowing dates and either no application of NPK or low and excess NPK doses. Among such combinations, the greatest values were obtained for plant seeds from seeds are sown in Nov. 1st week and received the fourth level of NPK (178.0 g/3m²).

Otherwise, plant seeds from seeds of the fourth sowing date (Third week of Dec.) and did not receive NPK (0.0 level) resulted in the lowest content of mucilage in their seeds.

These results are expected, since there is a gradual decrease of temperature from October up to December. This gradual decrease makes a gradual tie up of the phosphorus in the soil and makes the phosphorus unavailable for plants as the temperature decrease. The shortage of phosphorus in the soil and the decrease of temperature during this time, restricts nitrogen and potassium uptake (El-Gamiely *et al.*, 1998).

Such findings were also in good agreement with the results obtained by Randhawa *et al.* (1978) on Isabgol, who reported that supplying Isabgol with 20 or 40 kg N/ha and sown on October 21 were favourable for good seed yield and high content of active constituents.

Table 5. Effect of interaction between the sowing dates and NPK levels on the mucilage content in seeds as percentage of *Plantago ovata* in 1995/96 and 1996/97 seasons.

Sowing Dates	NPK levels (g/cm ²)					
	0.0	47.3	118.4	178.0	236.8	297.2
1995/96 season						
Nov. 1 st week	29.28	29.73	32.52	33.86	29.89	29.78
Nov. 3 rd week	28.90	29.06	31.15	32.40	29.34	29.15
Dec. 1 st week	27.85	28.40	30.74	32.02	28.94	28.57
Dec. 3 rd week	27.57	27.96	30.02	31.52	28.40	28.12
LSD at 5%	1.31					
1996/97 season						
Nov. 1 st week	30.58	31.58	33.42	35.17	32.86	31.70
Nov. 3 rd week	29.60	30.13	32.33	34.10	31.90	30.81
Dec. 1 st week	28.34	29.01	31.96	33.21	30.71	29.90
Dec. 3 rd week	28.02	28.91	31.13	32.86	29.37	29.05
LSD at 5%	1.43					

The aforementioned results clearly proved the significant effect of sowing dates and NPK levels either solely or in combination on the active substance in Isabgol seeds. Therefore, the growers of such plant must give an attention on these factors. They provide early sowing date for seeds in the beginning and applied the most efficient dose of NPK during the growing season to produce an economic yield of mucilage content. This is in turn increase income from their Isabgol plantation.

REFERENCES

- Alex, L. (1969). Commercial flower. Forcing, 154-159 pp.
- Anderson, E. (1949). Endosperm mucilage of legumes occurrence and composition. Ind. Eng. Chem., 41:2887-2890.
- Anderson, J.W. (1987). Dietary fiber, lipids and atherosclerosis. American J. Cardiology, 60:17-22.
- Bloss, H.E. (1982). Formation of mycorrhizae in *Plantago ovata*. Mycologia, 74(3):505-508.
- Chiej, R. (1988). Medicinal Plants. MavDonald Co. Publishers Ltd. London, Sydney, P. 447.
- Cochran, W.G. and G.W. Cox (1957). Experimental Designs. 2nd Ed. John Wiley and Sons. In New York. USA, 595 P.
- Dessouky, M.M. (1995). Physiological studies on growth and active constituents of *Aloe vera*, L. Ph.D. Thesis, Fac. of Agric., Zagazig Univ.
- El-Gamiely, El.; H.M. Abd El-Naby; Hala. A.E. El-Sayed and A.E. Eata (1998). Effect of different planting dates and plant density on: 1. Vegetative and leaf chemical composition of peas. J. Agric. Sci. Mansoura Univ., 23(1):357-371.
- Gomez, K.A. and A.A. Gomez (1984). Statistical Procedures. Agric. Res., 2nd Ed. John Wiley and Sons, Inc, New York, USA.

- Harphool, S.; O.L. Sharma and R. Kumar (1997). Economics of nitrogen and phosphorus fertilization in Isabgol (*Plantago ovata* forsk). Crop Res., 11(2):246-247.
- Khater, M.R.; E.G. El-Ghauas and M.H. Abou Zied (1994). Effect of nitrogen and potassium fertilization on growth and yield of *Plantago psyllium*, L. Egypt. J. Appl. Sci., 9(10):796.
- Laid, R.A. and E.Q.V. Percival (1950). J. Chem. Soc., 2:1603.
- Marzouk, M.A. Amani (1995). A. Pharmacognotical study of certain plants containing mucilage and/or iridoids and growing in Egypt. M.Sc. Thesis, Fac. of Pharmact, Mansoura Univ., Egypt.
- Nordgaard, I.; H. Hove; M.R. Clausen and P.B. Mortensen (1996). Colonic production of butyrate in patients with previous colonic cancer during long-term treatment with dietary fiber (*Plantago ovata* seeds). Scandinavian J. Costroentology, 31:1011-1020.
- Patel, B.S.; S.G. Sadaria and J.C. Patel (1997). Influence of irrigation, nitrogen and phosphorus on yield, nutrient uptake and water-use efficiency on blond psyllium (*Plantago ovata*). J. Agron. Crop Sci., 17(3):145-216.
- Randhawa, G.S.; T. Sehata; D. Bains; V.P. Mahajan (1978). The effect of sowing date, seed rate and nitrogen fertilizer on the growth and yield of Isabgol (*Plantago ovata* F.). J. Agric. Sci. Cambridge, 80(2):341-343.
- Russell, T.E. (1975). Plantago wilt. Phytopathology, 65(3):359-360.
- Tackholm, V. (1974). Students Flora of Egypt. 2nd Ed. Cooperative Printing Company Press. Beirut, 516 PP.
- Trease, G.E. and W.C. Evans (1978). Pharmacogenosy. 11th Ed. Baillier, Tindall, London.
- Wallis, T.E. (1967). Textbook of Pharmacogenosy. J. and A. Churchill Ltd, 104 Glovcester Plance, London, 652 PP.

تأثير مواعيد الزراعة ومعدلات التسميد (النيتروجين - الفوسفور - البوتاسيوم)
على المحتوى الكيماوى لبيذور نباتات القاطونة
حكمت يحيى مسعود - حسين على أحمد - سهام محمد الجمل
قسم الخضر والزينة - كلية الزراعة - جامعة المنصورة

أجرى هذا البحث بمحطة بحوث ومعمل قسم الخضر والزينة - كلية الزراعة - جامعة المنصورة. وذلك خلال موسمين متتاليين ١٩٩٦/٩٥ و ١٩٩٧/٩٦. بهدف دراسة تأثير مواعيد الزراعة ومعدلات التسميد المختلفة (ن - فو - بو) والتفاعل بينهما على المحتوى الكيماوى للنبات من المواد المخاطية. تم دراسة أربعة مواعيد زراعة و٦ مستويات تسميد (ن فو بو) مختلفة. وكانت أهم النتائج المتحصل عليها مايلي:-

- ١- أعطت النباتات المنزرعة في ميعاد الزراعة الأول (الإسبوع الأول من نوفمبر) أعلى محتوى ونسبة مئوية من المسيلاج (المادة المخاطية).
- ٢- أدت معاملة النباتات بمستوى التسميد الرابع (١٧٨,٠ جم ن فو بو / ٣ م^٢) إلى الحصول على أعلى محتوى ونسبة مئوية من المسيلاج.
- ٣- تم الحصول على أعلى محتوى ونسبة مئوية من المسيلاج من التفاعل بين ميعاد الزراعة الأول ومستوى التسميد الرابع (١٧٨,٠ جم ن فو بو / ٣ م^٢).