

## EFFECT OF A TRANQUILIZING DRUG ADMINISTRATION ON SOME PHYSIOLOGICAL AND NUTRITIONAL RESPONSES OF BROILER CHICKS AT HIGH AMBIENT TEMPERATURES

Abd-Elaziz, A. M. S.<sup>1</sup>; A.M. Abdel-Maksoud<sup>1</sup>; Y. A. F. Hammouda<sup>2</sup> and Eman F.El-Daly<sup>2</sup>

1- Dept. Animal Prod., Fac. Agric., Ain Shams University, Cairo

2- National Res. Center, Dokki, Giza, Egypt.

### ABSTRACT

This experiment was conducted to study the possible beneficial effects of diazepam, a mild tranquilliser and appetite stimulator drug, on food intake and the physiological homeostasis of broiler chicks exposed to the high ambient temperatures of the summer months.

A total of 200, one day old, unsexed Hubbard broiler chicks were used. The chicks were allocated into five equal groups, one of them was kept as control while the others received different gradient levels of Diazepam for 6 weeks.

The results indicated that both live body weight and body weight gain at the end of the experiment (6 weeks) were significantly higher for birds at 0.5 gm/L of drinking water than the control and the other treatment groups. Changes in live body weight from 0 to 6 weeks of ages were 34.3; 37.7; 43.6; 40.2 and 37.9 folds for the control; 0.25; 0.50; 0.75 and 1.0 gm/L doses, respectively.

A significant increase in feed consumption was observed for birds treated by 0.50 gm/L of the drug at different ages. Similar trends were also observed for feed conversion ratio, protein consumption and efficiency of protein utilization in treated groups.

A significant dose-dependent increase in T4 and T3 levels at 4 and 6 weeks of age was also observed. Plasma total protein; triglycerides; cholesterol; creatinine; GOT and GPT levels were significantly different among treatment groups, except for the lower dose, and the control one.

Rectal temperature was significantly higher in the control and the 0.25 gm/L of diazepam groups than the other groups. Moreover, mortality rate was higher (10%) in the control group than in all treatment groups, where no mortalities occurred.

It could be concluded that, under hot summer conditions, water supplementation of 0.50 gm/L of diazepam is desirable to stimulate eating behaviour of broiler chicks, and to maintain their physiological equilibrium during high ambient temperature.

**Keywords:** heat stress, feed intake, blood picture, thyroid; growth, broilers.

### INTRODUCTION

One – of the first consequences of heat stress is a reduction in feed intake; lower growth; decreased feed efficiency and enhanced fat deposition (Geraert, *et al.*, 1993, 1996). Most studies deal with heat – induced hormonal changes and the endocrine control of thermogenesis. Plasma triiodothyronine (T3) levels appear to decrease, while thyroxine (T4) concentrations are not change (Klandorf, *et al.*, 1981; Sinurat, *et al.*, 1987) or even increase (Moss and Balnave, 1978).

In broilers, it has been shown that decreased rate of growth occurs (Howliger and Rose, 1989) when environmental temperature rises. This

negative effect is probably primarily due to reduced feed intake (Hurwitz, *et al.*, 1980). Moreover, Zuprizal, *et al.*, (1993) found that the true digestibilities of 12 amino acids were generally depressed when fed to broilers subjected to an increasing ambient temperature exposure from 21 to 32 °C. The growth reduction effect of heat stress was not related to feed intake (Dale and Fuller, 1980) and, thus, would have other origin.

Improving feed intake by dietary fat supplementation (Dale and Fuller, 1979) or by force feeding techniques up to 16% above adlib feed intake (Smith and Teeter, 1987) increased weight gain but was accompanied by enhanced fatness. Geraet *et al.*, (1996) reported that heat exposure appears to have an effect on growth performance independent of the decrease in feed intake. However, the reduction in growth is often greater than the reduction in feed intake.

Recent studies have been directed towards identifying specific brain sites and various neurotransmitters that are involved in food intake regulation (Denbow, 1989). Epinephrine and nor-epinephrine showed a central stimulatory effect on food intake in broilers (Denbow, *et al.*, 1981) but not in Leghorn (Denbow, *et al.*, 1983). On the contrary, serotonin has an inhibitory effect (Denbow, *et al.*, 1982, 1983).

Furthermore, there is evidence that Gamma – Aminobutyric acid (GABA), a neurotransmitter, can modulate food intake by opposing the inhibitory effect of serotonin (Morley, *et al.*, 1985), and may also play a role with angiotensin II in modulation of drinking behaviour (Swanson and Mogenson, 1981).

Pharmacological methods have been used to control appetite and body weight in broiler breeder birds (Oyawoya and Krueger, 1990). Since, recent advances in understanding the physiological control of feed intake have led to the identification of the potential veterinary products for regulating feed intake and body weight in chickens (El Halawani *et al.*, 1982). Tranquilizers have been suggested for use with poultry because of their stress ameliorating properties with possible beneficial effect on growth and feed efficiency, (Huston, 1959; Premachandra and Turner, 1960; Burger and Lorenz, 1960; Kicha, 1973; Kicka and Kamar, 1977; Abdel-Hakim, *et al.*, 1986; El-Habbak and Radwan, 1990). Since, diazepam, a mild tranquilizer, is used to stimulate eating behaviour and body weight of humans (Hoebel *et al.*, 1975); pigs (Dantzer, 1976); rats (Stapleton, *et al.*, 1979) and Japanese quail (Shoukry, 1993). This drug is classified as anxiolytic agent which potentiates the inhibitory effect of GABA on the serotonergic neurons, its receptors are present in both mammals and birds, its binding capacity to peripheral tissues are very low compared to brain tissues, and finally it has a prolonged half life (Feldman and Quenzer, 1984). These observations suggest that, diazepam may be functioning as an appetite – stimulating drug for broiler chicks exposed to high ambient temperature stress during the summer months. Since, the present study was undertaken to provide further information about the physiological and nutritional responses of broilers to diazepam administration.

## MATERIALS AND METHODS

### Birds and management:

A total number of 200 Hubbard, one day old, unsexed broiler chicks from a commercial supplier were brooded in floor pens during the period from July, 20<sup>th</sup> to September, 3<sup>rd</sup> / 1998. During the first week of age, the brooding temperature was maintained at 35±1°C with a constant photoperiod of 24h. Feed and water were provided, ad libitum, throughout the whole experimental period. Compositions of both the starter and finisher diets are shown in Table (1). The finisher diet was used after 3 weeks of age.

After the brooding week, the chicks were assigned randomly to five treatment groups, 40 chicks each. One group was a control and freely allowed to drink tap water with no additives, and the other four groups were given graded doses (via drinking water) of diazepam (Ratiopharm GmbH Arzneimittel, 7902 Blaubeuren, Germany). The doses were 0.25; 0.50; 0.75 and 1.0 gm/L of drinking water for the four groups, respectively.

**Table (1): Composition of the starter and finisher diets.**

Ingredients	Starter	Finisher
Yellow corn	65	74
Soybean meal (44%)	24	15
Herring fish meal	4	3
Corn gluten	4	4.5
Bone meal	1.6	2.0
Limest one	0.65	0.70
Premix (1)	0.25	0.25
NaCl	0.20	0.25
Lysine	0.14	0.20
DL - Methioine	0.16	0.10
Total	100	100
<b>Calculated analysis</b>		
CP	22.18	19.20
ME (Kcal/Kg)	3018	3110
C/P ratio	137	163
Ca%	1.12	1.10
P%	0.70	0.62
Methionine %	0.62	0.56
Lysine	1.20	1.06
Methionine + Cyst	0.96	0.73

(1) Broiler premix, each 1kg contained vit. A 12000 IU; Vit D3 2500 IU; Vit. E. 10 mg; Vit. K3 mg; Vit. B1 1 mg; Vit B2 4 mg; Pantothenic acid 10 mg; Folic acid 1 mg; D – biotin 0.5 mg; Niacine 40 mg; Vit, B6 – 3 mg; Vit. B12 – 20 mg; Mn – 62 mg; Fe – 44 mg; Zn – 56 mg; Cu – 5 mg and Se – 100 mg.

### Meteorological data:

Indoor ambient air temperature (Max.; Min.) and relative humidity (RH) were recorded daily. Averages of Max. and Min. ambient air temperatures and RH during the experimental period were 36.8°C; 25.4°C and 59%, respectively.

### Measurements:

Live body weight; weight gain; feed consumption; feed conversion; protein consumption, and efficiency of protein utilization were determined and

calculated in a biweekly basis (2, 4 and 6 weeks of age).

At 4 and 6 weeks of age, blood samples were withdrawn, via the wing vein, from ten birds of each treatment group. Blood serum was obtained, while EDTA was used as an anticoagulant. All samples were centrifuged at 6000 r. p. m. for 10 minutes, plasma or serum was then decanted and stored frozen at - 20°C until analysed.

Plasma thyroxine (T4) and triiodothyronine (T3) were determined by the enzymatic – colorimetric method, using available commercial kits (bio Merieux, 69280 Mary – l' Etoile, France). Total plasma protein, triglycerides; Cholesterol, Creatinine; serum glutamic oxaloacetic transaminase (GOT); and serum glutamic pyruvic transaminase (GPT) were also determined using available commercial kits. A drop of blood / bird was smeared, stained (Wright's stain) and then examined for the differential counts of blood leukocytes to calculate the heterophils (H) and lymphocytes (L) numbers and the H/L ratio.

**Statistical analysis:**

Data were subjected to the ANOVA using the SAS general Linear Models procedure (SAS Institute, 1994) and to multiple range test (Duncan, 1955) to test the differences among means.

## **RESULTS AND DISCUSSION**

### **Live body weight (LBW) and weight gain (WG):**

Table (2) shows the effect of diazepam on both LBW and WG of broiler chicks at 2, 4 and 6 weeks of age. It is clear from the results that the higher doses of diazepam (0.75 and 1 gm/L) gave the best LBW and WG of chicks at 2 weeks of age. However the treated groups have Significantly ( $P \leq 0.05$ ) higher body weights at the 4th and the 6<sup>th</sup> weeks of age.

At 4weeks of age, a significant increase in LBW of birds treated with 0.25 and 0.50 gm/L of the drug was observed. However, at 6 weeks of age birds that given the 0.50 and 0.75 gm/L doses surpassed the other treatments and recorded significantly ( $P \leq 0.05$ ) the highest LBW. Moreover, analysis of variance data (Table-2) revealed highly significant effects of treatment, age and their interaction on both LBW and WG.

The results show also that the body weight gain increased significantly as the dose of diazepam increased. The best values obtained, however, were for birds that treated with 0.50 and 0.75 mg/L doses, respectively. It is of interest to notice that the LBW of all treatment groups increased dramatically from hatch to the marketable age (6 weeks). The percentages of change that were about 34.3; 37.7; 43.6; 40.2 and 37.9 folds for the control, 0.25; 0.50; 0.75 and 1 gm/L of diazepam treated groups, respectively. It appears from these results that diazepam administration (0.50 or 0.75 gm/L) via drinking water improved LBW and WG of broiler chicks exposed to high ambient temperatures. The averages of the maximum and minimum ambient temperature that recorded throughout the experimental period were 36.8°C and 25.4°C, respectively.

Table (2): Body weight and weight gain (WG) in grams of broiler chicks as affected by treatments at different ages.

Age(wk) Dose(G/L)	One day	2	WG	4	WG	6	WG
0	43.2 ± 0.53 <sup>a</sup>	310.6 ± 30.2 <sup>b</sup>	267.8.3 ± 15.1	896.3 ± 86.5 <sup>b</sup>	586.2 ± 36.6	1480.2 ± 116.2 <sup>c</sup>	583.9 ± 46.6 <sup>e</sup>
2.5	42.6 ± 0.75 <sup>a</sup>	303.2 ± 18.8 <sup>b</sup>	262 ± 18.6	1060.2 ± 104.2 <sup>c</sup>	756.4 ± 58.3	1604.8 ± 131.8 <sup>c</sup>	546.8 ± 40.2 <sup>c</sup>
5	42.8 ± 0.55 <sup>a</sup>	318.5 ± 23.4 <sup>bc</sup>	277.8 ± 20.8	1126.0 ± 88.2 <sup>a</sup>	810.5 ± 51.3	1866.5 ± 128.2 <sup>a</sup>	740.4 ± 50.2
7.5	43.0 ± 0.64 <sup>a</sup>	330.1 ± 29.8 <sup>a</sup>	286.3 ± 19.7	968.8 ± 65.8 <sup>b</sup>	647.8 ± 41.2	1730.3 ± 139.8 <sup>b</sup>	761.2 ± 67.4
10	42.8 ± 0.66 <sup>a</sup>	321.7 ± 16.6 <sup>ac</sup>	278.81 ± 19.8	987.3 ± 80.6 <sup>b</sup>	666.5 ± 44.6	1620.4 ± 108.5 <sup>c</sup>	633.2 ± 41.6

Significance of effects (n=200)

Treatment (T)	NS	*	*	**	***	***	**
Age (A)	NS	**	*	***	**	**	**
T x A	NS	NS	NS	**	**	**	**

1\* P ≤ 0.05; \*\* P ≤ 0.01; \*\*\* P ≤ 0.001; NS - Not Significant  
a, b, ... Means with different superscripts within a column are significantly different (P ≤ 0.05)

Although this temperature was reported to reduce the growth of broilers (Smith and Teeter, 1987; Howliger and Rose 1989 and Geraert, *et al.*, 1996), the present results clearly indicate a pronounced increase in broilers body weight which reflects the importance of diazepam administration during hot months. It seems that diazepam enhances food intake and feed utilization of birds by a direct effect on birds thermobalance during hot weather. It may also exerts an effect on the food regulating centers in the brain either by decreasing serotonin secretion or by a main effect on the brain satiety centers. These results were supported by the findings of many authors in different mammalian and avian species (Kicka, 1973; Hoebel, *et al.*, 1975; Dantzer, 1976; Kicka and Kamar, 1976; Stapleton, *et al.* ; 1979; EL-Halawany, *et al.* 1982 and others.

**Some nutritive responses to diazepam administration:**

The dose response of feed consumption and efficiency to diazepam supplement were found to be significantly different during the experimental period. Table (3) shows that feed consumption was significantly ( $P \leq 0.05$ ) higher for the 0.25 and 1.0 gm/L groups than the other treatments and control groups at 2 weeks of age. These differences were more obvious at 4 and 6 weeks of age, where feed consumption increased significantly ( $P \leq 0.05$ ) for the 0.50; 0.75 and 1.0 gm/L of drinking water groups.

At doses of 0.50 and 0.75 gm/L of drinking water, diazepam significantly improved feed efficiency of birds at 2, 4 and 6 weeks of age. There were significant differences between the treated and the control groups, since the ANOVA data revealed a highly significant ( $P \leq 0.01$ ) effect of treatment, age and their interaction on all nutritional traits studied.

Similar trends were also observed for protein consumption and utilization, which may suggest that diazepam, enhances the digestibility and or absorption of dietary proteins. Although the efficiency of protein utilization was reported to be decreased (Zuprazil, *et al.*, 1993) in broilers reared at high ambient temperatures, the results obtained herein indicate that diazepam can alleviate these deleterious effects. It seems that diazepam administration at 0.50 or 0.75 gm/L dose would protect birds against the anorectic effects due to heat exposure, by stimulating the eating behaviour of chickens.

That diazepam improved feed consumption and subsequently the birds performance, may be due to its inhibitory effect on the serotonergic neurons in the higher brain centers, and / or its direct or indirect effect on the thyroid releasing hormone (TRH) secreting nucleus in the hypothalamus. It has been reported that diazepam potentiates GABA which in turn antagonize the inhibitory effect of serotonin on food intake (Stapleton, *et al.*, 1979 and Morley, *et al.*, 1985). Since the results indicate that diazepam effects were neural rather than metabolic.

Moreover, there is evidence that diazepam decreased plasma glucose concentration in birds (Dantzer, 1976; Swanson and Mogenson, 1981; and Shoukry, 1993). Since the role of plasma glucose level on food intake was reported to be mediated through its action on hepatic receptors rather than central via brain receptors. Such a reduction in plasma glucose may cause the bird to respond by increasing food intake to obtain or to

compensate for its energy requirements. In agreement with the present results were the findings of Denbow, *et al.*, (1983); Denbow, (1984 and 1989); Gonzalez, *et al.*, (1984); Ida *et al.*, (1985); Klandorf and Harvey. (1985); Oyawoye and Krueger (1990) and others.

**Effects of diazepam on some blood constituents:**

Plasma levels of T4-T3 and the T4-T3 ratio in both the control and diazepam - supplemented groups at 4 and 6 weeks of age are presented in Table (4). It is clear that T4 level increased significantly ( $P \leq 0.05$ ) in birds at the 0.50 and 0.75 gm/L groups, respectively. This increase reflects a stimulating effect of the drug on thyroid activity.

A similar trend was also observed for T3 levels at 4 and 6 weeks of age. The results reveal that the drug administration would be beneficial in maintaining an euthyroid status for birds exposed to high ambient temperature, which was around 36.8°C during this experiment. Since, the depressing effects of an ambient temperature above 29°C on the thyroid activity could be alleviated by diazepam administration.

Most studies deal with heat-induced hypothyroidism in broilers which improved the physiological adaptation of birds to heat stress by the modulation of metabolic rates (Klandorf, *et al.*, 1981; Sinurat, *et al.*, 1987 and Geraert, *et al.*, 1996). Although the results of Shoukry (1993) indicate that plasma T4 decreased by a single dose of diazepam injection (in crop) in Japanese quail, no such decrease was observed in the present study. This may be due to the different magnitude and duration of the drug administration and also to the different avian species used.

Concerning plasma total protein (TP) level, the data indicate a significant increase in TP of birds treated with 0.25; 0.50 and 0.75 gm/L of the drug than those treated with 1.0 gm/L and the control group. In addition, a significant dose-dependent increase in Plasma triglycerides, cholesterol and creatinine levels were also observed. These results indicate a considerable changes in the nutrients metabolism as a consequence of the drug supplementation. This may also indicate that diazepam can affect the lipolytic action of the adrenergic neurons. Diazepam administration has been reported to affect the adrenergic receptors and lipid and protein mobilization and degradation in birds which confirms the results of the present study (Denbow, *et al.*, 1981; Colasanti, 1982 and Feldman and Quenzer, 1984).

There were also significant in the serum GOT and GPT among the control and the treated groups. A dose - dependent increases in both enzymes were observed at 4 and 6 weeks of age. The increased SGOT and SGPT levels may be attributed to the effect of diazepam on liver function.

This was evident as some birds of the higher doses groups of the drug (above 0.5 gm/L) show signs (at autopsy) of liver subcutaneous blood hematomases, and hypertrophy of liver, spleen and the right kidney. These signs, however, did not cause any mortality losses in all the treated groups, although the mortality rate of the control group was higher (10%) as a result of heat stress imposed during the experimental period. Table (4) shows also that the drug administration improved the resistance of birds to heat stress as indicated by the lower H/L ratio obtained for group treated with doses of 0.50, 0.75 and 1.0 gm/L of diazepam at 4 and 6 weeks of age.

Table (3): Effect of treatments on feed consumption (FC) feed efficiency (FE) ; Protein consumption (PC) and efficiency of protein utilization (EPU) of chicks at 2, 4 and 6 weeks of age.

Age(wk)	2			4			6					
	FC(g)	FE	PC(g)	EPU	FC(g)	FE	PC(g)	EPU	FC(g)	FE	PC(g)	EPU
Dose(G/L)	350 ± 28 <sup>b</sup>	1.12 ± 0.5 <sup>c</sup>	77 ± 9 <sup>2</sup>	0.29 ± 0.03 <sup>b</sup>	1318 ± 64 <sup>c</sup>	1.47 ± 0.15 <sup>b</sup>	290 ± 21 <sup>b</sup>	0.49 ± 0.13 <sup>b</sup>	3285 ± 62 <sup>d</sup>	2.23 ± 0.09 <sup>c</sup>	690 ± 43 <sup>d</sup>	1.31 ± 0.20 <sup>b</sup>
0	371 ± 22 <sup>a</sup>	1.23 ± 0.09 <sup>a</sup>	81.8 ± 8.6 <sup>a</sup>	0.32 ± 0.06 <sup>a</sup>	1420 ± 53 <sup>b</sup>	1.30 ± 0.12 <sup>a</sup>	312 ± 18 <sup>a</sup>	0.41 ± 0.09 <sup>a</sup>	3461 ± 89 <sup>f</sup>	2.14 ± 0.12 <sup>b</sup>	727 ± 38 <sup>e</sup>	1.35 ± 0.18 <sup>c</sup>
2.5	366 ± 18 <sup>ab</sup>	1.15 ± <sup>b</sup>	80.6 ± 10 <sup>ac</sup>	0.29 ± 0.09 <sup>b</sup>	1481 ± 48 <sup>a</sup>	1.28 ± 0.09 <sup>a</sup>	328 ± 24 <sup>a</sup>	0.40 ± 0.14 <sup>a</sup>	3660 ± 101 <sup>a</sup>	1.96 ± 0.08 <sup>a</sup>	769 ± 56 <sup>a</sup>	1.03 ± 0.12 <sup>a</sup>
5.0	355 ± 26 <sup>b</sup>	1.07 ± 0.06 <sup>b</sup>	78.5 ± 11.2 <sup>bc</sup>	0.27 ± 0.10 <sup>b</sup>	1340 ± 66 <sup>c</sup>	1.38 ± 0.14 <sup>a</sup>	295 ± 20 <sup>b</sup>	0.46 ± 0.12 <sup>b</sup>	3559 ± 74 <sup>b</sup>	2.05 ± 0.14 <sup>b</sup>	747 ± 62 <sup>b</sup>	0.98 ± 0.20 <sup>a</sup>
7.5	380 ± 26 <sup>a</sup>	1.18 ± 0.12 <sup>a</sup>	83.6 ± 8.5 <sup>a</sup>	0.29 ± 0.08 <sup>b</sup>	1375 ± 58 <sup>c</sup>	1.40 ± 0.12 <sup>b</sup>	303 ± 27 <sup>b</sup>	0.46 ± 0.12 <sup>b</sup>	3491 ± 93 <sup>b</sup>	2.15 ± 0.12 <sup>c</sup>	733 ± 42 <sup>c</sup>	1.26 ± 0.21 <sup>b</sup>
10												
Significance of effects												
Treatment (T)	*	*	*	*	**	*	*	*	**	**	*	**
Age(A)	NS	**	*	*	**	**	**	*	**	*	**	*
T X A	NS	*	*	*	**	*	*	*	**	*	*	*

a-c: Means within a column having different superscripts are significantly different (P ≤ 0.05)

Table (4): Effect of drug administration on some blood parameters at 4 and 6 weeks of age.

Dose gm/L Item	0		2.5		5.0		7.5		10.0	
	4wk	6	4	6	4	6	4	6	4	6
T <sub>4</sub>	12.6 ± 1.6 <sup>b</sup>	14.2 ± 2.2 <sup>b</sup>	12.9 ± 1.4 <sup>b</sup>	15.4 ± 1.8 <sup>b</sup>	20.9 ± 2.4 <sup>a</sup>	20.5 ± 3.4 <sup>a</sup>	17.5 ± 3.2 <sup>a</sup>	18.7 ± 3.5 <sup>a</sup>	10.9 ± 4.2 <sup>b</sup>	15.9 ± 2.8 <sup>b</sup>
T <sub>1</sub>	3.9 ± 0.91 <sup>a</sup>	2.8 ± 1.2 <sup>b</sup>	4.2 ± 1.0 <sup>a</sup>	3.2 ± 1.2 <sup>b</sup>	4.4 ± 0.9 <sup>a</sup>	4.2 ± 1.3 <sup>a</sup>	3.7 ± 1.2 <sup>b</sup>	4.2 ± 1.2 <sup>a</sup>	3.7 ± 1.7 <sup>a</sup>	3.9 ± 1.4 <sup>a</sup>
T <sub>3</sub> /T <sub>4</sub>	0.30 ± 0.06 <sup>a</sup>	0.19 ± 0.03 <sup>b</sup>	0.32 ± 0.08 <sup>a</sup>	0.20 ± 0.04 <sup>b</sup>	0.21 ± 0.06 <sup>b</sup>	0.20 ± 0.05 <sup>b</sup>	0.22 ± 0.03 <sup>a</sup>	0.21 ± 0.05 <sup>ab</sup>	0.35 ± 0.02 <sup>b</sup>	0.24 ± 0.06 <sup>a</sup>
T protein	4.2 ± 1.0 <sup>a</sup>	3.9 ± 1.01 <sup>a,b</sup>	4.2 ± 0.93 <sup>a</sup>	4.01 ± 0.9 <sup>ab</sup>	4.7 ± 1.0 <sup>a</sup>	4.9 ± 1.1 <sup>a</sup>	4.7 ± 1.2 <sup>a</sup>	4.4 ± 1.1 <sup>a</sup>	3.9 ± 1.1 <sup>b</sup>	3.7 ± 1.3 <sup>b</sup>
Triglyc.	320.6 ± 52.8 <sup>b</sup>	380.8 ± 46.6 <sup>b</sup>	375.6 ± 44.6 <sup>a</sup>	396.0 ± 60.2 <sup>b</sup>	430.5 ± 56.4 <sup>a</sup>	465.1 ± 70.0 <sup>a</sup>	395.6 ± 62.2 <sup>b</sup>	474.2 ± 73.2 <sup>a</sup>	418.8 ± 74.3 <sup>b</sup>	484.5 ± 81.2 <sup>a</sup>
Cholest	82.5 ± 18.8 <sup>ab</sup>	91.6 ± 15.4 <sup>c</sup>	100.6 ± 16.8 <sup>a</sup>	94.5 ± 16.8 <sup>a</sup>	125.8 ± 23.2 <sup>b</sup>	134.8 ± 22.5 <sup>b</sup>	118.9 ± 19.8 <sup>b</sup>	142.6 ± 29.2 <sup>b</sup>	102.5 ± 21.0 <sup>a</sup>	198.9 ± 25.2 <sup>a</sup>
Creatinine	0.64 ± 0.08 <sup>b</sup>	0.72 ± 0.11 <sup>c</sup>	0.71 ± 0.12 <sup>b</sup>	0.77 ± 0.14 <sup>c</sup>	0.86 ± 0.09 <sup>a</sup>	0.88 ± 0.16 <sup>b</sup>	0.85 ± 0.16 <sup>b</sup>	0.98 ± 0.2 <sup>b</sup>	1.1 ± 0.32 <sup>b</sup>	1.26 ± 0.28 <sup>a</sup>
GOT	60.1 ± 9.2 <sup>b</sup>	71.8 ± 11.3 <sup>c</sup>	78.3 ± 10.5 <sup>a</sup>	81.8 ± 12.4 <sup>c</sup>	92.2 ± 18.6 <sup>a</sup>	98.5 ± 13.1 <sup>b</sup>	96.4 ± 16.2 <sup>a</sup>	101.4 ± 19.3 <sup>b</sup>	106.8 ± 21.8 <sup>b</sup>	126.5 ± 20.8 <sup>a</sup>
GPT	7.8 ± 1.4 <sup>a</sup>	10.2 ± 9.4 <sup>c</sup>	10.8 ± 1.1 <sup>ab</sup>	12.0 ± 1.2 <sup>c</sup>	22.3 ± 2.6 <sup>b</sup>	29.9 ± 3.2 <sup>b</sup>	18.9 ± 1.2 <sup>b</sup>	31.4 ± 3.3 <sup>b</sup>	32.7 ± 3.6 <sup>b</sup>	43.5 ± 4.9 <sup>a</sup>
H/L	0.46 ± 0.02 <sup>a</sup>	0.68 ± 0.09 <sup>a</sup>	0.53 ± 0.1 <sup>a</sup>	0.58 ± 0.06 <sup>b</sup>	0.50 ± 0.12 <sup>a</sup>	0.37 ± 0.08 <sup>c</sup>	0.35 ± 0.09 <sup>b</sup>	0.35 ± 0.12 <sup>c</sup>	0.31 ± 0.05 <sup>b</sup>	0.24 ± 0.06 <sup>d</sup>
RT	41.3 ± 0.08 <sup>a</sup>	42.6 ± 0.06 <sup>b</sup>	41.2 ± 0.07 <sup>a</sup>	42.3 ± 0.12 <sup>b</sup>	41.2 ± 0.10 <sup>a</sup>	41.5 ± 0.06 <sup>b</sup>	40.8 ± 0.13 <sup>b</sup>	41.2 ± 0.09 <sup>b</sup>	40.5 ± 0.11 <sup>b</sup>	41.03 ± 0.08 <sup>b</sup>

a-d, Means with different superscripts within a row in the same ages are significantly different ( $P \leq 0.05$ ).

This effect was also accompanied by a lower rectal temperature of birds in the higher doses groups. The desirable effects of the drug in decreasing both the H/L ratio and RT may suggest the drug administration during the summer months where the air temperature rises. It is also suggested that H/L ratio may be used as a criterion to measure the resistance of birds to stress condition. This was supported by the findings of Zulkifli and Siegel (1995) who reported that an L/H ratio lower than 0.5 is a good indicator that the bird is more heat tolerant.

It is concluded, from the present results, that diazepam administration at 0.50 gm/L of drinking water during the hot summer months would be a suitable appetite - stimulator and a growth enhancing agent for broilers. It would be suggested to use diazepam in the drinking water of broilers to maintain their physiological equilibrium and productive performance under stressful conditions. The role that the drug plays at the brain receptors and opioid peptides need, however, further investigations.

## REFERENCES

- Abdel-Hakim, N.F.; A.A. Amer and N.M. EL-Naggar (1986.) Effect of reserpine on performance of broiler chicks. 11th International cong. Statis., comput., Sci., Demog. Res., Cairo, 283-296.
- Burger, R. E. and F. W. Lorenz (1960) pharmacologically induced resistance to heat shock. *Poult. Sci.*, 39:468 - 473.
- Colasanti, B. K. (1982). Parasympathomimetic drugs in: *Modern pharmacology* Edited by C.R. Craig and R. E. Stitzel. Published by Little, Brown and Company, Boston, 157-163.
- Dale, N. M. and H. L. Fuller (1979). Effects of diet composition on feed intake and growth of chicks under heat stress. I-Dietary fat levels. *Poult. Sci.*, 58:1529 - 1534.
- Dale, N. M. and H. L. Fuller (1980). Effects of diet composition on feed intake and growth of chicks under heat stress. II-Constant vs. cycling temperatures. *Poult. Sci.*, 59:1434 - 1441.
- Dantzer R. (1976). Effect of Diazepam on performance of pigs in a progressive ration schedule. *Physiology and Behavior*, 17: 161 - 163.
- Denbow, D. M. (1989). Peripheral and central control of food intake. *Poultry Science*, 68:938-947
- Denbow, D. M. (1984) Body temperature and food intake of turkeys following ICV injections of serotonin *Nutr. Behav.*, 1:301- 308.
- Denbow, D. M.; J. A. Cherry; P. B. Siegel and H. P. Van kery (1981). Eating, drinking and temperature response of chicks to brain catecholamine injections. *Physiol. Behav.*, 27:265- 269.
- Denbow, D. M.; H. P. van kery and J. A. Cherry (1982). Feeding and drinking response of young chicks to injections of serotonin into the lateral ventricle of the brain. *Poultry Science*, 61: 150-155
- Denbow, D. M.; H. P. Van kery; M.P. Lacy and T. J. Dietrick (1983). Feeding, drinking and body temperature of Leghorn chicks: Effects of ICV injection of biogenic amines. *Physiol. Behav.*, 31 :85-90

- Duncan, D. B. (1955). Multiple range and multiple "F" test. *Biometrics*, 11: 1-42.
- El- Halabak, M. M. E. and A. A. Radwan (1990). Some physiological responses of adult cocks to the tranquilizing drug reserpine. *Egypt. Poultry Sci.*, 10: 239- 252
- EL- Halawani, M. E.; W. H. Bruke; P. T. Dennison and J. L. Silsby (1982). Neuropharmacological aspects of neural regulation of avian endocrine function in: *Aspects of Avian Endocrinology: practical and theoretical implications*. Edited by Scams, G. Cr; Ottinger, M. A; Kenny A.D; Balthazrt, J.; Cronshaw, J. and Chester, I Jones. Published by Texas Tec. Press, Texas, pp.33-60
- Feldman R. S. and L. F. Quenzer (1984). *Fundamentals of Neuropsychopharmacology*. published by Sinaur Associates, Inc., Massachusetts, 268-285
- Geraert, P.A.; J. C. Padilha and S. Guillaumen (1993). Metabolic and endocrine changes induced by heat exposure in chickens. *Proc. Nutr. Soc.*, 52:165
- Geraert, P. A.; J. C. Padilha and S. Guillaumen (1996). Metabolic and endocrine changes induced by chronic heat exposure in broiler chickens: growth performance, body composition and energy retention. *Brit. J. Nutrition*, 75: 195-204
- Gonzalez, Y.; M.P. Fernandez; F. Sanchez and J. Rio (1984). Antagonism of diazepam - induced feeding in rats by anti-sera to opioid peptides. *Life Science*, 35:1423-1429.
- Hoebel, B. G.; I. K. Krauss and D. Willard (1975). Body weight decreased in human by phenylpropanolamine taken before meals. *Obesity Bariatric Medicine*, 4: 200-206
- Howlader, M. A. and S. P. Rose (1989). Rearing temperature and the meat yield of broilers. *Brit Poultry Sci.*, 30:61 – 67
- Hurwitz, S. I.; M. Weiselberg; V. Eisner; I. Bartov; G. Riesenfeld and S. Bornstein(1980). The energy requirements and performance of growing chickens and turkeys as affected by environmental temperature. *Poultry Sci.*, 59:2290-2299
- Huston, T. M. (1959) The effect of serpasil on controlled and naturally occurring climatic thermal stresses on immature domestic fowl *Poultry Sci.*, 43: 120-126
- Ida, Y.; M. Tanaka; A. Tsuda; S. Tsujimaru and N. Nagasaki(1985). Attenuating effect of diazepam on stress- induced increases in noradrenaline turnover in specific brain regions of rats: Antagonism by Ro-15-1788. *Life Science*, 37: 2489- 2491.
- Kicka, M. A. M. (1973). The role of vitamin C and tranquilizers on heat regulation and subsequent productivity of chickens. Ph.D. thesis, Univ. of Cairo.
- Kicka, M. A. M. and G. A. R. Kamar (1977). Effect of tranquilizers and bird density on chicken growth under heat stress. *Egypt J. Anim. Prod.*, 17:55-61

- Klandorf, H. and S. Harvey (1985). Food intake regulation and circulating thyroid hormones in domestic fowl. *General and Comparative Endocrinology*, 60:162- 170.
- Klandorf, H.; P.J. Sharp and M. G. Macleod (1981). The relationship between heat production and concentrations of plasma thyroid hormones in the domestic hen. *Gen. Compar. Endocrin.*, 45:513- 522
- Morley, J. E.; T. J. Bartness; Gosnell and A. S. Levine (1985). Peptidergic regulation of feeding. *International Rev. Neuroendocrin*, 27: 207-281.
- Moss, B. and D. Balnave (1978). The influence of elevated environmental temperature and nutrient intake on thyroid status and hepatic enzyme activities in immature male chicks. *Compar. Bioch. Physiol.*, 157- 161
- Oyawaye, E. O. and W. F. Krueger (1990). Potential of chemical regulation of food intake and body weight of broiler breeder chicks. *Brit. Poul. Sci.*, 31: 735 -742
- Premachandra, B. N. and C. W. Turner (1960). Reserpine and thyroid activity in fowls. *Proc. Soc. Exp. Biol & Med.*, 104: 306 - 309.
- SAS Institute (1994). *SAS / STAT ® User's Guide Statistics*, Ver. 604, Fourth Ed. SAS Institute Inc., Carry, NC.
- Shoukry, H. M. S. (1993). The effects of single dose of anxiolytic agent (Diazepam) on food intake in male Japanese quail. *Al-Azhar J. Agric. Res.*, 17:301 -316.
- Sinurat, A. P.; D. Balnave and G. H. McDowell (1987). Growth performance and concentrations of thyroid hormones and growth hormone in plasma of broilers at high temperatures. *Aust. J. Biol. Sci.*, 40: 443 - 450.
- Smith, M. O. and R. G. Teeter (1987). Force Feeding methodology and equipment for poultry. *Poult. Sci.*, 66 (Suppl. 1), 185 (Abstract).
- Stapleton, J. M.; D. L. Marcia; J.M. Vicki and D. R. Larry (1979). Naloxone inhibits Diazepam-induced feeding in rats. *Life Science*, 24: 2421-2426.
- Swanson, L. W. and G. J. Mogenson (1981). Neural mechanisms for the functional coupling of autonomic, endocrine and somatomotor responses in adaptive behavior. *Brain Research Reviews*, 3: 1-34
- Zulkifli, I. and P. B. Siegel (1995). Is there a positive side to stress. *World's Poul. Sci. J.*, 51:63-76.
- Zuprazil, M. L.; A. M. Chagneau and P. A. S. Geraert (1993). Influence of ambient temperature on true digestibility of protein and amino acids of rapeseed and soybean meals in broilers. *Poult. Sci.*, 72: 289-295.

تأثير إعطاء أحد المواد المهدنة على بعض الاستجابات الفسيولوجية الغذائية في  
دجاج اللحم المربي تحت ظروف الحرارة المرتفعة .  
أحمد محمد سعيد عبد العزيز<sup>١</sup>، عبد المجيد مخيمر عبد المقصود<sup>١</sup>، ياسر أحمد فتحى حمودة<sup>٢</sup>،  
إيمان فرج الدالي<sup>٢</sup>  
١- قسم الإنتاج الحيواني - كلية الزراعة - جامعة عين شمس - القاهرة  
٢- المركز القومي للبحوث - الدقى - الجيزة .

اجريت هذه الدراسة لمعرفة تأثير أحد العقاقير المهدنة على بعض الصفات الإنتاجية والفسيولوجية  
في دجاج اللحم المربي على درجة حرارة مرتفعة أثناء شهر الصيف. استخدم في التجربة ٢٠٠ كتكوت  
تسمين قُسمت إلى خمسة مجموعات إحداهما للمقارنة والباقي حسب الجرعة المعطاة هي ٠، ٢،٥ ، ٥ ، ٧،٥ ،  
١٠ ملليجرام لكل لتر من مياه الشرب واستمرت المعاملة حتى نهاية التجربة.  
اتضح من النتائج وجود زيادة معنوية في وزن الجسم ومعدل النمو للمجموعات التي عوملت  
بالمهدئ وكانت أفضل النتائج عند الجرعات ٥ و ٧،٥ ملليجرام / لتر.  
كان للمعاملات تأثير معنوي على زيادة معدل استهلاك الغذاء (رغم ارتفاع الحرارة) وذلك في جميع  
الأعمار كما ارتفع معدل التحويل الغذائي خاصة عند الجرعتان ٥ و ٧،٥ ملليجرام.  
انخفضت درجة حرارة المستقيم معنوياً بزيادة الجرعة وكذلك نسبة النفوق.  
لوحظ أن مستوى هرمونات الغدة الدرقية قد ارتفع معنوياً في الدم بزيادة الجرعة وذلك عند عمر ٤  
، ٦ أسابيع كما تأثر مستوى بروتين البلازما والجلوسيدات الكلية والكولسترول والكرياتينين والانزيمات  
الناقلة لمجموعة الأمين وذلك بزيادة الجرعة أيضاً.  
لم تؤثر المعاملات على المظهر التشريحي للطيور بصفة عامة وإن وجدت بعض حالات فردية قليلة  
في المجموعات التي أعطيت جرعات مرتفعة (٧،٥ و ١٠ ملليجرام) ممثلة في وجود ارتشاح دموي بسالكيد  
مع زيادة في وزنه . وكذلك تضخم في الكلية اليسرى والطحال دون ظهور أي أعراض على الطائر.  
وتوضح نتائج التجربة بصفة عامة أن إضافة المهدئ إلى مياه الشرب خلال شهر الصيف يعطى  
تأثيراً جيداً على الاتزان الفسيولوجي لدجاج اللحم وخاصة على معدلات النمو ونشاط الغدة الدرقية ودرجة  
حرارة المستقيم وكذلك كيمياء الدم وبالتالي توصى النتائج بإمكانية استخدامه عند ارتفاع درجات الحرارة في  
حالة تربية دجاج اللحم في العنابر المفتوحة.