



Microbes and Infectious Diseases

Journal homepage: <https://mid.journals.ekb.eg/>

Original article

Prevalence, intensity, and risk factors of urinary schistosomiasis among primary school children in Silame, Sokoto, Nigeria

Solomon Matthias Gamde ^{*1,2}, Aminu Garba ³, Princewill Tongywam ², Solomon Dashal Gamde ⁴, Daniel Dansy Agom ⁵, Isaac Igbino ⁵

1- Health Initiative for Safety and Stability in Africa.

2- Department of Medical Laboratory Sciences, Faculty of Health Sciences and Technology, University of Jos, Nigeria.

3- National Health Insurance Scheme Authority, Sokoto State Office, Nigeria.

4- Department of Microbiology, Faculty of Natural Sciences, Abubakar Tafawa Balewa University, Bauchi State, Nigeria.

5- Hospital Services Management Board Sokoto, Nigeria.

ARTICLE INFO

Article history:

Received 20 August 2022

Received in revised form 24 August 2022

Accepted 27 August 2022

Keywords:

Urinary schistosomiasis

Hematuria

Silame

Nigeria

ABSTRACT

Background: Nigeria is the global hotspot for schistosomiasis despite several rounds of school-based preventive chemotherapy because reliable data regarding its geographical distribution are lacking and there is a need to know the current prevalence for the control plans. **Objective:** This study aimed to determine the current prevalence, intensity, and risk factors for urinary schistosomiasis among vulnerable primary school children in Silame, Sokoto, Nigeria. **Methods:** A cross-sectional laboratory-based study was carried out in April 2021 on 188 primary school children in Silame, Sokoto State, Nigeria. Urine samples were collected and examined using the sedimentation technique for the presence of *Schistosoma haematobium* eggs. Hematuria was tested using a urine dipstick. **Results:** Overall, 40 (21.3%) were positive for urinary schistosomiasis with a mean geometric count of 452 eggs/10 ml of urine. Gender was associated ($p=0.0036$) with *Schistosoma haematobium* infection and males 27 (14.4%) recorded higher prevalence than females 13 (6.9%). Hematuria was significantly associated with *Schistosoma haematobium* infection ($p<0.001$) and males (13.8%) recorded higher frequency than females (7.5%). Participants with light intensity of infections 26 (65%) were more than moderate 12 (30%), and heavy infections 2 (5%). Factors significantly associated with *Schistosoma haematobium* infection are gender ($p=0.0036$) and water source ($p<0.001$). **Conclusion:** The present study showed that a significant number of primary school children in Silame suffer from urinary schistosomiasis. Males are at higher risk of infections than females. Strengthening schistosomiasis surveillance systems to identify hotspots, sustainable chemotherapeutic intervention, and improving health education reduces schistosomiasis prevalence.

Introduction

Schistosomiasis caused by *Schistosoma haematobium* (*S.haematobium*) is a public health problem in several parts of the world, especially in tropical and sub-tropical countries characterized by

poverty where there are inadequate supply of safe water, poor sanitation, and unhygienic practices such as open urination. Of the world's 250 million estimated schistosomiasis cases and estimated

280,000 deaths reported annually, it is estimated that at least 93% of individuals affected and requiring treatment for schistosomiasis live in Africa [1-4]. In this region, Nigeria is the major contributor to the schistosomiasis burden with 25.8 million estimated cases and 101.3 million people at risk of the infection [5]. The endemicity of schistosomiasis is attributed to the lack of knowledge of the risks, deprived socio-economic status, inadequate sanitation, and open water activities in snail-infected rivers, streams, and ponds [6].

Schistosomiasis is essentially an infection in the rural and agricultural areas, transmitted by freshwater snails of the genus *Bulinus* where the people's lifestyle promotes the contamination of inland water with human excreta [7]. An individual becomes infected during contact with water containing the infective cercariae of the parasite that penetrate the skin and develops [8,9]. Adult male and female worms are paired together to form a copula within the pelvic venous plexus and the eggs produced are deposited in the bladder wall [10]. Children under 15 years of age remain the most vulnerable and represent the target group for most control interventions [11]. Early symptoms of the disease include haematuria, dysuria, anemia, retardation of growth, and development in children. At an advanced stage, schistosomiasis results in cancer of the bladder, hepatic, and renal malfunctions [12-14].

To date, concerted efforts to control schistosomiasis in Nigeria have failed woefully because reliable data regarding its geographical distribution are lacking due to the lack of political will and poor funding for the control programs [15]. It seems that eliminating schistosomiasis from the country by 2030 is a difficult task. In this context, addressing the underscored issues will escalate the journey towards the goal of schistosomiasis elimination [16]. Therefore, this study aimed to determine the current prevalence, intensity, and risk factors for urinary schistosomiasis among primary school children in Silame, Sokoto, Nigeria.

Materials and methods

Study area

This study was carried out in April 2021 in Silame, covering an area mass of 790 Km² along Sokoto River in Silame Local Government Area, Sokoto State, Nigeria. Sokoto River is located at latitude N13°2'20" and longitude E4°50'35" and provides water for domestic and agricultural uses. The river

is the major source of water for irrigation farming, fish farming, and recreational activities even for the neighboring communities [17].

Study design and population

This is a cross-sectional laboratory-based study was carried out in April 2021 on 188 primary school children attending government primary school Silame, Sokoto State. Age ranged from 5-19 years. The study was conducted according to the international guidelines of Strengthening the Reporting for Observational Studies in Epidemiology (STROBE).

Ethical approval, recruitment, and enrollment

This study was carried out in accordance with the Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments involving human subjects. The protocol was approved by the Sokoto State Ministry of Health, Research Ethics Committee with Ref. No. SLM/PHC/VOL.1/039. The study objectives, procedures, and benefits were explained to the school authorities, parents/guardians, and participants to get their cooperation and consent to conduct the study. A total of 188 primary school children with parental consent were used in the study.

Participants received clean labeled urine containers and were taught how to collect urine samples. A structured questionnaire was administered to each participant to obtain socio-demographic information which was analyzed to determine the associated risk factors for *Schistosoma* infection in the study area [18]. School children who were found to be infected with *S. haematobium* were administered praziquantel drugs provided by the World Health Organization through The Sokoto State Ministry of Health. On the other hand, school children not willing to take part in the study were excluded.


Urine sample collection and processing

Individual-based questionnaires were used to collect information on name, age, sex, water sources, and parental occupation. Urinalysis form was used to record urine parameters while clean, screw-capped, and labeled plastic universal bottles (20 ml) were presented to the respondents who produced terminal urine samples between the hours of 10:00 AM and 2:00 PM [19, 20] which coincided with the circadian rhythm of the parasite egg excretion [21]. In the laboratory, 10ml of each urine sample was measured into a test tube and centrifuged at 5000 rpm for 5 min. The supernatant was decanted off while the sediment was examined under a light microscope

using an x40 objective lens for the presence of terminal spine of *S. haematobium* eggs and expressed as the number of eggs/10ml of urine [22]. The intensity of infection was graded as light

between 1-9 eggs/10ml urine, moderate (10-49 eggs/10 ml urine), and heavy excreting above 50 eggs/10 ml urine. A negative urine sample was indicated by the absence of parasite eggs [1].

Figure 1. GPS map of Silame sampling site.



Statistical analysis

Data obtained from the study were entered into Microsoft Excel 2010 and analyzed with SPSS version 25.0 (SPSS Inc., Chicago, IL, USA). Associations between variables were determined using the Chi-square test and a statistical significance was determined at $p < 0.05$.

Results

Table 1: Sociodemographic characteristics of study participants

Of the 188 urine samples analyzed for urinary schistosomiasis, 51.1% (96/188) were males and 48.9% (92/188) were females, giving a gender ratio of 1:1.044 (males: females). Study participants were primary school children and the majority (65.4%) were of age 5–11 years. Farming is the major occupation of their parents (63.3%). It was obvious that most of the participants depend on boreholes/rivers for daily activities such as bathing and washing.

Table 2: Prevalence of *S. haematobium* among the study participants

The overall prevalence of *S. haematobium* infection was 21.3% (40/188) with a mean egg intensity of 23.1 egg/10 ml of urine. The infections were more prevalent among males 27 (14.4%) than in females 13 (6.9%). The infections were significantly associated with participants' gender ($p=0.036^*$).

Table 3: Age and gender distributions of *S. haematobium* among the study participants

Among the age groups, there was a positive association between the study population and *S. haematobium* infections. The highest frequency of *S. haematobium* infections 26 (13.8%) was recorded among those aged 5 to 11 years with a mean intensity of infections of 8.35 eggs/10 ml (20.9%), while participants aged 12 to 16 years had the least 14 (7.5%) prevalence with a mean intensity of 14.7eggs/10 ml (37.0%). However, the association was not significant. Males had the highest frequency 27 (14.4%) as compared to the females 13 (6.9%). The association was statistically significant ($p=0.036^*$).

Table 4: Age and gender distribution of hematuria among study participants

The frequency of hematuria was higher among those aged 5 to 11 years (13.3%) than older aged 12 to 16 years (7.5%). Hematuria was most frequent among males (13.8%) than females (7.5%).

Table 5: Association between *S. haematobium* infections and hematuria among study participants

Schistosoma haematobium infections are significantly associated with hematuria ($p<0.001^*$) with a mean parasite concentration of 3.0.

Table 6: Risk Factors of schistosomiasis among study participants

Schistosoma haematobium infections are significantly associated with participants' gender and sources of water ($p<0.001^*$). There was no significant association between the father's occupation and the infection ($p=0.75$).

Table 7: Intensity of *S. haematobium* among study participants

Of the 188 samples examined, 40 (21.3%) were found to be positive for the presence of *S. haematobium* eggs. Of the 40 (21.3%) positive, 26 (65%) were excreting between 1-9eggs/10ml, 12 (30%) were excreting between 10-49 eggs/10ml urine, and 2 (5%) were excreting above 50 eggs/10 ml urine.

Table 1. Socio-demographic characteristics among study participants.

Variables	Frequency	Percentage (%)
Age (years)		
5 to 11	123	65.4%
12 to 16	65	34.6%
Gender		
Male	96	51.1%
Female	92	48.9%
Parents occupation		
Crop farmers	119	63.3%
Traders	52	27.7%
Herders	17	9.0%
Water source		
Well water	33	17.6%
River water	10	5.3%
Borehole water	145	77.1%
Education		
Primary school	188	100%

Table 2. Prevalence of *S. haematobium* among study participants.

Variable	Parasite Status – n (%)		Total	P – value
	Negative	Positive		
Gender of the Pupils				
Male	69 (71.9)	27 (14.4%)	96 (100)	0.036 *
Female	79 (85.9)	13 (6.9%)	92 (100)	
Total	148 (78.7)	40 (21.3)	188 (100)	

Table 3. Age and gender distribution of *S. haematobium* among study participants.

Variable	Category	Number of negative <i>S. haematobium</i>	Number of positive <i>S. haematobium</i>	p-value
Age (years)	5 to 11	123 (65.4%)	26 (13.8%)	0.56
	12 to 16	65 (34.6%)	14 (7.5%)	
Gender	Male	69 (36.7%)	27 (14.4%)	0.036 *
	Female	79 (42.0%)	13 (6.9%)	

Table 4. Age and gender distribution of hematuria among study participants.

Variable	Frequency hematuria	Percentage distribution
Age (years)		
5 to 11	25	13.3%
12 to 16	15	7.9%
Gender		
Male	26	13.8%
Female	14	7.5%

Table 5. Association between *S. haematobium* infection and hematuria among study participants

Variable	Mean parasite	
	concentration	<i>p</i> – value
Hematuria		<0.001*
<i>S. haematobium</i>	3.0	

* Significantly associated with parasite status at $p < 0.05$.

Table 6. Risk Factors of schistosomiasis among study participants.

Variable	Category	Number of negative <i>S. haematobium</i>	Number of positive <i>S. haematobium</i>	<i>p</i> -value
Fathers' occupation	Farmers	119 (63.3%)	25 (13.3%)	0.753
	Traders	52 (27.7%)	11 (5.9%)	
	Civil servant	17 (9.0%)	4 (2.1%)	
Water sources	Well water	22 (11.7%)	2 (1.0%)	< 0.001*
	River water	36 (19.2%)	6 (3.2%)	
	Borehole water	130 (69.1%)	32 (17.0%)	
Level of Education	Primary	148 (78.7%)	40 (21.3%)	

Table 7. Intensity of schistosomiasis among study participants

Variable intensity	Age group		
	5-11 years	12-16 years	Total
Light (1-9eggs/10 ml)	18 (45%)	8 (20%)	26 (65%)
Moderate (10-49eggs/10 ml)	7 (17.5%)	5 (12.5%)	12 (30%)
Heavy (50eggs/10 ml)	1 (2.5%)	1 (2.5%)	2 (5%)
Mean Intensity (eggs/10ml)	8.35 (20.9%)	14.7 (37.0%)	23.1 (57.9%)

Discussion

The prevalence of *S. haematobium* infections among primary school children in Silame was 21.3%, a moderate risk prevalence which falls within the national prevalence range of 2 to 82.5% in Nigeria but higher than the national mean of 13%

[23]. The finding of moderate risk prevalence was in tandem with previous studies in Sokoto, North-western Nigeria [24], Kaduna, North-central Nigeria [25], and Maiduguri, North-eastern Nigeria [26]. However, the present study recorded a lower figure of moderate risk prevalence. The decline in the

prevalence of urinary schistosomiasis in the study areas could be attributed to the recent mass distribution of praziquantel drugs among school children [27].

The frequency of *S. haematobium* infection was higher among school age 5-11 years (13.8%) than 12-16 years (7.5%), supporting previous studies in Nigeria [8,28], Cameroon [29], Cote d'Ivoire [30], Swaziland [4], and Malawi [31]. However, the age difference in frequency was not significant. **Chu et al.** [4] and **Nwachukwu et al.** [32] reported the contrary at different locations. The high prevalence of schistosomiasis among younger children might be due to their unrestrained exposure to schistosome-infested water [33].

The frequency of *S. haematobium* infections was higher among males than females with a significant association ($p=0.0036$). This result is in tandem with other studies [19,30,33,34] but different from others that recorded higher prevalence among females [18,35]. Such differences could be traced to the different cultural and religious practices which grant male children unrestrained access to open waters.

Furthermore, the frequency of school children with a light intensity of infections was significantly higher than those with moderate and heavy infections. The high frequency of light intensity of infections in this study was in tandem with findings of **Uneke et al.** [36]; **Otuneme et al.** [18], an indication that most infected individuals harbor low schistosome. However, light-intensity of infections are lethal if not treated. Consistent with **Achanau et al.** [5] and **Degarege et al.** [27] reports, *S. haematobium* infection was associated with hematuria ($p<0.001$). Our findings also confirmed that the dipstick method could be used for rapid screening of *S. haematobium* infection in tandem with a previous report [37].

School children whose water sources for drinking, washing, bathing, and swimming are boreholes and rivers were significantly associated with schistosomiasis. Participants whose parental occupation is farming also showed a positive association with urinary schistosomiasis. This finding is supported by previous studies that associated schistosomiasis infections with cultural and traditional agricultural practices such as washing, fishing, and bathing influences schistosomiasis transmission in Nigeria [38].

Nonetheless, this study has some limitations and the results should be interpreted with care. We conceived that some light-intensity of infections were missed and the overall prevalence of *S. haematobium* might be higher than reported. We also speculate some reporting bias in the questionnaires administered to the participants, no specific information was collected on the hygiene behavior of the participants which are identified risk factors for schistosomiasis.

Conclusion

The present study showed that a significant number of primary school children in Silame suffer from urinary schistosomiasis. Males are at higher risk of infections than females in the study areas. Strengthening surveillance systems to identify schistosomiasis hotspots, sustainable chemotherapeutic intervention, and improving health education are recommended to reduce schistosomiasis prevalence.

Acknowledgments

We thank the Ministry of Health, Sokoto State for the ethical clearance, the principal of Government Primary School Silame for assisting during data collection, and the study participants for their cooperation during specimens and data collection.

Availability of data and materials

Data are available on reasonable request from the corresponding author.

Conflict of interest : None

Funding

Nothing to declare

References

- 1-**Siddiqui AA, Siddiqui SZ.** Sm-p80-based schistosomiasis vaccine: preparation for human clinical trials. Trends Parasitol 2017; 33(3):194–201.
- 2-**Frahm S, Anisuzzaman A, Prodjinotho UF, Vejzagić N, Verschoor A, Prazeres da Costa C.** A novel cell-free method to culture *Schistosoma mansoni* from cercariae to juvenile worm stages for in vitro drug testing. PLoS Negl Trop Di 2019; 13(1):e0006590.
- 3-**Anisuzzaman, Tsuji N.** Schistosomiasis and hookworm infection in humans: disease burden,

- pathobiology, and anthelmintic vaccines. *Parasitol Int* 2020;102051.
- 4-**Chu TB, Liao CW, D’Lamini P, Chang Peter WS, Chiu WT, Du WY, et al.** Prevalence of *Schistosoma haematobium* infection among inhabitants of Lowveld, Swaziland, an endemic area for the disease. *Tropical Biomedicine* 2010; 27(2): 337–342.
- 5-**Anchau ZG, Silas PC, Shaibu MA.** Prevalence of *Schistosoma haematobium* among Qur’anic School Pupils in Zaria, Kaduna State. *UJMR* 2020; 5(2): 92 – 97.
- 6-**Ezeh CO, Onyekwelu KC, Akinwale OP, Shan L, Wei H.** Urinary schistosomiasis in Nigeria: A 50-year review of prevalence, distribution, and disease burden. *Parasite* 2019; 26:19.
- 7-**Balla HJ, Zailani SB, Askira MM, Musa AB, Mursal A.** Prevalence of urinary schistosomiasis amongst “Almajiris” and Primary School Pupils in Gwange Ward of Maiduguri. *Borno Medical Journal (BOMJ)* 2010; 7(2):7-10.
- 8-**Mohammed K, Suwaiba M, Spencer HI, Nataala SU, Ashcroft OF, Nuhuand A, et al.** Prevalence of Urinary Schistosomiasis among Primary School Children in Kwankwalawa Area, Sokoto State, North-Western Nigeria. *Asian Journal of Research in Medical and Pharmaceutical Sciences* 2018; 3(1): 1-10.
- 9-**World Health Organization (WHO).** “Ending the neglect to attain the Sustainable Development Goals: a road map for neglected tropical diseases” World Health Organization; 2020. 2021–2030. Geneva. Available at: <https://www.who.int/publications/i/item/9789240010352>.
- 10-**Oluwafemi FS, Oluwadare T, Fasuba B, Faeji CO, Kukoyi O, Oni IO, et al.** Prevalence of Urinary Schistosomiasis Among School Children In Ago Aduloju Community, Ado LGA of Ekiti State, Nigeria. *Nigerian Journal of Parasitology* 2022; 43(1) 135-144.
- 11-**Olamiju F, Nebe OJ, Mogaji H, Marcus A, Amodu–Agbi P, Urude RO, et al.** Schistosomiasis outbreak during COVID-19 pandemic in Takum, Northeast Nigeria: Analysis of infection status and associated risk factors. *PLoS. One* 2022; 17(1): e0262524.
- 12-**Casmo V, Augusto G, Nala R, Sabonete A, Carvalho-Costa FA.** The effect of hookworm infection and urinary schistosomiasis on blood hemoglobin concentration of schoolchildren living in northern Mozambique. *Rev Inst Med Trop Sao Paulo* 2014; 56(3):219–24.
- 13-**Deribew K1, Tekeste Z, Petros B, Huat LB.** Urinary *schistosomiasis* and malaria-associated anemia in Ethiopia. *Asian Pac J Trop Biomed* 2013; 3(4):307–310.
- 14-**Atalabi TE, Lawal U, Ipinlaye SJ.** Prevalence and intensity of genito-urinary *schistosomiasis* and associated risk factors among junior high school students in two local government areas around Zobe Dam in Katsina State, Nigeria. *Parasites & Vectors* 2016; 9:388, 1-12.
- 15-**Adedeji AS, Okechukwu O, Adenike OF, Abolore AH, Kayode OS, Kayode OG.** Original Article Prevalence and risk factors of schistosomiasis infection among primary school pupils in Patigi Local Government Area, Kwara State, Nigeria. *Sudan Med J* 2017; 53(1); 1-13.
- 16-**Mushi V, Zacharia A, Shao M, Mubi M, Tarimo D.** Persistence of *Schistosoma haematobium* transmission among school children and its implication for the control of urogenital schistosomiasis in Lindi, Tanzania. *PLoS ONE* 2022; 17(2): e0263929.
- 17-**Yusha’u MA, Tsafe AK, Babangida SI, Lawal NI.** Problems and prospects of

- Integrated Almajiri Education in Northern Nigeria. *Scientific Journal of Pure and Applied science* 2013; 2(3), 125-134.
- 18-**Otuneme OG, Obebe OO, Sajobi TT, Akinleye WA, Faloye TG.** Prevalence of Schistosomiasis in a neglected Community, southwestern Nigeria at two points in time, spaced three years apart. *Afri Health Sci* 2019; 19(1): 1338-1345.
- 19-**Geleta S, Alemu A, Getie S, Mekonnen Z, Erko B.** Prevalence of urinary *schistosomiasis* and associated risk factors among Abobo Primary School children in Gambella Regional State, Southwestern Ethiopia: a cross-sectional study. *Parasit Vectors* 2015; 8 (1):1106-1114.
- 20-**Atalabi TE, Adoh SD, Eze KM.** The current epidemiological status of urogenital schistosomiasis among primary school pupils in Katsina State, Nigeria: An imperative for a scale-up of water and sanitation initiative and mass administration of medicines with Praziquantel. *PLoS Neglected Tropical Diseases* 2018; 12: 36.
- 21-**Ekpo UF, Laja-Deile A, Oluwole AS, Sam-Wobo SO, Mafiana CF.** Urinary schistosomiasis among preschool children in a rural community near Abeokuta, Nigeria. *Parasit Vectors* 2010; 3:58.
- 22-**Cheesbrough M.** "District Laboratory Practice in Tropical Countries" Part 2 Second Edition. Cambridge University Press, 2006.
- 23-**Uchen O, Oladoyin V, Idowu M, Adeyera O, Olabisi O, Oluwatosin O, et al.** Urinary schistosomiasis among vulnerable children in a rehabilitation home in Ibadan, Oyo state, Nigeria *Obioma. BMC Infect. Dis* 2017; 17:487.
- 24-**Solomon M. Gamde, Ahmed M. Ganau, Jimoh A. Abdulahi, Princewill J. Tongvwam, Dashed S. Gamde.** Prevalence and risk factors of urinary schistosomiasis amongst Almajiri pupils in Silame, Sokoto State, North-western Nigeria. *Umaru Musa Yar'adua University Journal of Microbiology Research* 2021; 6(1): 1-10.
- 25-**Omenesa HO, Bishop HG, Raji HM.** Prevalence of urinary schistosomiasis among pupils attending primary schools in Bomo village, Zaria- Nigeria. *International Journal of Research in Engineering and Science* 2015; 3(5):14-19.
- 26-**Balla HJ, Babagana DR, Baba S, Ibrahim H.** Incidence of Urinary Schistosomiasis amongst Out-of-School Pupils and "Almajiris" in Dikwa, North-Eastern Nigeria. *Global Journal of Medical Research, Microbiology, and Pathology* 2015; 15 (2): 9-13.
- 27-**Degarege A, Mekonnen Z, Levecke B, Legesse M, Negash Y, Vercruysse J, et al.** Prevalence of *Schistosoma haematobium* Infection Among School-Age Children in Afar Area, Northeastern Ethiopia. *PLoS ONE* 2015 10(8): e0133142.
- 28-**Sarkinfa F, Oyebanji AA, Sadiq IA, Ilyasu Z.** Urinary schistosomiasis in the Danjarima community in Kano, Nigeria. *J Infect Dev Ctries* 2009; 3(6):452-457.
- 29-**Sama MT, Oyono E, Ratard RC.** High-risk behaviors and *schistosomiasis* infection in Kumba, South-West Province, Cameroon. *Int J Environ ResPublic Health* 2007; 4(2):101-105.
- 30-**Angora EK, Boissier J, Menan H, Rey O, Tuo K, Touré AO, et al.** Prevalence and Risk Factors for Schistosomiasis among Schoolchildren in two Settings of Côte d'Ivoire. *Trop Med Infect Dis* 2019; 4: 110.
- 31-**Chipeta MG, Ngwira B, Kazembe LN.** Analysis of *schistosomiasis haematobium* infection prevalence and intensity in Chikhwawa, Malawi: An application of a two-

- part model. PLoS Negl Trop Dis 2013; 7(3): e2131.
- 32-Nwachukwu PC, Ohaeri CC, Ukpai OM, Irole-eze OP, Amaechi EC. Prevalence of *Schistosoma haematobium* infection among school-aged children in Afikpo North local government area, Ebonyi State, Nigeria. Sri Lankan J Biol 2018; 3(2):1.
- 33-Awosolu OB, Akinnifesi OJ, Salawu AS, Omotayo YF, Obimakinde ET, Olise C. Prevalence and intensity of urinary schistosomiasis among school-age children in Ikota, Southwestern Nigeria. Braz. J. Biol. Sci 2019; 6: 391–399.
- 34-Ahmed AM, Abbas H, Mansour FA, Gasim GI, Adam I. *Schistosoma haematobium* infections among schoolchildren in central Sudan one year after treatment with praziquantel. Parasites & Vectors 2012 5:108.
- 35-Hassan AO, Amoo AOJ, Akinwale OP, Deji-Agboola AM, Adeleke MA, Gyang PV. Human water contact activities and urinary schistosomiasis around Erinle and Eko-Ende dams. Glob Adv Res J Med Med. Sci 2012; 1: 77-84.
- 36-Uneke C, Oyibo P, Ugwuoru C, Nwanokwai A, Iloegbunam R. Urinary *Schistosomiasis* among School Age Children in Ebonyi State, Nigeria. The Internet Journal of Laboratory Medicine 2007; 2 (1).
- 37-King CH1, Bertsch D. Meta-analysis of urine heme dipstick diagnosis of *Schistosoma haematobium* infection, including low-prevalence and previously-treated populations. PLOS Negl Trop Dis 2013; 12:7:e2431.
- 38-Amuta EU, Houmsou RS, "Prevalence, the intensity of infection and risk factors of urinary *schistosomiasis* in pre-preschool and school-aged children in Guma Local Government Area, Nigeria. Asian Pac. J Trop Med 2014; 34-39.