**Original Paper****Prevalence of *Campylobacter jejuni* in fresh meat products**Ahmed M. AboShady¹, Ashraf A. Abd El-Tawab¹, Fatma I. El-Hofy¹, Mona M. Sobhy², Enas A. Soliman¹¹ Department of Bacteriology, Immunology and Mycology, Faculty of Veterinary Medicine, Benha University, Egypt² Department of Reproductive Diseases, Animal Reproduction Research Institute, Agriculture Research Centre (ARC), Cairo, Egypt.**ARTICLE INFO****Keywords**

C. jejuni
Fresh meat products
PCR

Received 11/11/2022

Accepted 04/01/2023

Available On-Line

15/01/2023

ABSTRACT

Campylobacter jejuni (*C. jejuni*) is one of great significance to public health as it is the main source of food borne *Campylobacter* enteritis in human. Raw and undercooked contaminated meat products are known as important sources of human campylobacteriosis. Contamination of meat can occur in different steps during production as preparation, processing, distribution, marketing and handling at transportation. The main object of study was to isolated and identified of *C. jejuni* in from fresh meat by conventional methods and confirmed by polymerase chain reaction (PCR). Sixty one samples were collected from fresh meat including minced meat (15), liver (14), meat (25) and sausage Baladi (7). The prevalence rate of *C. jejuni* in fresh meat by PCR was 32.7% compared with conventional methods (19.67%). Polymerase chain reaction targeting hip *O* gene specific for *C. jejuni* was used for phenotypically identified *C. jejuni* isolates. This study concluded that PCR was more specific and rapid than the conventional methods for identification of *C. jejuni* in fresh meat. Raw retail meats need control programs and consumer food safety education efforts by application of hygienic measuring.

1. INTRODUCTION

C. jejuni is a gram-negative bacterium. It is considered one of the most important from both a microbiological and public health perspective (Ryan and Ray, 2004), can cause infection and gastroenteritis (Brownsell *et al.*, 1989). *Campylobacter* species are widespread in most warm-blooded animals. They are prevalent in food animals such as poultry, cattle, pigs, sheep, ostriches, and shellfish as well as pets, including cats and dogs (World Health Organization, 2011). Consumption of contaminated or undercooked meat (especially poultry) is the major route of transmission in humans, (Center for Food Security and Public Health, 2013). Contamination of meat can occur in different steps along the food production chain including production, processing, distribution, marketing and handling or preparation (Zhao *et al.*, 2001). Thermophilic *Campylobacters*; *C. jejuni*, *C. coli*, *C. upsaliensis* and *C. lari* implicated as food borne infections (Iovine *et al.*, 2008). *C. jejuni* is the most of the food borne enteritis in human, followed by *C. coli* and lesser extent *C. lari* (Skirrow and Blaser, 2000). *C. jejuni* has been frequently reported as a cause of human campylobacteriosis (80-90%) compared to *C. coli* (5-10%) (EFSA, 2008)

Detection of thermophilic *Campylobacter* species based upon performing catalase test, susceptibility to nalidixic acid and cephaloxin and rapid hippurate hydrolysis test. Most of *Campylobacter* isolates were resistant to nalidixic acid, so that it is used in the distinction between *C. jejuni*, *C. lari* and *C. coli* (Megraud, 1987). Hippurate hydrolysis is the only phenotypic exam differentiating *C. jejuni* from other species of *Campylobacters*, especially the

thermophilic species. *C. jejuni* is capable of hydrolyzing sodium hippurate to benzoic acid and glycine (ISO, 2006). The main object of this study was to isolate and identify of *C. jejuni* from fresh meat by conventional methods and confirmed by polymerase chain reaction (PCR).

2. MATERIAL AND METHODS

Our study was performed during the period between August 2017 and January 2019 in Reproductive Diseases Department, Animal Reproduction Research Institute, ARC, El Haram, Giza

2.1. Samples

A total of 61 samples gathered from fresh meat products including minced meat (15), liver (14), meat (25) and sausage Baladi (7) from many supermarkets and slaughterhouses in Cairo Province

2.2. Culturing of prepared samples

All samples were homogenized and inoculated in thioglycolate broth then incubated at 42 ° C for 48 hrs under microaerobic condition (5% O₂, 10% CO₂ and 85% N₂) (Gebhart *et al.* 1985).

A loopful from each sample were cultured directly on thioglycolate broth medium and then sub-cultured in modified *Campylobacter* blood free selective medium. The plates were incubated in microaerophilic condition (CO₂ (10%), O₂ (5%) and nitrogen (85%)) at 37°C for 48 hours (Smibert, 1974).

2.3. Biochemical identification of *Campylobacter* colonies
Identification of suspected colonies was done depending on characteristics colony morphology, Gram stain Koneman *et al.*, (1995), motility test (Smibert, 1974) and biochemical tests (Catalase production test, Nitrate reduction test, Oxidase test, Urease test, H₂S production test with lead acetate paper, Temperature tolerance test, Glycine tolerance test, NaCl tolerance test, Hippurate hydrolysis test, sensitivity to cephaloxcin and nalidixic acid) Al-Gohary, (1998).

2.4. Molecular identification by PCR

Campylobacter DNA was extracted by using (Thermo Scientific Gene Jet Genomic DNA Purification Kit#K0721, #K0722). Species-specific primer targeting *hip O* gene specific for *C. jejuni* (Wang *et al.*, 2002) were 5`ACTTCTTTATTGCTTGCTGC3` (forward) and 5`GCCA-CAACAAGTAAAGAAGC3` (reverse). PCR amplification was done by using thermal cycler (Biometra) (Wang *et al.*, 2002). 1.5% agarose gel (Biometra) was used for electrophoresis of the amplified PCR products. The gel was photographed by Alpha Innotech system (El-Adawyet *et al.*, 2012).

3. RESULTS

Bacteriological examination of collected samples revealed isolation of *C. jejuni* from 12/61 (19.67%) fresh meat product samples. They were isolated from minced meat (5/15) (33.3%), meat muscles (5/25) (20%), sausage baladi (1/7) (14.3%) and liver (1/14) (7.1%). The samples confirmed by molecular PCR targeting *hip O* gene specific for *C. jejuni*. The occurrence of *C. jejuni* in fresh meat by PCR was 20/61 (32.7%) compared with conventional methods 12/61 (19.67%). The highest percentage for isolation of *C. jejuni* was in minced meat 33.3% by CM and 46.6% by PCR, the isolation rate of *Campylobacter jejuni* from sausage was 14.3% by CM and 28.6% by PCR. The results of *C. jejuni* in meat muscles were 20% by CM and 28% by PCR but, the incidence of *C. jejuni* in liver was 7.1% by C.M. and 28.6% by PCR (Table, 1 and Fig. 1).

Table (1): Incidence of *C. jejuni* in fresh meat product samples by CM and PCR

Type of Samples	No. of Samples	Fresh meat samples			
		Positive by CM		Positive by PCR	
		No	%	No	%
Minced meat	15	5	33.3%	7	46.6%
Liver	14	1	7.1%	4	28.6%
Meat	25	5	20%	7	28%
Sausage Baladi	7	1	14.3%	2	28.6%
Total	61	12	19.67%	20	32.7%




Fig (1): Electrophoresis profile of *C. jejuni* producing amplicons of average size 735 bp for *hip* gene, lane 3 positive control *Campylobacter jejuni* (ATCC 33291)

4. DISCUSSION

Campylobacters are one of the most important food bacteria leading to gastroenteritis in humans in developed and developing countries (Rahimi and Ameri, 2011).

Campylobacter jejuni causes more than 90% of the reported *Campylobacter* infections (NARMS, 2010). *C. jejuni* is distinct by rapid Sodium hippurate hydrolysis test which is the only phenotypic test differentiating *C. jejuni* from other species of *Campylobacters*, especially thermophilic species (ISO, 2006). Moreover, PCR targeting *hip O* (benzoylglycineamidohydrolase) has been used to confirm the detection and discrimination of *C. jejuni*. The gene *hip O* is specific for the hippurate activity and differentiates *C. jejuni* from other *Campylobacter* species (Englen *et al.*, 2003).

In this current study, the overall occurrence of *Campylobacter jejuni* was 19.67% (12/61) fresh raw meat samples by conventional methods while by PCR, its prevalence was 32.7% (20/61). The largest incidence rate of *C. jejuni* was identified in minced meat (5/15) 33.3%, followed by meat (5/25) 20%, Baladi sausage (1/7) 14.3% and liver (1/14) 7.1% (Table, 1).

Nearly similar prevalence rate of *C. jejuni* from beef meat obtained at retail outlets was 22% of examined samples (Datta *et al.*, 2003). Also, the prevalence of *Campylobacter* SPP. isolate was 16.66% in beef meat, 26.66% in beef liver and 22% in pediatric diarrhea. *C. jejuni* were 60% higher than *C. coli* (Parkhill *et al.*, 2000). Elgabry *et al.*, (2016) recorded 16% (117/733) from fast meat meals (shawarma, kofta, hamburger, sausage, offal, and liver) in five Egyptian governorates. The distribution of *C. jejuni* from fast meat meals was 25.3% (19/75) in Assuit, 16.8% (28/167) in Fayuom, 15.3% (26/170) in Qalubia, 14.7% (30/204) in Cairo, and 12% (14/117) in Bin-suef (Elgabry *et al.*, 2016).

Lower isolation rate of *C. jejuni* from beef meat was obtained in Great Britain; 1.6% from meat samples (Turnbull PC and Rose P. 1982), U.S. Washington; 4.7% (6/129 bovine meat) (Stern *et al.*, 1984), Poland; 0% (0/114 beef) (Kwiatk *et al.*, 1990), Assuit; 12% (3/25 slaughtered buffalo) (Refaie and Galal, 1991), the Greater Washington, D.C., area, including suburban Maryland; 0.5% (1/182 beef samples) (Zhao *et al.*, 2001), Istanbul, Turkey; 3% (6/198 beef samples from carcasses) (Bostan *et al.*, 2009), Northern Poland; 10.1% (35/347 beef meat) (Andrzejewska *et al.*, 2019), Italy; 0; 24% (3/1203 bovine meat samples (689 hamburgers and 514 knife-cut meat preparations) (Giannatale *et al.*, 2019) Africa; 4% (21/521 cattle meat) 4.47% (37/827 cattle carcass) (Thomas KM *et al.*, 2020), Mekelle, Ethiopia; 9% (19/210 cattle meat) (Hagos *et al.*, 2021), and in Belgium beef carcasses and cutting meat (3.3% and 5.0%, respectively) (Ghafir *et al.*, 2007). Several studies recorded absence of *Campylobacter* isolation in examined samples. As, examination of 133 retail beef samples, consisting of ground beef and whole muscle steaks and roasts in a rural Midwest city (Fargo) (Kegode *et al.*, 2008); retail ground beef samples (n ~ 100) in Alberta, Canada (Bohaychuk, *et al.*, 2006); a survey on beef carcasses conducted in 10 abattoirs in Northern Ireland (Madden *et al.*, 2001) and from veal carcasses in Belgium (Ghafir *et al.*, 2007).

However, higher findings of *C. jejuni* from beef meat were reported in Ghana 29.1% (32/110 cattle carcasses) (Karikari *et al.*, 2017), Ottawa, Ontario 50% (50/100). The distribution form of *C. jejuni* -positive animals, in decreasing order, was steers (55%) (42/76), bulls (40%) (4/10), heifers (40%) (2/5), and cows (22%) (2/9) (Garcia *et al.*, 1985), Belgium 100% of isolates were *C. jejuni* (5/5 beef meat) between 2000 and 2003, (Ghafir *et al.*, 2007).

The occurrence rate of *C. jejuni* in beef livers in our study was 7.1% (1/14 beef liver) by conventional methods and

28.6% (4/14 beef liver) by PCR. That was higher than reported in Brazil; 1.51% (2/132 chilled beef liver) (Takeuchi et al., 2022), and Turkey; 0%(0/325 Liver samples from apparently healthy cattle by PCR) (Açik and Çetinkaya, 2005). From the other hand, our isolation rate was lower than the previous studies conducted in Washington, D.C.; 15%(6/40 fresh beef liver) (Stern et al., 1984), Ottawa, Ontario; 12%(12/100) (Garcia et al., 1985), Assuit Governorate; 8%(2/25 fresh liver samples) (Refaie and Galal., 1991), Toukh, Kaliobia governorate; 16.66% (5/30 beef liver) (Khalifa, Nashwa. (2013)), Egypt; 18.2% (26/143 offal and liver) (Elgabry et al., 2016), 30% (3/10) from offal and liver was in Assuit, 22.9%(8/35) in Qaluobia, 21.2% (7/33) in Fayuom, 12.8%(5/39) in Cairo and 11.5%(3/26) in Bin-suef (Elgabry et al., 2016), and Africa; 16.66%(5/30 cattle liver) (Thomas et al., 2020). Garcia et al., (1985) reported 12% (12/100) from the examined liver samples (Garcia et al., 1985). Noor mohamed and Fakhr, (2013) reported the overall prevalence of *C. jejuni* in beef livers was 26% (13/50 beef livers) and 2/50 (4%) of the samples was contaminated with both *C. jejuni* and *C. coli*. Also, Ghafir et al., (2007) suggested that the highest *C. jejuni* recovery from livers probably because the liver surface stays moist suitable for foodborne pathogen. Fecal contamination of *C. jejuni* by slaughtered cows is a possible source of contaminated beef liver.

Although the overall prevalence of *C. jejuni* in minced meat in the present investigation was 33.3% (5/15) by conventional methods and 46.6% (7/15) by PCR, no positive result was found from minced meat from beef and veal in Belgium. Pork and beef minced meat were contaminated with *Campylobacter* at a very low level: on average, 2.5% and 0.6% in 25 g samples, respectively in Belgium (Ghafir et al., 2007). Furthermore, *C. jejuni* was not isolated from chilled minced meat 0% (0/138) in Brazil (Takeuchi et al., 2022). Moreover, only in one sample minced beef were both *Salmonella* and *Campylobacter* found together (Turnbull PC and Rose P. 1982). On the other hand; *C. jejuni* was isolated from all samples of minced meat. It is a common or regular finding in minced meat for sale in ordinary grocery stores (Svedhem et al., 1981). The isolation rate of *C. jejuni* in the current study in sausage baladi was 14.3% (1/7) by conventional methods and 28.6% (2/7) by PCR. That was in line with other study recorded by Elgabry et al., (2016) in Egypt, who reported prevalence rate 15.2% (27/178 sausage) from five Egyptian governorates. The highest incidence rate was detected in Assuit 20.8%(5/24) followed by 15.4%(6/39) in Fayuom, 15.2%(7/46) in Cairo, 13.2%(5/38) in Qaluobia and 12.9%(4/31) in Bin-Suef.

The maximum identification percentages of *C. jejuni* by PCR was in minced meat (7 /15) 46.6% followed by sausage (3/9) 33.3% then, liver (4 /14) 28.6%, and meat muscles (7/25) 28% (Table, 1 and Fig. 1).

The fluctuation in *Campylobacter* species isolation rate in various studies have been attributed to many reasons as, type of examined samples, location, climate factors, hygienic measures and isolation as well as identification techniques (Jorgensen et al., 2011 and Chatur et al., 2014). In addition to technical differences in protocols experiment, various other factors as size, breed of herd, season, animal ages, geography, or husbandry practices (Stanley and Jones, 2003 and Hakkinen and Hanninen, 2009).

5. CONCLUSION

Finally, it is concluded that the examined fresh meat was contaminated by *C. jejuni*. Consumption of undercooked or

cooked contaminated meat products is considered a possible risk factor of human campylobacteriosis. Raw retail meats are potential vehicles for transmitting foodborne diseases, need control programs and consumer food safety education efforts by applied hygienic measuring, continued applying of HACCP systems, and increased consumer food safety at home.

CONFLICT OF INTEREST

The authors declare that they have no conflicts of interest for current data

6. REFERENCES

1. Açik, M. and Çetinkaya, B. 2005. The heterogeneity of *Campylobacter jejuni* and *Campylobacter coli* strains isolated from healthy cattle. Letters in Applied Microbiology 41, 397-403.
2. Andrzejewska, M., Szczepańska, B., Śpica, D., and Klawe, J. J., 2019. Prevalence, Virulence, and Antimicrobial Resistance of *Campylobacter* spp. in Raw Milk, Beef, and Pork Meat in Northern Poland. Foods (Basel, Switzerland), 8(9), 420.
3. Bohaychuk, V.M., Gensler, G.E., King, R.K., Manninen, K.I., Sorensen, O., Wu, J.T., Stiles, M.E., McMullen, L.M. et al., 2006. Occurrence of pathogens in raw and ready-to-eat meat and poultry products collected from the retail marketplace in Edmonton, Alberta, Canada. J Food Prot. 69(9), 2176-2182.
4. Brownsell, V.L., C.J. Griffith and E. Jones, 1989. Applied Science for Food Studies. Longman Scientific and Technical, Hong Kong.
5. Center for Food Security and Public Health 2013. Zoonotic *Campylobacteriosis*, *Campylobacter* Enteritis, *Vibronic* Enteritis, *vibriosis*. [http:// www.cfsph.iastate.edu/IICAB/](http://www.cfsph.iastate.edu/IICAB/).
6. Chatur, Y.A., Brahmbhatt, M.N., Modi, S., Nayak, J.B. et al., 2014. Fluoroquinolone resistance and detection of topoisomerase gene mutation in *Campylobacter jejuni* isolated from animal and human sources. Int. J. Curr. Microbiol. App. Sci., 3(6), 773-783.
7. Dadi, Lemma & Asrat, Daniel. et al., 2009. Prevalence and antimicrobial susceptibility profiles of thermotolerant *Campylobacter* strains in retail raw meat products in Ethiopia. Ethiopian Journal of Health Development. 22. 10.4314/ejhd.v22i2.10072.
8. Datta, S., Niwa, H., and Itoh, K. 2003. Prevalence of 11 pathogenic genes of *Campylobacter jejuni* by PCR in strains isolated from humans, poultry meat and broiler and bovine feces. Journal of Medical Microbiology 52(4), 345-348.
9. EFSA, 2010. The community summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in the European Union in 2008. EFSA Journal 8(1), 1496.
10. El-Adawy, H., Hotzel, H., Tomaso, H., Neubauer, H., and Hafez, H.M., 2012. Elucidation of colonization time and prevalence of thermophilic *Campylobacter* species during turkey rearing using multiplex polymerase chain reaction. Poultry Sci.; 91(2):454-9.
11. Elgabry, A., Barakat, Ashraf, El-Razik, Khaled, Azza, S and Hassan, A., 2016. Incidence of zoonotic *Campylobacter jejuni* in fast meal meat, grill chickens and symptomatic Egyptians. African Journal of Microbiology Research 10, 608-615.
12. El-Gohary, A.H., (1998). Prospective studies on campylobacteriosis in human and animals in contact. Assiut Vet. Med. J., 38, 192-202.
13. Englen, M.D., Ladely, S.R., Fedorka-Cray, P.J., 2003. Isolation of *Campylobacter* and identification by PCR. Methods Mol. Biol., 216, 109-21.
14. Garcia, M.M., Lior, H., Stewart, R.B., Ruckerbauer, G.M., Trudel, J.R.R. and Skljarevski, A. 1985. Isolation, characterization, and serotyping of *Campylobacter jejuni* and *Campylobacter coli* from slaughter cattle. Applied and Environmental Microbiology 49, 667-672.
15. Gebhart, C., Edmonds, P., Word, G., Kurtz, H. and Bernner, D., 1985. *Campylobacter* hyointestinalis spp: A new species of *Campylobacter* found in the intestine of pigs and other animals. J. Clin. Microbiol. 21, 715-720.
16. Ghafir, Y., B. China, K. Dierick, L. De Zutter and G. Daube, 2007. A seven-year survey of *Campylobacter* contamination in

- meat at different production stages in Belgium. *Int. J. Food Microbiol.*, 116, 111-120.
17. Di Giannatale, E., Calistri, P., Di Donato, G., Decastelli, L., Goffredo, E., Adriano, D., Mancini, M.E., Galleggiante, A., Neri, D., Antoci, S., Marfoggia, C., Marotta, F., Nuvoloni, R., Migliorati, G. et al., 2019. Thermotolerant *Campylobacter* spp. in chicken and bovine meat in Italy: Prevalence, level of contamination and molecular characterization of isolates. *PLoS One*.14(12):e0225957. Published 2019 Dec 6. doi:10.1371/journal.pone.0225957.
 18. Hagos, Y., Gugsu, G., Awol, N., Ahmed, M., Tsegaye, Y., Abebe, N., and Bsrat, A. 2021. Isolation, identification, and antimicrobial susceptibility pattern of *Campylobacter jejuni* and *Campylobacter coli* from cattle, goat, and chicken meats in Mekelle, Ethiopia. *PLoS one*, 16(2), e0246755. <https://doi.org/10.1371/journal.pone.0246755>.
 19. Iovine, N.M., Pursnani, S., Voldman, A., Wasserman, G., Blaser, M.J., Weinrauch, Y., 2008. Reactive nitrogen species contribute to innate host defense against *C.jejuni*. *Infection and immunity* 76, 986-993.
 20. ISO, 2006. Microbiology of Food and Animal Feeding Stuffs – Horizontal Method for Detection and Enumeration of *Campylobacter* spp. Part 1: Detection Method. Geneva: International Organization for Standardization. [ISO 10272-1:2006.]
 21. Jorgensen, F., Ellis-Iversen, J., Rushton, S., Bull, S.A., Harris, S.A., Bryan, S.J., Gonzalez, A., Humphrey, T.J. 2011. Influence of Season and Geography on *Campylobacter jejuni* and *C. coli* Subtypes in Housed Broiler Flocks Reared in Great Britain. *Appl. Environ. Microbiol.*, 77(11),3741-3748.
 22. Karikari, A. B., Obiri-Danso, K., Frimpong, E. H., and Krogfelt, K. A., 2017. Antibiotic Resistance of *Campylobacter* Recovered from Faeces and Carcasses of Healthy Livestock. *BioMed Research International*, 2017, [4091856].
 23. Kashoma, I. P., Kassem, I. I., John, J., Kessy, B. M., Gebreyes, W., Kazwala, R. R., and Rajashekara, G. 2016. Prevalence and Antimicrobial Resistance of *Campylobacter* Isolated from Dressed Beef Carcasses and Raw Milk in Tanzania. *Microbial drug resistance (Larchmont, N.Y.)*, 22(1), 40–52.
 24. Kegode, R.B., Doetkott, D.K., Khaitsa, M.L. and Wesley, I.V. 2008. Occurrence of *Campylobacter* species, *Salmonella* species and generic *Escherichia coli* in meat products from retail outlets in the Fargo metropolitan area. *J. Food Safety*, 28: 111-125.
 25. Nashwa O. Khalifa, Jehan S.A. Afify and Nagwa S. Rabie 2013. Zoonotic and Molecular Characterizations of *Campylobacter jejuni* and *Campylobacter coli* Isolated from Beef Cattle and Children. *Global Veterinaria* 11 (5): 585-591.
 26. Koneman, E.W., Allen, S.D., Janda, W.M., Schreckenberger, P.C., and Winn, W.C., 1995. *Introduction to diagnostic Microbiology* 5thed. Lippincott. Company, Philadelphia, (USA).
 27. Korsak, D., Maćkiw, E., Rożynek, E., and Żyłowska, M., 2015. Prevalence of *Campylobacter* spp. in Retail Chicken, Turkey, Pork, and Beef Meat in Poland between 2009 and 2013. *Journal of food protection*, 78(5), 1024–1028.
 28. Kramer, J. M., Frost, J. A., Bolton, F. J., and Wareing, D. R. 2000. *Campylobacter* contamination of raw meat and poultry at retail sale: identification of multiple types and comparison with isolates from human infection. *Journal of food protection*, 63(12), 1654–1659.
 29. Kwiatek, K., Wojton, B., & Stern, N. J. 1990. Prevalence and Distribution of *Campylobacter* spp. on Poultry and Selected Red Meat Carcasses in Poland. *Journal of food protection*, 53(2), 127–130.
 30. Madden, R. H., Espie, W. E., Moran, L., McBride, J., and Scates, P. 2001. Occurrence of *Escherichia coli* O157:H7, *Listeria monocytogenes*, *Salmonella* and *Campylobacter* spp. on beef carcasses in Northern Ireland. *Meat science*, 58(4), 343–346.
 31. Megraud, F., 1987. Diagnostic bacteriologique des infections à *Campylobacter*. *Rev. Fr. Lab.*, 156:2-16.
 32. Mehdizadeh, M., Rahimifard, N., Shoeibi, S., Mirsalehian, A., Saadati, S., Noori, Z., Pirali- Hamedani, M., 2009. Detection of *Campylobacter jejuni* in raw meat. *Iran J Microbiol.* 1;1(3):43-44.
 33. NARMS, National Antimicrobial Resistance Monitoring System for Enteric Bacteria 2010. Human Isolates Final Report. U.S. Department of Health and Human Services, CDC. (Accessed at <http://www.cdc.gov/narms/reports.htm>).
 34. Noormohamed, A., and Fakhr, M. K., 2013. A higher prevalence rate of *Campylobacter* in retail beef livers compared to other beef and pork meat cuts. *International journal of environmental research and public health*, 10(5), 2058–2068.
 35. Parkhill, J., Wren, B. W., Mungall, K., Ketley, J.M., Churcher, C., Basham, D., Chillingworth, T., Davies, R.M., Feltwell, T., Holroyd, S. and Jagsels K., 2000. The genome sequence of the food-borne pathogen *Campylobacter jejuni* reveals hyper variable sequences. *Nature* 403,665–668.
 36. Rahimi, E., Ameri, M., and Kazemeini, H. R. 2010. Prevalence and antimicrobial resistance of *Campylobacter* species isolated from raw camel, beef, lamb, and goat meat in Iran. *Foodborne pathogens and disease*, 7(4), 443–447.
 37. Rahimi, E., Ameri, M., and Kazemeini, H. R. 2010. Prevalence and antimicrobial resistance of *Campylobacter* species isolated from raw camel, beef, lamb, and goat meat in Iran. *Foodborne pathogens and disease*, 7(4), 443–447.
 38. Rahimi, E. and Ameri, M., 2011. Antimicrobial resistance patterns of *Campylobacter* species isolated from raw chicken, turkey, quail, partridge, and ostrich meat in Iran. *Food Control*. 22, 1165- 1170.
 39. Refaie, R.S. and B. Galal, 1992. Survival and growth of *Campylobacter jejuni* in meat. *Assuit Vet. Med. J.*, 28, 174-179.
 40. Ryan, K.J. and Ray, C.G., 2004. *Sherris Medical Microbiology* (4th ed.). McGraw Hill. ISBN 978-0-8385-8529-0.
 41. Skirrow, M.B. and Blaser, M.J. (2000) *Clinical Aspects of Campylobacter Infection*. In: Nachamkin, I. and Blaser, M.J., 2nd Eds., *Campylobacter*. American Society for Microbiology, Washington DC, 11,69-88.
 42. Smibert, R.M. (1974): *Campylobacter*: In *Bergey's Manual of Determinative Bacteriology* 8th Ed., Baltimore, Williams and Wilkins 207-211.
 43. Stanley K. and Jones K. 2003. Cattle and sheep farms as reservoirs of *Campylobacter*. *J Appl Microbiol* 94(Suppl.), 104S–113S. 10.1046/j.1365-2672.94.s1.12.x
 44. Stern, N. J., Green, S. S., Thaker, N., Krout, D. J., and Chiu, J. (1984). Recovery of *Campylobacter jejuni* from Fresh and Frozen Meat and Poultry Collected at Slaughter. *Journal of food protection*, 47(5), 372–374.
 45. Svedham, A., B. Kaijser, and E. Sjogren. 1981. The occurrence of *Campylobacter jejuni* in fresh food and survival under different conditions. *J. Hyg.* 87:421-425.
 46. Takeuchi, M. G., de Melo, R. T., Dumont, C. F., Peixoto, J. L. M., Ferreira, G. R. A., Chueiri, M. C., Iasbeck, J. R., Timóteo, M. F., de Araújo Brum, B., and Rossi, D. A., 2022.
 47. Agents of *Campylobacteriosis* in Different Meat Matrices in Brazil. *International journal of environmental research and public health*, 19(10), 6087.
 48. Thomas, K. M., de Glanville, W. A., Barker, G. C., Benschop, J., Buza, J. J., Cleaveland, S., Davis, M. A., French, N. P., Mmbaga, B. T., Prinsen, G., Swai, E. S., Zadoks, R. N., and Crump, J. A., 2020. Prevalence of *Campylobacter* and *Salmonella* in African food animals and meat: A systematic review and meta-analysis. *International journal of food microbiology*, 315, 108382.
 49. Trokhymchuk, A., Waldner, C., Chaban, B., Gow, S., and Hill, J. E., 2014. Prevalence and diversity of *Campylobacter* species in Saskatchewan retail ground beef. *J. Food Prot.* 77, 2106–2110.
 50. Turnbull, P. C. B., and P. Rose. (1982). *Campylobacter jejuni* and *Salmonella* in raw meats. *J. Hyg.* 88:29-37.
 51. Vipham, J. L., Brashears, M. M., Loneragan, G. H., Echeverry, A., Brooks, J. C., Chaney, W. E., and Miller, M. F., 2012. *Salmonella* and *Campylobacter* baseline in retail ground beef and whole-muscle cuts purchased during 2010 in the United States. *Journal of food protection*, 75(12), 2110–2115.
 52. Wang, G., Clark, C. G., Taylor, T. M., Pucknell, C., Barton, C., Price, L., Woodward, D. L., and Rodgers, F. G., 2002.

- Colony multiplex PCR assay for identification and differentiation of *Campylobacter jejuni*, *C. coli*, *C. lari*, *C. upsaliensis*, and *C. fetus* subsp. *fetus*. *Journal of clinical microbiology*, 40(12), 4744–4747.
53. World Health Organization (WHO)(2011): *Campylobacter* FactsheetN°255October
54. Zhao, C., Ge, B., De Villena, J., Sudler, R., Yeh, E., Zhao, S., White, D. G., Wagner, D., and Meng, J., 2001. Prevalence of *Campylobacter* spp., *Escherichia coli*, and *Salmonella* serovars in retail chicken, turkey, pork, and beef from the Greater Washington, D.C., area. *Applied and environmental microbiology*, 67(12), 5431–5436.