



# Microbes and Infectious Diseases

Journal homepage: <https://mid.journals.ekb.eg/>

## Original article

## Detection of *Mycobacterium tuberculosis* in tongue Swab

Daniel Abigail Abi<sup>1</sup>, Bimba John Samson<sup>2</sup>, Solomon Matthias Gamde<sup>3\*</sup>, Simon Peter Abriba<sup>3</sup>

1- Department of Biology, School of Basic Studies, Bingham University Karu, Nasarawa State.

2- Department of Community Medicine/Zankli Research Centr, Bingham University Karu, Nasarawa State.

3- Department of medical laboratory science, Bingham University Karu, Nasarawa State

### ARTICLE INFO

#### Article history:

Received 19 July 2023

Received in revised form 12 August 2023

Accepted 14 August 2023

#### Keywords:

Sputum, GeneXpert MTB/RIF

Tongue swab

pulmonary tuberculosis

Prevalence

### ABSTRACT

**Background:** Sputum examination is the gold standard for the diagnosis of pulmonary tuberculosis. However, individuals particularly children, the aged, and immuno-compromised may not expectorate sputum due to some underlying conditions. Moreover, sputum produces infectious aerosols which exposes both the patients and care giver. It is also difficult to standardize sputum specimen for *Mycobacterium tuberculosis*. Therefore, the present study considered alternative specimen for pulmonary tuberculosis. **Aim:** To identify *Mycobacterium tuberculosis* in tongue swab. **Methods:** This was a cross-sectional laboratory-based study carried out in July-December 2022 on 384 tuberculosis patients attending health facilities in Karu, North central Nigeria. Using a multistage sampling technique, participants were added one at a time until the required sample size was gotten and analyzed via GeneXpert MTB/RIF. Performance of the tongue swab was compared to the combined reference standard of sputum XPERT. **Results:** The tongue swab XPERT ultra had a sensitivity of 78%, a specificity of 100%, a positive predictive value of 100%, and a negative predictive value of 97.4% relative to a sputum XPERT ultra reference standard. There were positive associations between the socio-demographic variables (age group, sex, family size, family type, marital status, religion, educational status, and occupation) and *Mycobacterium tuberculosis* infection. However, only the participant's educational status ( $p=0.027$ ) and occupation ( $p=0.05$ ) had statistically significant difference. **Conclusion:** Tongue swab is an alternate specimen for sputum-based molecular diagnosis for *Mycobacterium tuberculosis*. The procedure is less invasive, easier, and non-aerosol producing efficient alternate approach for pulmonary tuberculosis.

### Introduction

According to The World Health Organization (WHO), tuberculosis is among the major causes of morbidity and mortality in the world. Every year, more than 10 million people in their prime years are infected with active tuberculosis especially in poor regions of Africa and Asia [1]. Sputum analysis is the gold standard for the diagnosis of pulmonary tuberculosis. It is a viscous substance produced by the airways of

patients. However, individuals particularly children, the aged, and immuno-compromised patients may not expectorate sputum due to the underlying medical conditions [2, 3]. Moreover, sputum produces infectious aerosols which exposes both the patients and care giver. It is also difficult to standardize and prepare the sample for *Mycobacterium tuberculosis* DNA detection [3,4]. Therefore, modern research has looked at alternative samples such as exhaled breath concentration,

DOI: 10.21608/MID.2023.223973.1566

\* Corresponding author: Solomon Gamde, Matthias

E-mail address: [solomonmatthias85@gmail.com](mailto:solomonmatthias85@gmail.com)

© 2020 The author (s). Published by Zagazig University. This is an open access article under the CC BY 4.0 license <https://creativecommons.org/licenses/by/4.0/>.

blood, urine, and saliva [6]. Unfortunately, these samples were typically less sensitive and specific than the sputum [7].

Although, sputum samples from person's exhibiting symptoms of tuberculosis are used in the majority of contemporary diagnosis of tuberculosis, the non-sputum-based diagnosis have been identified by the World Health Organization as a high priority for diagnostic development to stop the spread of tuberculosis [2]. It is indispensable to identify and treat individuals with active tuberculosis as soon as possible. To stop the spread of tuberculosis, active case-finding demand a high-turnout sampling of a large number of persons but the use of sputum would worsen the existing problems. A simple, safe, non-invasive, and more effective alternative technique without non-aerosol such as the use of tongue swab is required to identify the instigating organism in order to support early detection and treatment of the disease.

Documented evidence abound that *Mycobacterium tuberculosis* can be recovered from the mouth of individuals living with pulmonary tuberculosis. Oral swabs have been shown in studies to be a suitable substitute to sputum testing. Oral swabs have the same sensitivity as a sputum sample [8,9]. Moreover, the existence of *Mycobacterium tuberculosis* in the bucal cavity is conceivable given that the bacteria is found in the tonsil and adenoid of patients having tonsillectomies, suggests a way to measure tuberculosis prevalence. In addition, tuberculosis testing efficiency would rise with the availability of the non-invasive oral swabs samples that are simple to collect outside the clinic while reducing the risk of exposure. In this context, addressing the underscored issues will escalate the journey towards the goal of tuberculosis elimination. Therefore, this study aimed to identify the diagnostic accuracy of *Mycobacterium tuberculosis* in tongue Swab.

## MATERIALS AND METHODS

### Study Design and Population

This is a prospective and cross-sectional laboratory-based study carried out in July/December 2022 on 384 tuberculosis patients of the age 18 years and above attending health facilities in Karu, Nasarawa State, Nigeria. Using a multistage sampling technique; stage 1 was the selection of Health Facilities, stage 2 was the selection of DOTS clinics, and stage 3 was the selection of study participants. Four (4) health facilities in Karu were

selected by simple random sampling technique. The health facilities are the major hospitals located within the state providing essential, specialized, and referral medical services to the residents and neighboring states. Of the four health facilities selected, two health facilities with DOTS clinics (Mararaba Medical Center and General Hospital Uke) were selected by simple random sampling. Participants were added one at a time until the required sample size was gotten to detect the diagnostic accuracy of *Mycobacterium tuberculosis* using tongue swab. The study was conducted on a pool of clinical specimens- tongue swabs according to the international guidelines of Strengthening the Reporting for Observational Studies in Epidemiology (STROBE).

### Ethical Approval, Recruitment, and Enrollment

This study was carried out in accordance with the Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments involving human subjects. The study procedures was approved by the Ethics Committee, Bingham University Teaching Hospital Jos with Ref. No. NHREC/21/05/2005/00965. Informed consent was obtained from each study subject and confidentiality of patient information and samples was maintained at all times. Spent samples were appropriately disposed by autoclaving.

A structured questionnaire was administered to each participant to obtain socio-demographic information which was analyzed to determine the associated risk factors for *Mycobacterium tuberculosis* infection in the study. The study participants were given standardized questionnaires upon enrolment that inquired about their socio-demographics, clinical features, and epidemiological/laboratory data. Each participant had a sample of their sputum and tongue swab taken early in the morning. Participants were asked to refrain from eating, drinking, brushing their teeth, and using mouthwash for at least 30 minutes prior to sample collection. For the tongue swab, participants were told to stick out their tongues, and the study researcher ran a sterile, individually wrapped swab over the front 2/3 of each participant's tongue, measuring its length and width. With just the right amount of pressure, the swab shafts might be slightly bent. Samples were obtained while continually spinning the swab for 15 to 20 seconds. The sample collection bottle cap was taken off as soon as the sample was collected and the shaft

section of the swab was carefully snapped off into the bottle containing 2ml normal saline so that the cap could seal completely. Sample collection bottle was sealed tightly. For the sputum sample, the participants were instructed to perform a deep cough and generate high-quality sputum in a sputum cup as part of the recruitment process. The sample containers for each participant were suitably labeled with their initials and date of collection. At the clinic, samples were kept inside a leak-proof bio-hazard bag and then into an ice-packed box before transporting to Zankli Research Center at Bingham University for Xpert MTB/RIF test.

### Expected Outcome

Performance of the tongue swab was compared to the combined reference standard of sputum XPERT, which is the WHO's suggested first line TB diagnosis. Participants who tested positive for Mycobacterium tuberculosis in their sputum were deemed to have Mycobacterium tuberculosis (TB), whereas those who tested negative were deemed to be TB negative. Swab XPERT positivity for Mycobacterium tuberculosis was used to determine the tongue swab positivity.

### Statistical Analysis

Data obtained from the study were entered into Microsoft Excel 2010 and analyzed with SPSS version 25.0 (SPSS Inc., Chicago, IL, USA). Appropriate descriptive statistic was used to summarize the socio-demographic data of the participants. Sensitivity and specificity of the tongue swab Xpert were calculated in reference to sputum Xpert for tuberculosis. Correlations between variables were determined using the Pearson Chi-square test and the statistical significance value was determined at  $p \leq 0.05$ .

## RESULTS

Section A: Socio-demographic characteristics

### Table 1: Socio-demographic Characteristics of the Study Population

Of the 384 samples analyzed for Mycobacterium tuberculosis, 57.8% (222/384) were males and 42.2% (162/384) were females, giving a gender ratio of 1:0.730 (males: females). Majority of the study participants 109 (28.4%) were of age 18–27 years while the least was the oldest age group  $\geq 57$  years, which included 40 (10.4%) of the individuals. It was obvious that most of the study population were literate. According to the findings of the study, 259 (67.4%) of the participants are

married, 2 (0.5%) are divorced and 7 (1.8%) are widowed. Furthermore, 116(30.2%) are single and 235(61.2%) have a family size of 5 and above.

### Table 2: Assessment of the Sputum for Mycobacterium tuberculosis

The positivity rate of Mycobacterium tuberculosis infection among the study population was age dependent, age group 18 – 27 years had the highest positivity rate of 13.5% (13/96) and the least was among the older age group of 57 years, 0.14% (5/35). There were positive associations between the socio-demographic variables (age group, sex, family size, family type, marital status, religion, educational status, and occupation) and Mycobacterium tuberculosis infection. However, only the educational status of the study population had a statistically significant relationship ( $p=0.030$ ).

### Table 3: Assessment of the throat Swab for Mycobacterium tuberculosis

The positivity rate of Mycobacterium tuberculosis infection was found among the age group 18 – 27 years, 17.2% (16/93) and the least was among the older age group 48 – 57 years, 14.3% (8/56). There were positive associations between the socio-demographic variables (age group, sex, family size, family type, marital status, religion, educational status, and occupation) and Mycobacterium tuberculosis infection. However, only the participant's educational status ( $p=0.027$ ) and occupation ( $p=0.05$ ) have a statistically significant difference.

## SECTION B: Clinical Characteristics

### Table 4: Clinical Characteristics of the Study Population

Majority 339(88.3%) of the respondents have not cough for more than two weeks. Again, 360(93%) said they don't cough with mucus. A total of 370(96.4%) respondents haven't experience weight loss over the last three months while 372(96.9%) respondents sweat at night. All the respondents have not been treated for TB nor 381(99.2%) of their immediate family has TB. About 347(90.4%) were not feeling sick.

Section C: Epidemiological and Laboratory Characteristics

### Table 5: Epidemiological Characteristics of the Study Participants

All participants 384(100%) identified pulmonary tuberculosis as their site of presumptive TB and DS –TB. The overall sensitivity of

Mycobacterium tuberculosis in sputum was 10.7% (41/384) while tongue swab was 8.3% (32/384).

**HIV Status of Participants**

The HIV status of the study participants 7(1.8%) were positive and 256(66.7%) were negative while 119(31%) do not know their HIV status.

Overall, tongue swab XPERT detected 32(8.3%) positive and 352(91.7%) negative while the sputum XPERT detected 41(10.7%) positive and 343(89.3%) negative for pulmonary tuberculosis.

**Table 6: Sensitivity Assessment of the Sputum and Tongue Swab**

Table 6 shows that the validity of the swab result is 78.0% sensitive and 100% specific. The swab result is also reliable with a positive predictive value of 100% and negative predictive value of 97.4% pulmonary tuberculosis detection using tongue swab in the study population at 95% CI.

Section A: Socio-demographic Characteristics of the Participants'

**Table 1.** Socio-demographic Characteristics of the Study Population (n = 384).

Variable		Frequency	Percentage (%)
<b>Age group</b>	18 – 27 years	109	28.4
	28 – 37 years	85	22.1
	38 – 47 years	86	22.4
	48 – 57 years	64	16.7
	>57 years	40	10.4
<b>Sex</b>	Male	222	57.8
	Female	162	42.2
<b>Family type</b>	Single	186	48.4
	Joint	198	51.6
<b>Family Size</b>	≤5	149	38.8
	>5 members	235	61.2
<b>Marital Status</b>	Single	116	30.2
	Married	259	67.4
	Separated	2	0.5
	Widow	7	1.8
<b>Religion</b>	Christianity	334	87.0
	Islam	50	13.0
<b>Occupation</b>	Farming	45	11.7
	Business	170	44.3
	Student	91	23.7
	Unemployed	52	13.5
	Others	26	6.8
<b>Educational Status</b>	None	19	4.9
	Primary	80	20.8
	Secondary	200	52.1
	Tertiary	85	22.1

**Table 2.** Assessment of the Sputum for *Mycobacterium tuberculosis*.

Variables		Sputum/XPERT Result (%)		X <sup>2</sup>	P-value
		Positive (n)	Negative (n)		
Age group	18 – 27 years	13 (3.39%)	96 (25%)	6.758	0.149
	28 – 37 years	2 (0.52%)	83 (21.61%)		
	38 – 47 years	7 (1.82%)	79 (20.57%)		
	48 – 57 years	5 (1.30%)	59 (15.36%)		
	>57 years	5 (1.30%)	35 (9.11%)		
Sex	Male	20 (5.21%)	202 (52.60%)	0.314	0.575
	Female	12 (3.13%)	150 (39.06%)		
Family type	Single	14 (3.65%)	172 (44.79%)	0.307	0.579
	Joint	18 (4.69%)	180 (46.88%)		
Family Size	<= 5	14 (3.65%)	135 (35.16%)	0.360	0.549
	>5 members	18 (4.69%)	217 (56.51%)		
Marital Status	Single	11 (2.86%)	105 (27.34%)	1.036	0.793
	Married	21 (5.47%)	238 (61.98%)		
	Separated	0 (0.00%)	2 (0.52%)		
	Widow	0 (0.00%)	7 (1.82%)		
Religion	Christianity	30 (7.81%)	304 (79.17%)	1.413	0.235
	Islam	2 (0.52%)	48 (12.50%)		
Occupation	Farming	3 (0.78%)	42 (10.94%)	9.071	0.059
	Business	11 (2.86%)	159 (41.4%)		
	Student	9 (2.34%)	82 (21.35%)		
	Unemployed	3 (0.78%)	49 (12.76%)		
	Others	6 (1.56%)	20 (5.20%)		
Educational Status	None	5 (1.30%)	14 (3.65%)	8.942	0.030
	Primary	7 (1.82%)	73 (19.01%)		
	Secondary	13 (3.39%)	187 (48.70%)		
	Tertiary	7 (1.82%)	78 (20.31%)		

\* Statistically significant values are determined at  $p \leq 0.05$ .**Table 3.** Assessment of the tongue swab for *Mycobacterium tuberculosis*.

Variables		Swab Result (%)		X <sup>2</sup>	P-value
		Positive	Negative		
Age group	18 – 27 years	16 (4.12%)	93 (24.22%)	7.561	0.109
	28 – 37 years	3 (0.78%)	82 (21.35%)		
	38 – 47 years	8 (2.08%)	78 (20.31%)		
	48 – 57 years	8 (2.08%)	56 (14.58%)		
	>57 years	6 (1.56%)	34 (8.85%)		
Sex	Male	24 (6.25%)	198 (51.56%)	0.010	0.921
	Female	17 (4.43%)	145 (37.76%)		
Family type	Single	19 (4.95%)	167 (43.49%)	0.081	0.776
	Joint	22 (5.73%)	176 (45.83%)		
Family Size	<= 5	18 (4.69%)	131 (34.11%)	0.503	0.473
	>5 members	23 (5.99%)	212 (55.20%)		
Marital Status	Single	15 (3.91%)	101 (26.30%)	1.804	0.614
	Married	26 (6.77%)	233 (60.68%)		
	Separated	0 (0.00%)	2 (0.52%)		
	Widow	0 (0.00%)	7 (1.82%)		
Religion	Christianity	39 (10.16%)	295 (76.8%)	2.687	0.101
	Islam	2 (0.52%)	48 (12.50%)		

<b>Occupation</b>	Farming	4 (1.04%)	41 (10.68%)	9,496	0.050
	Business	14 (3.65%)	156 (40.63%)		
	Student	12 (3.13%)	79 (20.57%)		
	Unemployed	4 (1.04%)	48 (12.50%)		
	Others	7 (1.82%)	19 (4.95%)		
<b>Educational Status</b>	None	6 (1.56%)	13 (3.89%)	9,175	0.027
	Primary	8 (2.08%)	72 (18.75%)		
	Secondary	19 (4.95%)	181 (47.14%)		
	Tertiary	8 (2.08%)	77 (20.05%)		

\* Statistically significant values are determined at  $p \leq 0.05$ .

SECTION B: Clinical Characteristics

**Table 4.** Clinical characteristics of the study population.

<b>Variables</b>		<b>Frequency</b>	<b>Percentage (%)</b>
<b>Cough for two (2) weeks</b>	Yes	45	11.7
	No	339	88.3
<b>Cough with mucus</b>	Yes	24	6.3
	No	360	93.8
<b>Weight loss within three (3) months</b>	Yes	14	3.6
	No	370	96.4
<b>Night sweating</b>	Yes	12	3.1
	No	372	96.9
<b>Previous TB treatment</b>	Yes	0	0.0
	No	384	100.0
<b>Immediate family members with TB</b>	Yes	3	0.8
	No	381	99.2
<b>Currently feeling sick</b>	Yes	37	9.6
	No	347	90.4
<b>History of smoking</b>	Yes	0	0.0
	No	384	100.0

Section C: Epidemiological and Laboratory Characteristics of the Study Participants

**Table 5.** Epidemiological characteristics of the study participants.

<b>Variables</b>		<b>Frequency</b>	<b>Percentage (%)</b>
<b>Site of Presumptive TB</b>	Pulmonary TB	384	100.0
<b>Type of Presumptive Tuberculosis</b>	DS-TB	384	100.0
<b>HIV Status</b>	Positive	7	1.8
	Negative	256	66.7
	Unknown	119	31.0
<b>Reason for Examination</b>	Diagnosis	384	100.0
<b>Test Requested</b>	Xpert/MTB/RIF	384	100.0
<b>Swab/XPERT Result</b>	Positive	32	8.3
	Negative	352	91.7
<b>Sputum/ XPERT Result</b>	Positive	41	10.7
	Negative	343	89.3

**Table 6.** Cross tabulation on sputum Xpert and swab Xpert.

	Sputum Xpert			
		Positive	Negative	Total
Swab Xpert	Positive	32= TP (%)	0= FP (%)	32= (TP+FP)
	Negative	9= FN (%)	343= TN (%)	352= (FN+TN)
	Total	41= (TP+FN)	343= (FP+TN)	384= N

Where TP is True Positive; FN is False Negative; FP is False Positive, TN is True Negative and N is total number of samples

**Sensitivity**=  $[TP/(TP+FN)] \times 100\% = 78.0\%$

**Specificity**=  $[TN/(TN+FP)] \times 100\% = 100\%$


**PPV**=  $[TP/(TP+FP)] \times 100\% = 100\%$

**NPV**=  $[TN/(TN+FN)] \times 100\% = 97.4\%$

**Accuracy**=  $(TP+TN)/N \times 100\% = 97.7\%$

**Figure 1.** Receiver's operating characteristic curve of sputum Xpert and swab Xpert

From the ROC graph, the area under the curve is  $c = 0.974$  which indicates good predictive power of the tongue swab.



**Youden index** = Sensitivity (%) + Specificity (%) - 100 = 78%. The cut-off point for having an acceptable Youden index is 50% hence, 78% is above 50% which denotes that the tongue swab meets empirical benchmarks for being administered for diagnostic purposes of tuberculosis.

## Discussion

Some patients, children, the aged, and immuno-compromised may not expectorate quality sputum for testing *Mycobacterium tuberculosis* due to underlying medical conditions [10, 11, 12]. Oral swabs provide several potential diagnostic benefits for the most part in community settings where sputum collection for TB is unfeasible. This study aimed to identify the diagnostic accuracy of *Mycobacterium tuberculosis* in tongue Swab. Our data showed that participants between the ages of 18 - 27 years had the highest positivity rate for *Mycobacterium tuberculosis*. This finding is tandem with previous reports [13, 15, 16].

In this cross-sectional laboratory-base study, we found that tongue swab has similar sensitivity for *Mycobacterium tuberculosis* in comparison to the sputum sample. Similar studies were reported by Luabeya et al. [11] in South Africa and [12]. Tongue swab could be used to diagnose *Mycobacterium tuberculosis*. A sensitivity rate of 45% was obtained in a study on cheek swab in Peru [4]. In another study of mass TB screening among prisoners in Brazil indicated a sensitivity rate of 51% on tongue swabs. Besides, another study showed a single swab sensitivity of 88% (44/50) on day 1 and 94.4% (17/18) on day 2 with a specificity of 79.2% (42/53) [14, 17, 18]. Erroneous TB

diagnoses especially in high-burden countries with limited access to healthcare resources could widen the case-detection gap and promotes transmission rate of tuberculosis [19]. In the present study, the tongue swab GeneXpert performance in the detection of Mycobacterium tuberculosis was higher than the sputum GeneXpert in the early morning sample collection before engagement in any oral activity. Moreover, tongue swab has the advantage of lowering biosafety concerns regardless of the bacterial viability. Consequently, this study adds to the growing body of knowledge regarding the use of tongue swabs in the diagnosis of tuberculosis. Even though the morning oral swab had the maximum sensitivity, it only detected 78.0% of individuals with laboratory-confirmed pulmonary TB.

According to our results, one oral swab's sensitivity was 78% valid. The significant absence of overlap in MTB detection across the samples collected from the participants was another intriguing finding. Pulmonary TB is the site of presumptive TB for all participants with DS-TB as the type of presumptive TB. A good number of the participants 119(31.0%) said they don't know their HIV status while the majority of the participants (66.7%) knew their status as HIV negative and 7(1.8%) as HIV positive. All the participants' reason for examination is the diagnosis for XPERT/MTB/RIF in which 91.7% tested negative on sputum XPERT/MTB/RIF and 89.3% tested negative on swab XPERT/MTB/RIF respectively.

From the correlation table above the value of 87.2% shows a positive relationship between the Swab result and the sputum gene Xpert result. The relationship, however, has a significant effect ( $0.001 < 0.01$ ) which means that there are sufficient reasons to uphold the alternate hypothesis. This shows that swab kits are valid and reliable in line with previous reports [20, 21, 22, 23].

## CONCLUSION

Our finding demonstrated that Mycobacterium tuberculosis could be found in tongue swab especially for individuals who had difficulty producing sputum. The potential for non-sputum base diagnostics could be enhanced by the use of tongue swab. The procedure is less invasive, easier, and non-aerosol producing efficient alternate approach for pulmonary tuberculosis and should be further investigated.

## Acknowledgement

The assistance of the laboratory Staff from Zankli Research Center, Bingham University Karu is cherished and worthy of our commendation.

## Conflict of Interest

Nothing to declare.

## Funding

None.

## References

- 1- **World Health Organization (WHO)**. Global tuberculosis report 2020. Available at: [https://www.who.int/tb/publications/global\\_report/en/](https://www.who.int/tb/publications/global_report/en/). Accessed October 9, 2021.
- 2- **Vijay N**. Tuberculosis diagnostics: challenges and opportunities. Lung India: Official Organ of Indian Chest Society. Division of Microbiology and Clinical Pathology, National AIDS Research Institute Pune, Maharashtra, India. Lung India 2012; 29(3): 259-266. Doi: 10.4103/0970-2113.99112.
- 3- **Shapiro AE, Olson AM, Kidoguchi L, Niu X, Ngcobo Z, Magcaba ZP, et al**. Complementary Nonsputum Diagnostic Testing for Tuberculosis in People with HIV Using Oral Swab PCR and Urine Lipoarabinomannan Detection. J Clin Microbiol. 2022 Aug 17;60(8):e0043122. doi: 10.1128/jcm.00431-22. Epub 2022 Aug 1. PMID: 35913145; PMCID: PMC9383113.
- 4- **Andama A, Whitman GR, Crowder R, Reza TF, Jaganath D, Mulondo J, et al**. Accuracy of Tongue Swab Testing Using Xpert MTB-RIF Ultra for Tuberculosis Diagnosis. J Clin Microbiol. 2022 Jul 20;60(7):e0042122. doi: 10.1128/jcm.00421-22. Epub 2022 Jun 27. PMID: 35758702; PMCID: PMC9297831.
- 5- Erickson BR, Sealy TK, Flietstra T, Morgan L, Kargbo B, Matt-Lebby VE, et al. Ebola virus disease diagnostics, Sierra Leone: analysis of real-time reverse transcription polymerase



- chain reaction values for clinical blood and oral swab specimens. *J Infect Dis* 2016; 214 (suppl. 3):258–62.  
<https://doi.org/10.1093/infdis/jiw296>.
- 6- **Balán I, Frasca T, Ibitoye M, Dolezal C, Carballo-Diéguez A.** Fingerprick versus oral swab: acceptability of blood-based testing increases if other STIs can be detected. *AIDS Behav* 2017; 21(2):501–4.  
<https://doi.org/10.1007/s10461-016-1497-4>.
- 7- **Daum LT, Peters RP, Fourie PB, Jonkman K, Worthy SA, Rodriguez JD, et al.** Molecular detection of *Mycobacterium tuberculosis* from sputum transported in Prime Store (R) from rural settings. *Int. J. Tuberc. Lung Dis* 2015; 19, 552–557.
- 8- **Cox H, Workman L, Bateman L, Franckling-Smith Z, Prins M, Luiz J, et al.** Oral Swab Specimens Tested with Xpert MTB/RIF Ultra Assay for Diagnosis of Pulmonary Tuberculosis in Children: A Diagnostic Accuracy Study. *Clin Infect Dis*. 2022 Dec 19;75(12):2145-2152. doi: 10.1093/cid/ciac332. PMID: 35579497; PMCID: PMC9761885.
- 9- **Ealand C, Peters J, Jacobs O, Sewcharran A, Ghoor A, Golub J, et al.** Detection of *Mycobacterium tuberculosis* Complex Bacilli and Nucleic Acids from Tongue Swabs in Young, Hospitalized Children. *Front Cell Infect Microbiol*. 2021 Jun 14;11:696379. doi: 10.3389/fcimb.2021.696379. PMID: 34195103; PMCID: PMC8238041.
- 10- **Fauci AS, Eisinger RW.** Reimagining the Research Approach to Tuberculosis. *Am J Trop Med Hyg* 2018; 98(3):650-2.  
<https://doi.org/10.4269/ajtmh.17-0999>  
 PMID:29363452.
- 11- **Luabeya AK, Wood RC, Shenje J, Filander E, Ontong C, Mabwe S, et al.** Noninvasive Detection of Tuberculosis by Oral Swab Analysis. *J. Clin. Microbiol* 2019; 57, e0184718.
- 12- **Byanyima P, Kaswabuli S, Musisi E, Nabakiibi C, Zawedde J, Sanyu I, et al.** Feasibility and Sensitivity of Saliva GeneXpert MTB/RIF Ultra for Tuberculosis Diagnosis in Adults in Uganda. *Microbiol Spectr*. 2022 Oct 26;10(5):e0086022. doi: 10.1128/spectrum.00860-22. Epub 2022 Sep 26. PMID: 36154664; PMCID: PMC9603304.
- 13- **Luabeya AK, Wood RC, Shenje J, Filander E, Ontong C, Mabwe S, et al.** Noninvasive Detection of Tuberculosis by Oral Swab Analysis. *J Clin Microbiol*. 2019 Feb 27;57(3):e01847-18. doi: 10.1128/JCM.01847-18. PMID: 30541931; PMCID: PMC6425180.
- 14- **Mesman AW, Roger C, Martin S, Julia C, Juan A, Milagros M, et al.** *Mycobacterium tuberculosis* detection from oral swabs with Xpert MTB/RIF ULTRA: a pilot study 2019; 12:349 <https://doi.org/10.1186/s13104-019-4385-y>
- 15- **Murray H, Tuazon C, Kirmani N, Sheagren J.** The adult respiratory distress syndrome associated with military tuberculosis. *Chest* 1998; 73:37–43.
- 16- **World Health Organization (WHO).** Global Tuberculosis Report. 2021. Available at: <https://www.who.int/publications/i/item/9789240037021>.
- 17- **Lima F, Santos AS, Oliveira RD, Silva CCR, Gonçalves CCM, Andrews JR, et al.** Oral swab testing by Xpert® MTB/RIF Ultra for mass tuberculosis screening in prisons. *J Clin Tuberc Other Mycobact Dis*. 2020 Feb 4;19:100148. doi: 10.1016/j.jctube.2020.100148. Erratum in: *J Clin Tuberc Other Mycobact Dis*. 2020 Sep

- 09;21:100177. PMID: 32099908; PMCID: PMC7031315.
- 18-**Nicol MP, Wood RC, Workman L, Prins M, Whitman C, Ghebrekristos Y, et al.** Microbiological diagnosis of pulmonary tuberculosis in children by oral swab polymerase chain reaction. *Sci. Rep* 2019; 9: 10789.
- 19-**Wang Y, He Y, Wang L, Zhang YA, Wang MS.** Diagnostic Yield of Nucleic Acid Amplification Tests in Oral Samples for Pulmonary Tuberculosis: A Systematic Review and Meta-analysis. *Open Forum Infect Dis.* 2023 Feb 14;10(3):ofad082. doi: 10.1093/ofid/ofad082. PMID: 36937246; PMCID: PMC10019807.
- 20-**Flores JA, Calderón R, Mesman AW, Soto M, Coit J, Aliaga J, et al.** Detection of Mycobacterium Tuberculosis DNA in Buccal Swab Samples from Children in Lima, Peru. *Pediatr Infect Dis J.* 2020 Nov;39(11):e376-e380. doi: 10.1097/INF.0000000000002828. PMID: 32675773; PMCID: PMC7578116.
- 21- **Mesman AW, Calderon RI, Pollock NR, Soto M, Mendoza M, Coit J, et al.** Molecular detection of Mycobacterium tuberculosis from buccal swabs among adult in Peru. *Sci Rep.* 2020 Dec 17;10(1):22231. doi: 10.1038/s41598-020-79297-9. PMID: 33335256; PMCID: PMC7746708.
- 22-**Molina-Moya B, Ciobanu N, Hernandez M, Prat-Aymerich C, Crudu V, Adams ER, et al.** Molecular Detection of *Mycobacterium tuberculosis* in Oral Mucosa from Patients with Presumptive Tuberculosis. *J Clin Med.* 2020 Dec 21;9(12):4124. doi: 10.3390/jcm9124124. PMID: 33371314; PMCID: PMC7767357.
- 23- **LaCourse SM, Seko E, Wood R, Bundi W, Ouma GS, Agaya J, et al.** Diagnostic performance of oral swabs for non-sputum based TB diagnosis in a TB/HIV endemic setting. *PLoS One.* 2022 Jan 13;17(1):e0262123. doi: 10.1371/journal.pone.0262123. PMID: 35025930; PMCID: PMC8758000.