

A STUDY OF INTUITIONISTIC FUZZY I -CONVERGENT DOUBLE SEQUENCE SPACES DEFINED BY COMPACT OPERATOR

VAKEEL A. KHAN, YASMEEN, AYHAN ESI, HIRA FATIMA AND MOBEEN AHAMAD

ABSTRACT. After the pioneering work of Zadeh[35], a huge number of research papers have been appeared on fuzzy theory and its applications as well as fuzzy analogues of the classical theories. Fuzzy set theory is a powerful hand set for modelling uncertainty and vagueness in various problems arising in field of science and engineering. It has a wide range of applications in various fields: population dynamics[1], chaos control[4], computer programming[5], nonlinear dynamical system[6], etc. Fuzzy topology is one of the most important and useful tools and it proves to be very useful for dealing with such situations where the use of classical theories breaks down. In this paper, we introduce ideal convergence of double sequences in intuitionistic fuzzy sequences defined by compact operator and study the fuzzy topology on the said spaces.

1. INTRODUCTION AND PRELIMINARIES:

The concept of intuitionistic fuzzy normed space[32] and of intuitionistic fuzzy 2-normed space[27] are the latest developments in fuzzy topology. Recently V. A. Khan and Yasmeen[[7],[9], [10],[11],[12]] studied the intuitionistic fuzzy Zweier I -convergent sequence spaces defined by paranorm, modulus function and Orlicz function. And recently Totur, Çanak and Önder [[2],[31], [34]] studied and defined some summable double sequence of fuzzy numbers.

The notion of statistical convergence is a very useful functional tool for studying the convergence problems of numerical problems/matrices(double sequences) through the concept of density. The notion of I -convergence, which is a generalization of statistical convergence[[3],[19],[20],[21],[22],[23],[24],[25],[26],[28]] was introduced by Kostyrko, Salat and Wilczynski[15] by using the idea of I of subsets of the set of natural numbers \mathbb{N} and further studied in[[30],[8],[13]]. Recently, the notion of statistical convergence of double sequences $x = (x_{ij})$ has been defined and studied by Mursaleen and Edely[18], and for fuzzy numbers by Savaş and Mursaleen[33], M. Mursaleen, H. M. Srivastava and S. K. Sharma[29]. Quite recently, Das et al.[16]

2010 *Mathematics Subject Classification.* 41A10, 41A25, 41A36, 40A30.

Key words and phrases. Ideal, filter, I -convergence, I_2 -Convergence in an intuitionistic fuzzy normed spaces, bounded linear operator, compact linear operator.

Submitted May 28, 2018. Revised July 15, 2018 .

studied the notion of I and I^* - convergence of double sequences in \mathbb{R} .

We recall some notations and basic definitions used in this paper.

Definition 1.1 Let $I \subset 2^{\mathbb{N}}$ be a non-trivial ideal in \mathbb{N} . Then a sequence $x = (x_k)$ is said to be I -convergent to a number L if, for every $\epsilon > 0$, the set

$$\{k \in \mathbb{N} : |x_k - L| \geq \epsilon\} \in I.$$

Definition 1.2 Let X be a non empty set. Then $\mathcal{F} \subset 2^X$ is said to be a *filter* on X if and only if $\phi \notin \mathcal{F}$, for $A, B \in \mathcal{F}$ we have $A \cap B \in \mathcal{F}$ and for each $A \in \mathcal{F}$ and $B \supset A$ implies $B \in \mathcal{F}$.

Definition 1.3 Let $I \subset 2^{\mathbb{N}}$ be a non-trivial ideal in \mathbb{N} . Then a sequence $x = (x_k)$ is said to be I -Cauchy if, for each $\epsilon > 0$, there exists a number $N = N(\epsilon)$ such that the set $\{k \in \mathbb{N} : |x_k - x_N| \geq \epsilon\} \in I$.

Definition 1.4(See [8]) The five-tuple $(X, \mu, \nu, *, \diamond)$ is said to be an intuitionistic fuzzy normed space (for short, IFNS) if X is a vector space, $*$ is a continuous t-norm, \diamond is a continuous t-conorm and μ, ν are fuzzy sets on $X \times (0, \infty)$ satisfying the following conditions for every $x, y \in X$ and $s, t > 0$:

- (1) $\mu(x, t) + \nu(x, t) \leq 1$,
- (2) $\mu(x, t) > 0$,
- (3) $\mu(x, t) = 1$ if and only if $x = 0$,
- (4) $\mu(\alpha x, t) = \mu(x, \frac{t}{|\alpha|})$ for each $\alpha \neq 0$,
- (5) $\mu(x, t) * \mu(y, s) \leq \mu(x + y, t + s)$,
- (6) $\mu(x, \cdot) : (0, \infty) \rightarrow [0, 1]$ is continuous,
- (7) $\lim_{t \rightarrow \infty} \mu(x, t) = 1$ and $\lim_{t \rightarrow 0} \mu(x, t) = 0$,
- (8) $\nu(x, t) < 1$,
- (9) $\nu(x, t) = 0$ if and only if $x = 0$,
- (10) $\nu(\alpha x, t) = \nu(x, \frac{t}{|\alpha|})$ for each $\alpha \neq 0$,
- (11) $\nu(x, t) \diamond \nu(y, s) \geq \nu(x + y, t + s)$,
- (12) $\nu(x, \cdot) : (0, \infty) \rightarrow [0, 1]$ is continuous,
- (13) $\lim_{t \rightarrow \infty} \nu(x, t) = 0$ and $\lim_{t \rightarrow 0} \nu(x, t) = 1$.

In this case (μ, ν) is called an intuitionistic fuzzy norm.

Definition 1.5 Let $(X, \mu, \nu, *, \diamond)$ be an IFNS. Then a sequence $x = (x_k)$ is said to be convergent to $L \in X$ with respect to the intuitionistic fuzzy norm (μ, ν) if, for every $\epsilon > 0$ and $t > 0$, there exists $k_0 \in \mathbb{N}$ such that $\mu(x_k - L, t) > 1 - \epsilon$ and $\nu(x_k - L, t) < \epsilon$ for all $k \geq k_0$. In this case we write $(\mu, \nu) - \lim x = L$.

Definition 1.6 Let $(X, \mu, \nu, *, \diamond)$ be an IFNS. Then a sequence $x = (x_k)$ is said to be a Cauchy sequence with respect to the intuitionistic fuzzy norm (μ, ν) if, for every $\epsilon > 0$ and $t > 0$, there exists $k_0 \in \mathbb{N}$ such that $\mu(x_k - x_l, t) < \epsilon$ and $\nu(x_k - x_l, t) < \epsilon$ for all $k, l \geq k_0$.

Definition 1.7 Let K be the subset of natural numbers \mathbb{N} . Then the asymptotic density of K , denoted by $\delta(K)$, is defined as

$$\delta(K) = \lim_n \frac{1}{n} |\{k \leq n : k \in K\}|,$$

where the vertical bars denotes the cardinality of the enclosed set.

A number sequence $x = (x_k)$ is said to be statistically convergent to a number ℓ if, for each $\epsilon > 0$, the set $K(\epsilon) = \{k \leq n : |x_k - \ell| > \epsilon\}$ has asymptotic density zero, i.e.

$$\lim_n \frac{1}{n} |\{k \leq n : |x_k - \ell| > \epsilon\}| = 0.$$

In this case we write $st - \lim x = \ell$.

Definition 1.8 A number sequence $x = (x_k)$ is said to be statistically Cauchy sequence if, for every $\epsilon > 0$, there exists a number $N = N(\epsilon)$ such that

$$\lim_n \frac{1}{n} |\{j \leq n : |x_j - x_N| \geq \epsilon\}| = 0.$$

The concepts of statistical convergence and statistical Cauchy for double sequences in intuitionistic fuzzy normed spaces have been studied by Mursaleen and Mohiuddine[28].

Definition 1.9 Let $I \subset 2^{\mathbb{N}}$ be a non trivial ideal and $(X, \mu, \nu, *, \diamond)$ be an IFNS. A sequence $x = (x_k)$ of elements of X is said to be I -convergent to $L \in X$ with respect to the intuitionistic fuzzy norm (μ, ν) if for every $\epsilon > 0$ and $t > 0$, the set

$$\{k \in \mathbb{N} : \mu(x_k - L, t) \geq 1 - \epsilon \text{ or } \nu(x_k - L, t) \leq \epsilon\} \in I.$$

In this case L is called the I -limit of the sequence (x_k) with respect to the intuitionistic fuzzy norm (μ, ν) and we write $I_{(\mu, \nu)} - \lim x_k = L$.

Definition 1.10(See [14]) Let X and Y be two normed linear spaces and $T : \mathcal{D}(T) \rightarrow Y$ be a linear operator, where $\mathcal{D} \subset X$. Then, the operator T is said to be bounded, if there exists a positive real k such that

$$\|Tx\| \leq k\|x\|, \text{ for all } x \in \mathcal{D}(T).$$

The set of all bounded linear operators $\mathcal{B}(X, Y)$ [17] is a normed linear spaces normed by

$$\|T\| = \sup_{x \in X, \|x\|=1} \|Tx\|$$

and $\mathcal{B}(X, Y)$ is a Banach space if Y is a Banach space.

Definition 1.11(see [14]) Let X and Y be two normed linear spaces. An operator $T : X \rightarrow Y$ is said to be a compact linear operator (or completely continuous linear operator), if

- (1) T is linear,
- (2) T maps every bounded sequence (x_k) in X on to a sequence $(T(x_k))$ in Y which has a convergent subsequence.

The set of all compact linear operators $\mathcal{C}(X, Y)$ is a closed subspace of $\mathcal{B}(X, Y)$ and $\mathcal{C}(X, Y)$ is Banach space, if Y is a Banach space.

2. I_2 -CONVERGENCE IN AN IFNS

Definition 2.1 Let $(X, \mu, \nu, *, \diamond)$ be an IFNS. Then, a double sequence $x = (x_{ij})$ is said to be statistically convergent to $L \in X$ with respect to the intuitionistic fuzzy norm (μ, ν) if, for every $\epsilon > 0$ and $t > 0$,

$$\delta(\{(i, j) \in \mathbb{N} \times \mathbb{N} : \mu(x_{ij} - L, t) \leq 1 - \epsilon \text{ or } \nu(x_{ij} - L, t) \geq \epsilon\}) = 0.$$

or equivalently

$$\lim_{\frac{1}{mn}} \frac{1}{mn} |\{i \leq m, j \leq n, : \mu(x_{ij} - L, t) \leq 1 - \epsilon \text{ or } \nu(x_{ij} - L, t) \geq \epsilon\}| = 0.$$

In this case we write $st_{(\mu, \nu)}^2 - \lim x = L$.

Definition 2.2 Let $(X, \mu, \nu, *, \diamond)$ be an IFNS. Then, a double sequence $x = (x_{ij})$ is said to be statistically Cauchy with respect to the intuitionistic fuzzy norm (μ, ν) if, for every $\epsilon > 0$ and $t > 0$, there exist $N = N(\epsilon)$ and $M = M(\epsilon)$ such that for all $i, p \geq N$ and $j, q \geq M$,

$$\delta(\{(i, j) \in \mathbb{N} \times \mathbb{N} : \mu(x_{ij} - x_{pq}, t) \leq 1 - \epsilon \text{ or } \nu(x_{ij} - x_{pq}, t) \geq \epsilon\}) = 0.$$

Definition 2.3 Let I_2 be a non trivial ideal of $\mathbb{N} \times \mathbb{N}$ and $(X, \mu, \nu, *, \diamond)$ be an intuitionistic fuzzy normed space. A double sequence $x = (x_{ij})$ of elements of X is said to be I_2 convergent to $L \in X$ with respect to the intuitionistic fuzzy norm (μ, ν) if, for each $\epsilon > 0$ and $t > 0$,

$$\{(i, j) \in \mathbb{N} \times \mathbb{N} : \mu(x_{ij} - L, t) \leq 1 - \epsilon \text{ or } \nu(x_{ij} - L, t) \geq \epsilon\} \in I_2.$$

In this case we write $I_2^{(\mu, \nu)} - \lim x = L$.

In this article we introduce the following sequence spaces:

$${}_2S_{(\mu, \nu)}^I(T) = \{(x_{ij}) \in {}_2\ell_\infty : \{(i, j) \in \mathbb{N} \times \mathbb{N} : \mu(T(x_{ij}) - L, t) \leq 1 - \epsilon \text{ or } \nu(T(x_{ij}) - L, t) \geq \epsilon\} \in I_2\};$$

$${}_2S_{0(\mu, \nu)}^I(T) = \{(x_{ij}) \in {}_2\ell_\infty : \{(i, j) \in \mathbb{N} \times \mathbb{N} : \mu(T(x_{ij}), t) \leq 1 - \epsilon \text{ or } \nu(T(x_{ij}), t) \geq \epsilon\} \in I_2\}.$$

We also define an open ball with centre x and radius r with respect to t as follows:

$${}_2B_x(r, t)(T) = \{(y_{ij}) \in {}_2\ell_\infty : \{(i, j) \in \mathbb{N} \times \mathbb{N} : \mu(T(x_{ij}) - T(y_{ij}), t) \leq 1 - \epsilon \text{ or } \nu(T(x_{ij}) - T(y_{ij}), t) \geq \epsilon\} \in I_2\}.$$

3. MAIN RESULTS

Theorem 3.1 ${}_2S_{(\mu, \nu)}^I(T)$ and ${}_2S_{0(\mu, \nu)}^I(T)$ are linear spaces.

Proof. We shall prove the result for ${}_2S_{(\mu, \nu)}^I(T)$. The proof for the other space will follow similarly. Let $x = (x_{ij}), y = (y_{ij}) \in {}_2S_{(\mu, \nu)}^I(T)$ and α, β be scalars. Then for a given $\epsilon > 0$, we have

$$A_1 = \left\{ (i, j) \in \mathbb{N} \times \mathbb{N} : \mu\left(T(x_{ij}) - L_1, \frac{t}{2|\alpha|}\right) \leq 1 - \epsilon \text{ or } \nu\left(T(x_{ij}) - L_1, \frac{t}{2|\alpha|}\right) \geq \epsilon \right\} \in I_2;$$

$$A_2 = \left\{ (i, j) \in \mathbb{N} \times \mathbb{N} : \mu\left(T(y_{ij}) - L_2, \frac{t}{2|\beta|}\right) \leq 1 - \epsilon \text{ or } \nu\left(T(y_{ij}) - L_2, \frac{t}{2|\beta|}\right) \geq \epsilon \right\} \in I_2.$$

$$A_1^c = \left\{ (i, j) \in \mathbb{N} \times \mathbb{N} : \mu\left(T(x_{ij}) - L_1, \frac{t}{2|\alpha|}\right) > 1 - \epsilon \text{ or } \nu\left(T(x_{ij}) - L_1, \frac{t}{2|\alpha|}\right) < \epsilon \right\} \in \mathcal{F}(I_2);$$

$$A_2^c = \left\{ (i, j) \in \mathbb{N} \times \mathbb{N} : \mu\left(T(y_{ij}) - L_2, \frac{t}{2|\beta|}\right) > 1 - \epsilon \text{ or } \nu\left(T(y_{ij}) - L_2, \frac{t}{2|\beta|}\right) < \epsilon \right\} \in \mathcal{F}(I_2).$$

Define the set $A_3 = A_1 \cup A_2$, so that $A_3 \in I_2$. It follows that A_3^c is a non-empty set in $\mathcal{F}(I_2)$. We shall show that for each $(x_{ij}), (y_{ij}) \in {}_2S_{(\mu, \nu)}^I(T)$.

$$A_3^c \subset \left\{ (i, j) \in \mathbb{N} \times \mathbb{N} : \mu\left((\alpha T(x_{ij}) + \beta T(y_{ij})) - (\alpha L_1 + \beta L_2), t\right) > 1 - \epsilon \text{ or } \nu\left((\alpha T(x_{ij}) + \beta T(y_{ij})) - (\alpha L_1 + \beta L_2), t\right) < \epsilon \right\}.$$

Let $(m, n) \in A_3^c$. In this case

$$\mu\left(T(x_{mn}) - L_1, \frac{t}{2|\alpha|}\right) > 1 - \epsilon \text{ or } \nu\left(T(x_{mn}) - L_1, \frac{t}{2|\alpha|}\right) < \epsilon$$

and

$$\mu\left(T(y_{mn}) - L_2, \frac{t}{2|\beta|}\right) > 1 - \epsilon \text{ or } \nu\left(T(y_{mn}) - L_2, \frac{t}{2|\beta|}\right) < \epsilon.$$

We have

$$\begin{aligned} & \mu\left((\alpha T(x_{mn}) + \beta T(y_{mn})) - (\alpha L_1 + \beta L_2), t\right) \\ & \geq \mu\left(\alpha T(x_{mn}) - \alpha L_1, \frac{t}{2}\right) * \mu\left(\beta T(x_{mn}) - \beta L_2, \frac{t}{2}\right) \\ & = \mu\left(T(x_{mn}) - L_1, \frac{t}{2|\alpha|}\right) * \mu\left(T(x_{mn}) - L_2, \frac{t}{2|\beta|}\right) \\ & > (1 - \epsilon) * (1 - \epsilon) = 1 - \epsilon. \end{aligned}$$

and

$$\begin{aligned} & \nu\left((\alpha T(x_{mn}) + \beta T(y_{mn})) - (\alpha L_1 + \beta L_2), t\right) \\ & \leq \nu\left(\alpha T(x_{mn}) - \alpha L_1, \frac{t}{2}\right) \diamond \nu\left(\beta T(x_{mn}) - \beta L_2, \frac{t}{2}\right) \\ & = \mu\left(T(x_{mn}) - L_1, \frac{t}{2|\alpha|}\right) \diamond \mu\left(T(x_{mn}) - L_2, \frac{t}{2|\beta|}\right) \\ & < \epsilon \diamond \epsilon = \epsilon. \end{aligned}$$

This implies that

$$A_3^c \subset \left\{ (i, j) \in \mathbb{N} \times \mathbb{N} : \mu\left((\alpha T(x_{ij}) + \beta T(y_{ij})) - (\alpha L_1 + \beta L_2), t\right) > 1 - \epsilon \text{ or } \nu\left((\alpha T(x_{ij}) + \beta T(y_{ij})) - (\alpha L_1 + \beta L_2), t\right) < \epsilon \right\}.$$

Hence ${}_2S_{(\mu, \nu)}^I(T)$ is a linear space.

Theorem 3.2 Every open ball ${}_2B_x(r, t)(T)$ is an open set in ${}_2S_{(\mu, \nu)}^I(T)$.

Proof. Let ${}_2B_x(r, t)(T)$ be an open ball with centre x and radius r with respect to t . That is

$${}_2B_x(r, t)(T) = \{y = (y_{ij}) \in {}_2\ell_\infty : \{(i, j) \in \mathbb{N} \times \mathbb{N} : \mu(T(x_{ij}) - T(y_{ij}), t) \leq 1 - r \text{ or } \nu(T(x_{ij}) - T(y_{ij}), t) \geq r\} \in I_2\}.$$

Let $y \in {}_2B_x^c(r, t)(T)$. Then $\mu(T(x_{ij}) - T(y_{ij}), t) > 1 - r$ and $\nu(T(x_{ij}) - T(y_{ij}), t) < r$. Since $\mu(T(x_{ij}) - T(y_{ij}), t) > 1 - r$, there exists $t_0 \in (0, t)$ such that $\mu(T(x_{ij}) - T(y_{ij}), t_0) > 1 - r$ and $\nu(T(x_{ij}) - T(y_{ij}), t_0) < r$. Putting $r_0 = \mu(T(x_{ij}) - T(y_{ij}), t_0)$, we have $r_0 > 1 - r$, there exists $s \in (0, 1)$ such that $r_0 > 1 - s > 1 - r$. For $r_0 > 1 - s$, we have $r_1, r_2 \in (0, 1)$ such that $r_0 * r_1 > 1 - s$ and $(1 - r_0) \diamond (1 - r_0) \leq s$. Putting $r_3 = \max\{r_1, r_2\}$. Consider the ball ${}_2B_y^c(1 - r_3, t - t_0)(T)$. We prove that

${}_2B_y^c(1 - r_3, t - t_0)(T) \subset {}_2B_x^c(r, t)(T)$. Let $z = (z_{ij}) \in {}_2B_y^c(1 - r_3, t - t_0)(T)$, then $\mu(T(y_{ij}) - T(z_{ij}), t - t_0) > r_3$ and $\nu(T(y_{ij}) - T(z_{ij}), t - t_0) < 1 - r_3$. Therefore

$$\begin{aligned} \mu(T(x_{ij}) - T(z_{ij}), t) &\geq \mu(T(x_{ij}) - T(y_{ij}), t_0) * \mu(T(y_{ij}) - T(z_{ij}), t - t_0) \\ &\geq (r_0 * r_3) \geq (r_0 * r_1) \geq (1 - s) \geq (1 - r) \end{aligned}$$

and

$$\begin{aligned} \nu(T(x_{ij}) - T(z_{ij}), t) &\leq \nu(T(x_{ij}) - T(y_{ij}), t_0) \diamond \nu(T(y_{ij}) - T(z_{ij}), t - t_0) \\ &\leq (1 - r_0) \diamond (1 - r_3) \leq (1 - r_0) \diamond (1 - r_2) \leq s \leq r. \end{aligned}$$

Thus $z \in {}_2B_x^c(r, t)(T)$ and hence

$${}_2B_y^c(1 - r_3, t - t_0)(T) \subset {}_2B_x^c(r, t)(T).$$

Remark 3.3 ${}_2S_{(\mu, \nu)}^I(T)$ is an IFNS.

Define ${}_2\tau_{(\mu, \nu)}^I(T) = \{A \subset {}_2S_{(\mu, \nu)}^I(T) : \text{for each } x \in A \text{ there exists } t > 0$

and $r \in (0, 1)$ such that ${}_2B_x(r, t)(T) \subset A\}$.

Then ${}_2\tau_{(\mu, \nu)}^I(T)$ is a topology on ${}_2S_{(\mu, \nu)}^I(T)$.

Theorem 3.4 The topology ${}_2\tau_{(\mu, \nu)}^I(T)$ on ${}_2S_{0(\mu, \nu)}^I(T)$ is first countable.

Proof. $\{{}_2B_x(\frac{1}{n}, \frac{1}{n})(T) : n = 1, 2, 3, \dots\}$ is a local base at x , the topology ${}_2\tau_{(\mu, \nu)}^I(T)$ on ${}_2S_{0(\mu, \nu)}^I(T)$ is first countable.

Theorem 3.5 ${}_2S_{(\mu, \nu)}^I(T)$ and ${}_2S_{0(\mu, \nu)}^I(T)$ are Hausdorff spaces.

Proof. We prove the result for ${}_2S_{(\mu, \nu)}^I(T)$. Similarly the proof follows for ${}_2S_{0(\mu, \nu)}^I(T)$.

Let $x, y \in {}_2S_{(\mu, \nu)}^I(T)$ such that $x \neq y$. Then $0 < \mu(T(x) - T(y), t) < 1$ and $0 < \nu(T(x) - T(y), t) < 1$. Putting $r_1 = \mu(T(x) - T(y), t)$, $r_2 = \nu(T(x) - T(y), t)$ and $r = \max\{r_1, 1 - r_2\}$. For each $r_0 \in (r, 1)$ there exists r_3 and r_4 such that $r_3 * r_4 \geq r_0$ and $(1 - r_3) \diamond (1 - r_4) \leq (1 - r_0)$. Putting $r_5 = \max\{r_3, 1 - r_4\}$ and consider the open balls ${}_2B_x(1 - r_5, \frac{t}{2})$ and ${}_2B_y(1 - r_5, \frac{t}{2})$. Then clearly ${}_2B_x^c(1 - r_5, \frac{t}{2}) \cap {}_2B_y^c(1 - r_5, \frac{t}{2}) = \phi$. For if there exists $z \in {}_2B_x^c(1 - r_5, \frac{t}{2}) \cap {}_2B_y^c(1 - r_5, \frac{t}{2})$, then

$$r_1 = \mu(T(x) - T(y), t) \geq \mu(T(x) - T(z), \frac{t}{2}) * \mu(T(z) - T(y), \frac{t}{2}) \geq r_5 * r_5 \geq r_3 * r_3 \geq r_0 > r_1$$

and $r_2 = \nu(T(x) - T(y), t) \leq \nu(T(x) - T(z), \frac{t}{2}) \diamond \nu(T(z) - T(y), \frac{t}{2})$

$$\leq (1 - r_5) \diamond (1 - r_5) \leq (1 - r_4) \diamond (1 - r_4) \leq (1 - r_0) < r_2$$

which is a contradiction. Hence ${}_2S_{(\mu, \nu)}^I(T)$ is Hausdorff.

Theorem 3.6 ${}_2S_{(\mu, \nu)}^I(T)$ is an IFNS and ${}_2\tau_{(\mu, \nu)}^I(T)$ is a topology on ${}_2S_{(\mu, \nu)}^I(T)$.

Then a sequence $(x_{ij}) \in {}_2S_{(\mu, \nu)}^I(T)$, $x_{ij} \rightarrow x$ if and only if $\mu(T(x_{ij}) - T(x), t) \rightarrow 1$ and $\nu(T(x_{ij}) - T(x), t) \rightarrow 0$ as $k \rightarrow \infty$.

Proof. Fix $t_0 > 0$. Suppose $x_{ij} \rightarrow x$. Then for $r \in (0, 1)$, there exists $(m_0, n_0) \in \mathbb{N} \times \mathbb{N}$ such that $(x_{ij}) \in {}_2B_x(r, t)(T)$ for all $(i, j) \geq (m_0, n_0)$,

$${}_2B_x(r, t)(T) = \{(i, j) \in \mathbb{N} \times \mathbb{N} : \mu(T(x_{ij}) - T(x), t) \leq 1 - r \text{ or } \nu(T(x_{ij}) - T(x), t) \geq r\} \in I_2,$$

such that ${}_2B_x^c(r, t)(T) \in \mathcal{F}(I_2)$. Then $1 - \mu(T(x_{ij}) - T(x), t) < r$ and $\nu(T(x_{ij}) - T(x), t) < r$. Hence $\mu(T(x_{ij}) - T(x), t) \rightarrow 1$ and $\nu(T(x_{ij}) - T(x), t) \rightarrow 0$ as $i \rightarrow \infty, j \rightarrow \infty$.

Conversely, if for each $t > 0, \mu(T(x_{ij}) - T(x), t) \rightarrow 1$ and $\nu(T(x_{ij}) - T(x), t) \rightarrow 0$ as $i \rightarrow \infty, j \rightarrow \infty$, then for $r \in (0, 1)$, there exists $(m_0, n_0) \in \mathbb{N} \times \mathbb{N}$ such that $1 - \mu(T(x_{ij}) - T(x), t) < r$ and $\nu(T(x_{ij}) - T(x), t) < r$, for all $i \geq m_0, j \geq n_0$. It follows that $\mu(T(x_{ij}) - T(x), t) > 1 - r$ and $\nu(T(x_{ij}) - T(x), t) < r$ for all $i \geq m_0, j \geq n_0$. Thus $(x_{ij}) \in {}_2B_x^c(r, t)(T)$ for all $i \geq m_0, j \geq n_0$ and hence $x_{ij} \rightarrow x$.

Theorem 3.7 A sequence $x = (x_{ij}) \in {}_2S_{(\mu, \nu)}^I(T)$ is I_2 -convergent if and only if for every $\epsilon > 0$ and $t > 0$ there exists numbers $M = M(x, \epsilon, t)$ and $N = N(x, \epsilon, t)$ such that

$$\{(M, N) \in \mathbb{N} \times \mathbb{N} : \mu(T(x_{MN}) - L, \frac{t}{2}) > 1 - \epsilon \text{ or } \nu(T(x_{MN}) - L, \frac{t}{2}) < \epsilon\} \in \mathcal{F}(I_2).$$

Proof. Suppose that $I_2^{(\mu, \nu)} - \lim x = L$ and let $\epsilon > 0$ and $t > 0$. For a given $\epsilon > 0$, choose $s > 0$ such that $(1 - \epsilon) * (1 - \epsilon) > 1 - s$ and $\epsilon \diamond \epsilon < s$. Then for each $x \in {}_2S_{(\mu, \nu)}^I(T)$,

$$A = \{(i, j) \in \mathbb{N} \times \mathbb{N} : \mu(T(x_{ij}) - L, \frac{t}{2}) \leq 1 - \epsilon \text{ or } \nu(T(x_{ij}) - L, \frac{t}{2}) \geq \epsilon\} \in I_2,$$

which implies that

$$A^c = \{(i, j) \in \mathbb{N} \times \mathbb{N} : \mu(T(x_{ij}) - L, \frac{t}{2}) > 1 - \epsilon \text{ or } \nu(T(x_{ij}) - L, \frac{t}{2}) < \epsilon\} \in \mathcal{F}(I_2).$$

Conversely let us choose $(M, N) \in A$. Then

$$\mu(T(x_{MN}) - L, \frac{t}{2}) > 1 - \epsilon \text{ or } \nu(T(x_{MN}) - L, \frac{t}{2}) < \epsilon.$$

Now we want to show that there exists a number $M = M(x, \epsilon), N = N(x, \epsilon, t)$ such that

$$\{(M, N) \in \mathbb{N} \times \mathbb{N} : \mu(T(x_{ij}) - T(x_{MN}), t) \leq 1 - s \text{ or } \nu(T(x_{ij}) - T(x_{MN}), t) \geq s\} \in I_2.$$

For this, define for each $x \in {}_2S_{(\mu, \nu)}^I(T)$

$$B = \{(i, j) \in \mathbb{N} \times \mathbb{N} : \mu(T(x_{ij}) - T(x_{MN}), t) \leq 1 - s \text{ or } \nu(T(x_{ij}) - T(x_{MN}), t) \geq s\} \in I_2.$$

Now we have to show that $B \subset A$. Suppose that BA . Then there exists $(m, n) \in B$ and $n \notin A$. Therefore we have

$$\mu(T(x_{mn}) - T(x_{MN}), t) \leq 1 - s \text{ or } \mu(T(x_{mn}) - L, \frac{t}{2}) > 1 - \epsilon.$$

In particular $\mu(T(x_{MN}) - L, \frac{t}{2}) > 1 - \epsilon$. Therefore we have

$$1 - s \geq \mu(T(x_{mn}) - T(x_{MN}), t) \geq \mu(T(x_{mn}) - L, \frac{t}{2}) * \mu(T(x_{MN}) - L, \frac{t}{2}) \geq (1 - \epsilon) * (1 - \epsilon) > 1 - s,$$

which is not possible. On the other hand

$$\nu(T(x_{mn}) - T(x_{MN}), t) \geq s \text{ or } \nu(T(x_{mn}) - L, \frac{t}{2}) < \epsilon$$

In particular $\nu(T(x_{MN}) - L, \frac{t}{2}) < \epsilon$. Therefore we have

$$s \leq \nu(T(x_{mn}) - T(x_{MN}), t) \leq \nu(T(x_{mn}) - L, \frac{t}{2}) \diamond \nu(T(x_{MN}) - L, \frac{t}{2}) \leq \epsilon \diamond \epsilon < s,$$

which is not possible. Hence $B \subset A$. $A \in I_2$ implies $B \in I_2$.

4. CONCLUSION:

In the present paper we have studied the concept of ideal convergence of double sequence spaces in intuitionistic fuzzy sequence spaces defined by compact operator and studied the fuzzy topology on the said spaces.

5. AUTHORS CONTRIBUTIONS

All authors of the manuscript have read and agreed to its content and are accountable for all aspects of the accuracy and integrity of the manuscript.

6. AUTHOR DETAILS

Vakeel A. Khan received the M.Phil. and Ph.D. degrees in Mathematics from Aligarh Muslim University, Aligarh, India. Currently he is a Associate Professor at Aligarh Muslim University, Aligarh, India. A vigorous researcher in the area of Sequence Spaces, he has published a number of research papers in reputed national and international journals, including Numerical Functional Analysis and Optimization (Taylors and Francis), Information Sciences (Elsevier), Applied Mathematics Letters (Elsevier), A Journal of Chinese Universities (Springer- Verlag, China).

Yasmeen recieved M.Sc and M.Phil degrees from Aligarh Muslim University, and is currently a Ph.D., scholar at Aligarh Muslim University, Aligarh, India.

Ayhan Esi got his B.Sc. from Inonu University in 1987 and M. Sc. and Ph.D. degree in pure mathematics from Elazig University, Turkey in 1990 and 1995, respectively. Currently he is a Professor at Department of Mathematics of Adiyaman University. His research interests include Summability Theory, Sequences and Series in Analysis and Functional Analysis.

Hira Fatima received B.Sc and M.Sc. degrees from Aligarh Muslim University, and is currently a Ph.D. scholar at Aligarh Muslim University, Aligarh, India.

Mobeen Ahamad recieved M.Sc and M.Phil degrees from Aligarh Muslim University, and is currently a Ph.D., scholar at Aligarh Muslim University, Aligarh, India.

7. COMPETING INTERESTS

The authors declare that they have no competing interests.

8. ACKNOWLEDGMENT

The authors would like to record their gratitude to the reviewer for his/her careful reading and making some useful corrections which improved the presentation of the paper.

REFERENCES

- [1] L. C. Barros, R. C. Bassanezi, P. A. Tonelli, Fuzzy modelling in population dynamics, *Ecol. Model.*, 128(2000)27-33.
- [2] İ. Çanak, Ü. Totur, Z.Önder, A Tauberian theorem for $(C,1,1)$ summable double sequences of fuzzy numbers, *Iran. J. Fuzzy Syst.*, 14 (1), 61-75 (2017).
- [3] H. Fast, Sur la convergence statistique, *Colloq. Math.*,2(1951), 241-244.
- [4] A. L. Fradkov, R. J. Evans, Control of chaos: Methods of applications in engineering, *Chaos, solution and Fractals* (29)(2005),33-56.
- [5] R. Giles, A computer program for fuzzy reasoning, *Fuzzy sets and systems* (4)(1980),221-234.
- [6] L. Hong, J. Q. Sun, Bifurcations of fuzzy non-linear dynamical systems, *Commun. Nonlinear Sci.Numer. Simul.*, (1)(2006), 1-12.
- [7] V. A. Khan, K. Ebadullah and Yasmeen, On Zweier I -convergent sequence spaces, *Proyecciones Journal of Mathematics* Vol.(3)(33)(2014),259-276.
- [8] V. A. Khan, K. Ebadullah and R.K.A. Rababah, Intuitionistic fuzzy zweier I -convergent sequence spaces, *Functinal Analysis: Theory, methods and applications*, Vol. (1)(2015), 1-7.
- [9] V. A. Khan, Yasmeen, On Paranorm Type Intuitionistic Fuzzy Zweier I -convergent Sequence Spaces, *Annals of Fuzzy Mathematics and Informatics*(Article in press).
- [10] V. A. Khan, Yasmeen, Intuitionistic Fuzzy Zweier I -convergent Double Sequence Spaces, *New Trends in Mathematical Sciences*, Volume: 4, Issue: 2, (MAR 2016), pp:240-247.
- [11] V. A. Khan, Yasmeen, Intuitionistic Fuzzy Zweier I -convergent Sequence Spaces defined by Modulus function, *Cogent mathematics* (Taylors and Francis),vol . 3 Issue 2 (2016).
- [12] V. A. Khan, Yasmeen, Intuitionistic Fuzzy Zweier I -convergent Sequence Spaces defined by Orlicz function, *Annals of Fuzzy Mathematics and Informatics*, Vol. 12, 469-478(2016).
- [13] Vakeel A. Khan, Yasmeen, Hira Fatima, Henna Altaf and Q.M. Danish Lohani, Intuitionistic Fuzzy I -convergent Sequence Spaces defined by Compact Operator, *Cogent Mathematics* (2016), 3: 1267904.
- [14] V. A. Khan, Mohd. Shafiq and Bernardo Lafuerza-Guillen, On paranorm I -convergent sequence spaces defined by a compact operator, *Afrika Matematika, Journal of the African Mathematical Union*(Springer), vol.25, Issue 4(2014), 12 Pages, DOI 10.1007/s13370-014-0287-2.
- [15] P. Kostyrko, T. Salat and W. Wilczynski, I -convergence, *Real Analysis Exchange* (26)(2000), No. 2, 669-686.
- [16] P. Das, P. Kostyrko, W. Wilczynski, P. Malik, I and I^* - convergence of double sequences, *Math. Slovaca* (58)(2008), 605-620.
- [17] E. Kreyszig, *Introductory functional analysis with application*, John Wiley and Sons, Inc. New York-Chicheste-Brisbane-Toronto, 1978.
- [18] M. Mursaleen, Osama H. H. Edely, Statistical convergence of double sequences, *J. Math. Anal. Appl.* (288)(2003), 223-231.
- [19] M. Mursaleen and S. A. Mohiuddine, On ideal convergence in probabilistic normed spaces, *Mathematica Slovaca*, 62(1) (2012) 49-62.
- [20] M. Mursaleen and S. A. Mohiuddine, On ideal convergence of double sequences in probabilistic normed spaces, *Mathematical Reports*, 12(62) 4(2010) 359-371.
- [21] S. A. Mohiuddine, Alotaibi, and S. M. Alsulami, Ideal convergence of double sequences in random 2-normd spaces, *Advances in Difference Equations*, vol. 2012, article 149, 8 pages(2012).
- [22] B. Hazarika and S. A. Mohiuddine, Ideal convergence of random variables, *Journal of Function Spaces and Applications*, vol. 2013, Article ID 148249, 7 pages, 2013.
- [23] A. Alotaibi, B. Hazarika and S. A. Mohiuddine, On the ideal convergence of double sequences in locally solid Riesz spaces, *Abstract and Applied Analysis*, Vol 2014, Article ID 396254, 6 pages (2014).
- [24] M. Mursaleen and S. A. Mohiuddine, On lacunary statistical convergence with respect to the intuitionistic fuzzy normed spaces, *Journal of Computational and Applied Mathematics* 233 (2009) 142-149.
- [25] M. Mursaleen and S. A. Mohiuddine, Statistical convergence of double sequences in intuitionistic fuzzy normed spaces, *Chaos, Solitons and Fractals* 42(2009) 1731-1737.
- [26] S. A. Mohiuddine and Q. M. D. Lohani, On generalized statistical convergence in intuitionistic fuzzy normed spaces, *Chaos, Solitons and Fractals* 41(2009) 142-149.

- [27] M. Mursaleen, Q.M.D. Lohni, Intuitionistic fuzzy 2-normed space and some related concepts, Chaos, solution and Fractals (42)(2009), 331-344.
- [28] M. Mursaleen, S.A. Mohiuddine and Osama H.H. Edely, On the ideal convergence of double sequences in intuitionistic fuzzy normed spaces, Computers and Mathematics with Application 59(2010)603-611.
- [29] M. Mursaleen, H. M. Srivastava and S. K. Sharma, Generalized statistically convergent sequences of fuzzy numbers, J. Intelligent Fuzzy System 30 (2016), 1511-1518.
- [30] A. Nabiev, S. Pehlivan, M. Gürdal, On I - Cauchy sequence, Taiwanese J. Math. (11)(2)(2007), 569-576.
- [31] Z. Önder, İ. Çanak, Ü. Totur, Tauberian theorems for statistically $(C,1,1)$ summable double sequences of fuzzy numbers, Open Math., 15 (1), 157-178 (2017).
- [32] R. Saddati, J. H. Park, On the intuitionistic fuzzy topological spaces, Chaos, solution and Fractals (27)(2006), 331-344.
- [33] E. Savaş, M. Mursaleen, On statistical convergent double sequences of fuzzy numbers, Inform. Sci. 162(2004), 183-192.
- [34] Ü. Totur, İ. Çanak, Tauberian theorems for the statistical convergence and the statistical $(C,1,1)$ summability, Filomat, 32 (1), 101-116 (2018).
- [35] L. A. Zadeh, Fuzzy sets, Inform Control, (8)(1965), 338-353. 1, 1.8.

VAKEEL A. KHAN

DEPARTMENT OF MATHEMATICS, ALIGARH MUSLIM UNIVERSITY, ALIGARH-202002, INDIA.

E-mail address: vakhanmaths@gmail.com

YASMEEN

DEPARTMENT OF MATHEMATICS, ALIGARH MUSLIM UNIVERSITY, ALIGARH-202002, INDIA.

E-mail address: yasmeen@gmail.com

AYHAN ESI

DEPARTMENT OF MATHEMATICS, ADIYAMAN UNIVERSITY, ADIYAMAN , TURKEY.

E-mail address: aesi23@hotmail.com

HIRA FATIMA

DEPARTMENT OF MATHEMATICS, ALIGARH MUSLIM UNIVERSITY, ALIGARH-202002, INDIA.

E-mail address: hirafatima2014@gmail.com

MOBEEN AHAMAD

DEPARTMENT OF MATHEMATICS, ALIGARH MUSLIM UNIVERSITY, ALIGARH-202002, INDIA.

E-mail address: gc4889@myamu.ac.in