

PROPERTIES OF SOME NEW BILATERAL MOCK THETA FUNCTIONS

MOHAMMAD AHMAD, SIRAZUL HAQ AND ABDUL HAKIM KHAN

ABSTRACT. Bilateral mock theta functions were obtained and studied in [10]. We express them in terms of Lerch's transcendental function $f(x, \xi; q, p)$. We also express some bilateral mock theta functions as sum of other mock theta functions. We generalize these functions and show that these generalizations are F_q functions. We give an integral representation for these generalized functions.

1. INTRODUCTION

The mock theta functions were first introduced by Ramanujan [3] in his last letter to G H Hardy in January 1920 . He provided a list of seventeen mock theta functions and labeled them as of third, fifth and seventh order without mentioning the reason for his labeling. Watson [18] added to this set three more third order mock theta functions. His general definition of a mock theta functions is a functions $f(q)$ defined by a q series convergent when $q < 0$ which satisfies the following two conditions.

(a) For every root ξ of unity, there exists a theta function $\theta_\xi(q)$ such that the difference between $f(q)$ and $\theta_\xi(q)$ is bounded as $q \rightarrow \xi$ radially.

(b) there is no single theta function which works for all ξ i.e for every theta function $\theta_\xi(q)$ there is some root of unity ξ for which $f(q)$ minus the theta function $\theta_\xi(q)$ is unbounded as $q \rightarrow \xi$ radially.

Andrews and Hickerson [15] announced the existence of eleven more identities given in the 'lost' note book of Ramanujan involving seven new functions which they labeled as mock theta functions of order six. Y.S. Choi [1] has discovered four functions which he called the mock theta function of order ten. B.Gordon and R.J.McIntosh [30] have announced the existence of eight mock theta functions of order eight and R.J.McIntosh [5] has announced the existence of three mock theta

2010 *Mathematics Subject Classification.* 33D15, 11B65.

Key words and phrases. Mock theta functions, bilateral mock theta functions, Lerch transcendent, F_q -function.

Submitted Feb. 18, 2019. Revised Aug. 24, 2019.

functions of order two.

Hikami [13, 14] has introduced one mock theta function of order two, one of order four and two of order eight. Very recently Andrews [16] while studying q -orthogonal polynomials found four new mock theta functions and Bringmann et al [12] have also found two more new mock theta functions but they did not mention about the order of their mock theta functions.

Watson [19] has defined four bilateral series, which he called the 'Complete' or 'Bilateral' forms for four of the ten mock theta functions of order five and expressed them in terms of the transcendental functions $f(x, \xi; q, p)$ studied by M. Lerch [7]. S.D. Prasad [2] in 1970 has defined the 'Complete' or 'Bilateral' forms of the five generalized third order mock theta functions. The 'Complete' sixth order mock theta functions were studied by A. Gupta [31] Bhaskar Srivastava [26, 27, 28, 29] have studied bilateral mock theta functions of order five, eight, two and new mock theta functions by Andrew [6] and Bringmann et al [12].

Truesdell [25] calls the functions which satisfy the equations $\frac{d}{dz}F(z, \alpha) = F(z, \alpha + 1)$ as F -functions. He has tried to unify the study of these F -functions. The function which satisfy the q -analogue of the equation $D_{q,z}F(z, \alpha) = F(z, \alpha + 1)$ where $zD_{q,z}F(z, \alpha) = F(z, \alpha) - F(zq, \alpha)$ are called F_q -functions.

Mohammad Ahmad, Sirazul Haq and Abdul Hakim Khan have obtained and studied the following bilateral mock theta functions in [10].

$$f_{0,cs}(q) = \sum_{-\infty}^{\infty} \frac{q^{4n^2-3n}}{(-q; q)_n} \tag{1}$$

$$f_{1,cs}(q) = \sum_{-\infty}^{\infty} \frac{q^{4n^2-2n}}{(-q; q)_n} \tag{2}$$

$$F_{0,cs}(q^2) = \sum_{-\infty}^{\infty} \frac{q^{8n^2-6n}}{(q; q^2)_n} \tag{3}$$

$$F_{1,cs}(q^4) = \sum_{-\infty}^{\infty} \frac{q^{16n^2-8n}}{(q^6; q^4)_n} \tag{4}$$

$$\psi_{0,cs}(q) = \sum_{-\infty}^{\infty} q^{\frac{7(n^2+3n)}{2}} (-q; q)_n \tag{5}$$

$$\phi_{1,cs}(q^2) = \sum_{-\infty}^{\infty} q^{7n^2+14n} (-q; q^2)_n \tag{6}$$

$$\phi_{0,cs}(q^2) = \sum_{-\infty}^{\infty} \frac{q^{8n^2}}{(-q; q^2)_n} \tag{7}$$

$$\psi_{1,cs}(q) = \sum_{-\infty}^{\infty} \frac{q^{4n^2+4n}}{2(-q; q)_n} \tag{8}$$

The paper is divided as follows: In section 2 we list few important definitions. In section 3 we develop certain properties of these functions by expressing some of them as sums of other mock theta functions. We also express these functions in terms of the Lerch transcendental function $f(x, \xi; q, p)$. In section 4 we generalize these functions which are then proved to be F_q functions. We further give an integral representation of these functions.

2. NOTATION AND DEFINITIONS

We use the following q notation. Suppose q and z are complex numbers and n is an integer if $n \geq 0$ we define $(z)_n = (z, q)_n = \prod_{i=0}^{n-1} (1 - q^i z)$ if $n \leq 0$ and

$$(z)_{-n} = (z; q)_{-n} = \frac{(-z)^{-n} q^{\frac{n(n+1)}{2}}}{(q/z; q)_n}$$

and more generally

$$(z_1, z_2, \dots, z_r; q)_n = (z_1)_n (z_2)_n, \dots, (z_r)_n$$

For $|q^k| < 1$ let us define

$$(z; q^k)_n = (1 - z)(1 - z q^k), \dots, (1 - z q^{k(n-1)}) \quad n \geq 1 \quad (z; q^k)_0 = 1$$

$$(z; q^k)_\infty = \lim_{n \rightarrow \infty} (z; q^k)_n = \prod_{i \geq 0} (1 - q^{ki} z)$$

and more generally

$$(z_1, z_2, \dots, z_r; q^k)_\infty = (z_1; q^k)_\infty, \dots, (z_r; q^k)_\infty$$

A basic hypergeometric series ${}_{r+1}\phi_r$ on a base q^k is defined as

$${}_{r+1}\phi_r \left[\begin{matrix} a_1, a_1, \dots, a_r, ; \\ b_1, b_2, \dots, b_r \end{matrix} ; q^k, z \right] = \sum_{n=0}^{\infty} \frac{(a_1, a_2, \dots, a_r; q^k)_n z^n}{(q^k; q^k)_n (b_1, b_2, \dots, b_r; q^k)_n}, \quad (|z| < 1)$$

and a bilateral basic hypergeometric series ${}_r\psi_r$ is defined as

$${}_r\psi_r \left[\begin{matrix} a_1, a_1, \dots, a_r, ; \\ b_1, b_2, \dots, b_r \end{matrix} ; q, z \right] = \sum_{n=-\infty}^{\infty} \frac{(a_1, a_2, \dots, a_r; q^k)_n z^n}{(b_1, b_2, \dots, b_r; q^k)_n}, \quad \left(\left| \frac{b_1, \dots, b_r}{a_1, \dots, a_r} \right| < |z| < 1 \right)$$

Lerch transcendental function $f(x, \xi; q, p)$ is defined by :

$$f(x, \xi; q, p) = \sum_{-\infty}^{\infty} \frac{(p q)^{n^2} (x \xi)^{-2n}}{(-p \xi^{-2}; p^2)_n}$$

and by

$$f(x, \xi; q, p) = \sum_{-\infty}^{\infty} (-\xi^2 p; p^2)_n q^{n^2} x^{2n}$$

3. CERTAIN IDENTITIES AND LERCH REPRESENTATIONS

The following identities between the bilateral mock theta functions given in equations (1), (5), (6), (7) and the corresponding mock theta functions may be verified by hypergeometric transformations:

$$f_{0,c_8}(q) = f_{0,8}(q) + 2q^7\psi_{0,8}(q) \tag{9}$$

$$\phi_{0,c_8}(q^2) = \phi_{0,8}(q^2) + q^7\phi_{1,8}(q^2) \sum_0^\infty (1 + q^{2n+1}) \tag{10}$$

Here $f_{0,8}(q), \psi_{0,8}(q), \phi_{0,8}(q^2), \phi_{1,8}(q^2)$ are the corresponding mock theta functions. The bilateral mock theta functions defined in section 1 can be expressed in terms of the Lerch transcendent by means of the following lemma.

Lemma 3.1: For $\epsilon = \pm 1$,

$$\sum_{n=-\infty}^\infty \frac{q^\alpha n^2 q^{\beta n}}{(\epsilon q^\gamma; q^\delta)_n} = f\left((-\epsilon)^{-1/2} q^{\frac{2\gamma-2\beta-\delta}{4}}, (-\epsilon)^{1/2} q^{\frac{\delta-2\gamma}{2}}; q^{\frac{2\alpha-\delta}{2}}, q^{\delta/2} \right)$$

$$\sum_{n=-\infty}^\infty (-q; q^\gamma)_n q^{\alpha n^2} q^{\beta n} = f(q^{\beta/2} q^{\gamma/2}; q^\alpha, q^{\gamma/2}).$$

Proof. The proof follows from direct substitution and use of basic hypergeometric transformations. As an example we note that $f_{1,c_8}(q) = \sum_{-\infty}^\infty \frac{q^{4n^2-2n}}{(-q; q)_n} = f(q^{5/4}, q^{-1/4}, q^{7/2}, q^{1/2})$ by taking $\alpha = 4, \beta = -2, \epsilon = -1, \gamma = \delta = 1$ and $\phi_{1,c_8}(q^2) = \sum_{-\infty}^\infty q^{7n^2+14n} (-q; q^2)_n = f(q^7, 1, q^7, q)$ by taking $\alpha = 7, \beta = 14, \gamma = 2$ in the above lemma. In this way all other bilateral mock theta functions defined by equations (1) to (8) can be expressed in terms of the Lerch transcendent. \square

4. GENERALIZATION OF BILATERAL MOCK THETA FUNCTIONS

: We generalize the functions given by Equations (1) to (8) by introducing two parameters α, z . For $\alpha = 1, z = 0$ these are reduced to the original functions.

$$f_{0,c_8}(z, \alpha; q) = \frac{1}{(z)_\infty} \sum_{-\infty}^\infty \frac{(z)_n q^{4n^2+n\alpha-4n}}{(-q; q)_n} \tag{11}$$

$$f_{1,c_8}(z, \alpha; q) = \frac{1}{(z)_\infty} \sum_{-\infty}^\infty \frac{(z)_n q^{4n^2+n\alpha-3n}}{(-q; q)_n} \tag{12}$$

$$F_{0,c_8}(z, \alpha; q^2) = \frac{1}{(z)_\infty} \sum_{-\infty}^\infty \frac{(z)_n q^{8n^2+n\alpha-7n}}{(-q; q^2)_n} \tag{13}$$

$$F_{1,c_8}(z, \alpha; q^4) = \frac{1}{(z)_\infty} \sum_{-\infty}^\infty \frac{(z)_n q^{16n^2+n\alpha-9n}}{(q^6; q^4)_n} \tag{14}$$

$$\psi_{0,c_8}(z, \alpha; q) = \frac{1}{(z)_\infty} \sum_{-\infty}^\infty (z)_n q^{\frac{7n^2}{2}+n\alpha+\frac{19n}{2}} (-q; q)_n \tag{15}$$

$$\phi_{1,c_8}(z, \alpha; q^2) = \frac{1}{(z)_\infty} \sum_{-\infty}^{\infty} (z)_n q^{7n^2+n\alpha+13n} (-q; q^2)_n \quad (16)$$

$$\phi_{0,c_8}(z, \alpha; q^2) = \frac{1}{(z)_\infty} \sum_{-\infty}^{\infty} \frac{(z)_n q^{8n^2+n\alpha-n}}{(-q; q^2)_n} \quad (17)$$

$$\psi_{1,c_8}(z, \alpha; q) = \frac{1}{(z)_\infty} \sum_{-\infty}^{\infty} \frac{(z)_n q^{4n^2+n\alpha+3n}}{2(-q; q)_n} \quad (18)$$

We now show that these generalized functions are F_q functions.

Theorem 4.1: The functions defined by the Equations (1)- (18) are F_q functions.

Proof. We give the proof only for $f_{0,c_8}(z, \alpha; q)$. The remaining cases are similar. For $f_{0,c_8}(z, \alpha; q)$ note that

$$\begin{aligned} zD_{q,z}f_{0,c_8}(z, \alpha; q) &= f_{0,c_8}(z, \alpha; q) - f_{0,c_8}(z, \alpha; q) \\ &= \frac{1}{(z)_\infty} \sum_{-\infty}^{\infty} \frac{(z)_n q^{4n^2+n\alpha-4n}}{(-q; q)_n} \\ &\quad - \frac{1}{(zq)_\infty} \sum_{-\infty}^{\infty} \frac{(zq)_n q^{4n^2+n\alpha-4n}}{(-q; q)_n} \\ &= \frac{1}{(z)_\infty} \sum_{-\infty}^{\infty} \frac{(z)_n q^{4n^2+n\alpha-4n}}{(-q; q)_n} (1 - (1 - zq^n)) \\ &= \frac{z}{(z)_\infty} \sum_{-\infty}^{\infty} \frac{(z)_n q^{4n^2+(n+1)\alpha-4n}}{(-q; q)_n} \\ &= f_{0,c_8}(z, \alpha + 1; q) \end{aligned}$$

and hence $f_{0,c_8}(z, \alpha; q)$ is a F_q function. We now give integral representations of the generalized functions. Jackson (on page 23 of [17]) defined the q -integral on $(0, \infty)$ by

$$\int_0^\infty f(t) d_q t = (1-q) \sum_{-\infty}^{\infty} f(q^n) q^n$$

Now let $f(t) = t^{x-1}(tq; q)_\infty$ for some fixed x . we have

$$\begin{aligned} \int_0^\infty t^{x-1}(tq; q)_\infty d_q t &= (1-q) \sum_{-\infty}^{\infty} (q^{n+1}; q)_\infty q^{nx} \\ &= (1-q) \frac{(q; q)_\infty}{(q^x; q)_\infty} \end{aligned}$$

and so

$$\frac{1}{(q^x; q)_\infty} = \frac{(1-q)^{-1}}{(q; q)_\infty} \int_0^\infty t^{x-1}(tq; q)_\infty d_q t. \tag{19}$$

□

We now use equation (19) to give an integral representations of the F_q functions (11) to (18) . We let $a = q^\alpha$ for convenience .

$$f_{0,cs}(q^z, \alpha; q) = \frac{(1-q)^{-1}}{(q; q)_\infty} \int_0^\infty t^{z-1}(tq; q)_\infty f_{0,cs}(0, at; q) d_q t. \tag{20}$$

$$f_{1,cs}(q^z, \alpha; q) = \frac{(1-q)^{-1}}{(q; q)_\infty} \int_0^\infty t^{z-1}(tq; q)_\infty f_{1,cs}(0, at; q) d_q t. \tag{21}$$

$$F_{0,cs}(q^z, \alpha; q^2) = \frac{(1-q)^{-1}}{(q; q)_\infty} \int_0^\infty t^{z-1}(tq; q)_\infty F_{0,cs}(0, at; q^2) d_q t. \tag{22}$$

$$F_{1,cs}(q^z, \alpha; q^4) = \frac{(1-q)^{-1}}{(q; q)_\infty} \int_0^\infty t^{z-1}(tq; q)_\infty F_{1,cs}(0, at; q^4) d_q t. \tag{23}$$

$$\psi_{0,cs}(q^z, \alpha; q) = \frac{(1-q)^{-1}}{(q; q)_\infty} \int_0^\infty t^{z-1}(tq; q)_\infty \psi_{0,cs}(0, at; q) d_q t. \tag{24}$$

$$\phi_{1,cs}(q^z, \alpha; q^2) = \frac{(1-q)^{-1}}{(q; q)_\infty} \int_0^\infty t^{z-1}(tq; q)_\infty \phi_{1,cs}(0, at; q^2) d_q t. \tag{25}$$

$$\phi_{0,cs}(q^z, \alpha; q^2) = \frac{(1-q)^{-1}}{(q; q)_\infty} \int_0^\infty t^{z-1}(tq; q)_\infty \phi_{0,cs}(0, at; q^2) d_q t. \tag{26}$$

$$\psi_{1,cs}(q^z, \alpha; q) = \frac{(1-q)^{-1}}{(q; q)_\infty} \int_0^\infty t^{z-1}(tq; q)_\infty \psi_{1,cs}(0, at; q) d_q t. \tag{27}$$

Theorem 4.2: Equations (20) to (27) hold.

Proof. We prove only (20). The remaining cases are similar. Writing q^z for z and a for q^α in (11) we have,

$$\begin{aligned} f_{0,cs}(q^z, \alpha; q) &= \frac{1}{(q^z; q)_\infty} \sum_{-\infty}^\infty \frac{a^n (q^z; q)_n q^{4n^2-4n}}{(-q; q)_n} \\ &= \sum_{-\infty}^\infty \frac{a^n q^{4n^2-4n}}{(-q; q)_n (q^{n+z}; q)_\infty} \\ &= \frac{(1-q)^{-1}}{(q; q)_\infty} \sum_{-\infty}^\infty \frac{a^n q^{4n^2-4n}}{(-q; q)_n} \int_0^\infty t^{n+z-1}(tq; q)_\infty d_q t \\ &= \frac{(1-q)^{-1}}{(q; q)_\infty} \int_0^\infty t^{z-1}(tq; q)_\infty \sum_{-\infty}^\infty \frac{(at)^n q^{4n^2-4n}}{(-q; q)_n} d_q t \\ &= \frac{(1-q)^{-1}}{(q; q)_\infty} \int_0^\infty t^{z-1}(tq; q)_\infty f_{0,cs}(0, at; q) d_q t \end{aligned}$$

which completes the proof. We remarks that for $at = q$ the function $f_{0,c_8}(0, at; q)$ reduces to the bilateral mock theta function $f_{0,c_8}(q)$ defined previously. \square

5. ACKNOWLEDGEMENT:

Dr Mohammad Ahmad gratefully acknowledges the support and blessings of Air Marshal I P Vipin VM Commandant NDA, Rear Admiral S K Grewal VSM,NM Deputy Commandant and Chief Instructor NDA and Prof. OP Shukla Principal NDA Khadakwasla pune.

REFERENCES

- [1] Y.S. Choi, Tenth order mock theta functions in Ramanujan's lost note book, *Inventiones Mathematicae* 136(1999), 497-569.
- [2] S.D. Prasad, Certain extended mock theta functions and generalized basic hypergeometric transformation *Cambridge Mathematical Library, Math Scand.* 27(1970), 237-244.
- [3] S. Ramanujan, *Collected Papers*, Cambridge University Press (1927), reprinted by Chelsea New York Functi (1960).
- [4] R.P. Agarwal, Mock theta functions- An analytical point of view, *Proc. Nat. Acad. Sci. India* 64(A).I.(1994) 95-106.
- [5] R. J. McIntosh, Second order mock theta functions, *Canadian Mathematical Bulletin* 50(2) (2007), 284-290.
- [6] N. J. Fine, basic hypergeometric series and applications, *Mathematical Surveys and Mono Graphs*, Providence, R.I. Amer. Math. Soc. 27(1988).
- [7] M. Lerch, Nov analogie rady theta a nekte zvltn hypergeometrick rady Heineovy. *Rozpravy*, 3(1893), 1-10.
- [8] M. Ahmad, On the Behaviour of Bilateral Mock Theta Functions-I, *Algebra and Analysis: Theory and Applications*, Narosa Publishing House New Delhi, (2015), 259-274.
- [9] M. Ahmad and Shadab Faruqi, Some Bilateral Theta Function and Their Lerch Representations, *The Aligarh Bulletin of Mathematics*, Vol.34, Numbers1-2(2015)75-92.
- [10] M. Ahmad, Sirazul Haq and Abdul Hakim Khan, Bilateral Mock Theta Functions and Further Properties, to appear in *Electronic Journal of Mathematical Analysis and Application* Vol.7(2) July 2019.
- [11] L. A. Dragonette, Some asymptotic formulae for the mock theta series of Ramanujan, *Trans. Amer. Math. Soc.* 72 (1952),474-500.
- [12] K. Bringmann, K. Hikami and J. Lovejoy, On the modularity of the unified WRT invariants of certain Seifert maifolds, *Adv. Appl. Maths.* 46,1 (2011),86-93.
- [13] K. Hikami, Mock (false) theta functions as quantum invariants, *Regular and Chaotic Dynamic*, 10(2005), 509-530.
- [14] K. Hikami, Transformation formulae of the second order mock theta function, *Lett. Math. Phys.* 75(1), (2006),93-98.
- [15] G. E. Andrews and D. Hickerson, The sixth order mock theta functions, *Adv. Maths.* 89(1991), 60-105.
- [16] G. E. Andrews, q-orthogonal polynomials, Roger-Ramanujan identities and mock theta functions, *Proc. Steklov Institute of Math.* 276(1) (2012),21-32.
- [17] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, Cambridge University Press U. K.(2004).
- [18] G. N. Watson, The final problem, An account of the mock theta functions, *Journal of the London Math, Soc.* 11(1936), 55-80.
- [19] G. N. Watson, The Mock Theta Functions(2), *Proc. London Math, Soc.* 42(1937), 274-304.
- [20] D.P. Shukla and M. Ahmad, Bilateral mock theta functions of order seven, *Math. Sci. Res. J.*7(1)(2003), 8-15.
- [21] D. P. Shukla and M. Ahmad, On the behaviour of bilateral mock theta functions of order seven, *Math. sci. Res. J.* 7(1)(2003), 16-25.
- [22] D. P. Shukla and M. Ahmad, Bilateral mock theta functions of order eleven, *Proc. Jang Jeon Math.Soc*,6(2003),No.1, 59-64.
- [23] D. P. Shikla and M. Ahmad, Bilateral mock theta functions of order thirteen.*Proc. Jang Jeon Math.Soc*,6(2003),No.2, 167-183.

- [24] D.P. Shukla and M. Ahmad, Bilateral mock theta functions of order $2r+1$ and their behaviour of the unit circle, *Ganita* Vol.60, No.1, (2009),27-56.
- [25] Clifford Truesdell. An essay toward a unified theory of special functions, 18 Princeton university press, 1948.
- [26] Bhaskar Srivastava, Certain bilateral basic hypergeometric transformations and mock theta functions, *Hiroshima Maths. J.*29(1999),19-26.
- [27] Bhaskar Srivastava, A Study of bilateral forms of the mock theta functions of order eight, *Journal of the Chungcheon Math Soc.* Vol.18 No.2(2005),117-129.
- [28] Bhaskar Srivastava, A mock theta function of second order, *International Journal of Mathematics and Mathematical science*, Vol.(2009) ArtID978425,15pages.
- [29] Bhaskar Srivastava, A Study of bilateral new mock theta functions, *American Journal of Mathematics and Statistics* (2014)2(4),64-69.
- [30] B. Gordon and R.J. McIntosh, Some eight order mock theta functions, *Journal of the London Math. Soc.* 62(2)(2000),321-335.
- [31] Anju Gupta, On certain Ramanujan's mock theta functions, *Proc. Indian Acad. Sci.* 103(1993), 257-267.

MOHAMMAD AHMAD

DEPARTMENT OF MATHEMATICS, NATIONAL DEFENCE ACADEMY, KHADAKWASLA, PUNE, INDIA

E-mail address: mahmad-786@rediffmail.com

SIRAZUL HAQ

DEPARTMENT OF APPLIED MATHEMATICS, FACULTY OF ENGINEERING AND TECHNOLOGY, ALIGARH

MUSLIM UNIVERSITY, INDIA

E-mail address: sirajulhaq007@gmail.com

ABDUL HAKIM KHAN

DEPARTMENT OF APPLIED MATHEMATICS, FACULTY OF ENGINEERING AND TECHNOLOGY, ALIGARH

MUSLIM UNIVERSITY, INDIA.

E-mail address: ahkhan.amu@gmail.com