

CHEMICAL AND TECHNOLOGICAL STUDIES ON BLACK MULBERRY JUICE AND ITS CONCENTRATE

A. A. El-Bedawe⁽¹⁾, E. H. Rahma⁽¹⁾, E.S. Abd-Elwahab⁽²⁾ and Naglaa M. Ali⁽²⁾

⁽¹⁾ Dept. of Food Science, Faculty of Agric. Minufiya Univ. Egypt

⁽²⁾ Food Technology Res. Inst., Agric. Res. Center, Giza, Egypt

(Received: Dec. 8, 2014)

ABSTRACT: Fresh Black mulberry was mechanically extracted and clarified by using gelatin and (UF) ultrafiltration process. The clarified juice was concentrated using rotary evaporator and freezing process. Physico-chemical properties as well as chemical constituents and microbiological aspects was determined, The juice yield of 62.80 % with a moisture content was 81.20 %, Total soluble solids (TSS%), pH value and acidity were 16.4%, 5.49 and 0.341% respectively. Color index at (420nm) was 1.259. Clarity at (660 nm) was 35.9%. The major components were total sugars which recorded 12.80 %. Reducing sugars represents 11.20%. The juice contained 28.22 mg/100g fresh weight of L-ascorbic acid. Total antioxidant capacity (%), total flavonoids, total anthocyanin (mg/100g) and total phenols, were 53.72 %, 52.32 (mg/100g), 212.77 (mg/100g) and 434.61 (mg GAE /100g) respectively. Fourteen phenolic compounds were detected by HPLC. The obtained results during and after storage at room temperature 25C ° ± 5C ° for four months, declared that the clarified mulberry juice previously clarified gelatin ,having a lower stability than that clarified by Ultrafiltration process. Generally the juice was microbiologically safe. The sensory properties of mulberry nectar indicated that the prepared nectar from its concentrate after the end of storage is accepted and cope with the previous obtained data.

Key word: *Black mulberry juice, clarification, reconstitution, gelatin agent, ultrafiltration process.*

INTRODUCTION

The consumers interest for healthier and natural products has been growing and contributing to the consumption of lighter and refreshing products, such as fruit juices and bases fruit drinks (nectar, cocktails and drinks) (Matta *et al.*, 2004). From the Numerous literatures dealing to mulberry fruits, the various cultivars of mulberry fruits have nutritional and therapeutically properties, as they contain the phytochemical compound particularly the natural antioxidant agents. In the future, we should care for accelerating planting mulberry trees in places, which can not affect the agricultural area.

In Egypt, approximately 3 million old mulberry trees in the Delta area along the canals and roadsides were counted (Megalla, *et al* 1997). Mulberry, a member of the family *Moraceae*, is a deciduous tree distributed in the warm, moderate, and subtropical regions of Asia, Africa, North

America, and Southern Europe. There are more than 20 species of mulberry reported so far, and five of the species *Morus alba*, *M. nigra*, *M. rubra*, *M. indica*, and *M. laevigata* have edible fruit (Awasthi *et al.*, 2004).

Black mulberry fruits are consumed as fresh fruit or in the form of various confectionary products such as jam, marmalade, frozen desserts, pulp, juice, paste, ice cream, and wine. It is expected that the consumption of black mulberry fruit, with its delicious slightly acid flavor, extraordinary taste, and medicinal use will increase in the near future Koyuncu (2004). Black mulberry is a fruit known not only for its nutritional qualities and its flavor, but also for its traditional use in natural medicine, as it has a high content of active therapeutic compounds. (*Morus. nigra*) fruit is a good source of several phytonutrients and contains high amounts of total phenolics, total flavonoids, and ascorbic acid. Also, the

fruit has a pleasant taste with a slightly acidic flavor and an attractive dark red color Ozgen *et al* (2009).

Amin and Attia (2003) reported that the weight of black mulberry fruit was 1.89 g, fruit dimensions (length 1.93 cm and diameter 1.23 cm), size 1.66 cm³/fruit, juice yield was 76.54 %, pomace yield 23.46%, total soluble solids (T.S.S) 17.70% and the pH value was 5.70. Black mulberry has a moisture content of 79.68%, crude protein 9.50%, ash 5.22%, titratable acidity (as malice acid) 0.34%, reducing sugars 13.48%, non-reducing sugars 0.22%, total sugars 13.7, crude fiber 5.36%, nitrogen free extract 73.87% d.w. They also added that the black mulberry fruits are rich in anthocyanin (200.96 mg/100gm) on fresh weight basis.

Darias-Martín *et al* (2003) stated that the physical parameters of black mulberry were as follow, volume mass 1.06-1.007 g/cm, pH value was 3.10-3.36 and the TSS ranged between 13.00 to 17.50 %. They also found that the black mulberry has 3.12- 4.27 g/L of tannins, while the anthocyanin content was 2000 to 2600 mg/L.

Clarification process is very important technique for several fruit juices, to improve their qualities such as overall acceptability for the consumers and overcome the disadvantage of hazing and turbidity phenomena of some fruit juices especially during storage. (Stocke. 1998).

Ultrafiltration process (UF) has been used successfully in the clarification of pressed or refiltered of apple juice and other pressed fruit juices (Kirk *et al.*, 1963). Clarified juice coming from the Ultrafiltration process can be commercialized or submitted to a concentration process in order to obtain a product suitable for the preparation of juice and beverages, (Cassano *et al.*, 2003).

The concentration of fruit juices requires the part removal of water without changes in the composition of solids, leaving all the original solid components, such as fruit sugars, minerals, and vitamins, to the more concentrated solution. Concentration of fruit

juices, a major unit operation in the fruit-processing industry, is of critical importance as it determines the quality of the final product such as flavor, color, aroma, appearance and mouth feel. Color kinetics is the most important aspect of the successful processing of mulberry fruit, color is usually the first property the consumer observes. Fazaeli *et al* (2011).

Because of the black mulberry juice is considered an important a new and untraditional juice, which has a pleasant taste and aroma, but it has more of hazing precursors, so some of attempts of this work were conducted to reduce this phenomena, followed by the concentration process. The effect of storage at room temperature on the properties of reconstituted mulberry juice, as well as the sensory properties of mulberry nectar, were determined.

The aim of the current research was to study:

The physico-chemical constituents of black mulberry fruits. The optimum concentration of its effect on some properties of clarified juice. The influence of Ultrafiltration process and also gelatin agent on some estimated parameters of clarified juices. The effect of various techniques of concentration processes using (Rotary evaporator and basket centrifuge equipment's) on physico-chemicals constituents and sensory properties of reconstituted juices. The influence of storage conditions on physico-chemical constituents and microbial evaluation on reconstituted prickly pear juices as well as mulberry juice. The evaluation of sensory properties of the nectar prepared from concentrated prickly pear juices and mulberry juice after the end of storage period.

MATERIALS AND METHODS

Materials:-

Black mulberry fruits (*Morus nigra* L.) was obtained from Giza district Giza Governorate "Egypt". The all fruits were at ripe stage and collected in the year of 2012/2013.

Chemical and technological studies on black mulberry juice and its concentrate

Other materials used:

White sucrose sugar was purchased from the local market. Citric acid, sodium benzoate, potassium sorbate, Gelatin and celite were obtained from El- Gamhouria Trading Chemicals and Drugs Company, Egypt. The microbiological media (total bacterial count (TBC) using nutrient agar medium and yeast & mold count using potato dextrose agar (PDA), were obtained from El- Gamhouria Trading Chemicals and Drugs Company, Egypt.

Methods:

Technological methods

Extraction of black mulberry juice:

The ripe Black mulberry fruits were carefully sorted, graded, then proper washing, and extracted by pulper machine, which applied at (Food Technology Research Institute, Agriculture Research Center). The resultant pulp was homogenized, sieved with pressing well. The filtered juice was subjected to further processing.

Clarification process by gelatin agent:

Gelatin agent (0.15 g/L) was added to the homogenized juice (5000 rpm. for 5 min.). Sodium benzoate at the concentration of 0.025 % and also 0.025 % of potassium sorbate were added as preservative agents, and then left for 12 hours in a refrigerator. Celite as filter aid was used to facilitate the filtration operation by filter paper, using Buchner funnel. The clear juice was pasteurized at 95°C for 5 min. in a water bath and directly cooled with running tap water to the ambient temperature.

Ultra- filtration: (UF) Clarification:

Ultra-filtration was performed by using a laboratory pilot plant supplied by Tech-sep (Richelieu, (Québec) Canada) equipped with a Koch tubular membrane carboxer modules (type Series-Cor HFM-7 , PVDF, nominal molecular weight cut-off 50 (10³ Dalton), surface membrane area 0.16 m², No. tubes 7, UF membranes M8, membrane

diameter 6 mm, bursting pressure 40 bar, service pressure 15 bar, service pH 0-14). Experiments were carried out in the batch concentration mode to concentrate the juice up to a recovery factor of 42%. The membrane module was rinsed with tap water for 30 min after the treatment of the juice; then it was submitted to a cleaning procedure with the alkaline detergent Ultrasil 10 (El- Gamhouria Trading Chemicals and Drugs Company, Egypt) at a concentration of 0.2 % w/w % and at a temperature of 40 °C for 60 min. A final rinse of the system with tap water for at least 20 min was carried out. Body device was manufactured in the Synergy Egypt Company (Mohandeseen, Giza, Egypt) .Sodium benzoate at the concentration of 0.025 % and also 0.025 % of potassium sorbate were added as preservative agents, then the clear juice was pasteurized at 95°C for 5 min. in a water bath and directly cooled with running tap water to the ambient temperature.

Vacuum concentration:

Clarified mulberry juice was concentrated by rotary evaporator apparatus using an oil vacuum pump at 28 Hg and a temperature not exceeding 45°C. The process was continued until the concentration process reached 35 Brix of total soluble solids. Then the concentrated clear juice was packed into 250 ml. sterilized brown bottles, pasteurized at 90°C for 1min. in water bath , and cooled directly using tap water. The juices were subjected to the proper analysis.

Freezing concentration method: (Basket centrifuge):

Clarified mulberry juice were put inside the refrigerator cubes and frozen at -18°C for 24 hours. The frozen juice was placed in Basket centrifuge at 7000 rpm under atmospheric conditions to separate the concentrated juice from the formed ice, which remains inside the basket. This operation was repeated 3 times to obtain a concentrated juice of about 25 Brix.

The concentrated clear juice was packed into 250 ml. sterilized brown bottles, pasteurized at 90 °C for 1 min. in water bath, and cooled directly using tap water.

All porously previously juices were stored at ambient temperature (25C+5C) for four months and subjected to the proper analysis.

Reconstitution process

To analyze and evaluate the physico-chemical constituents as well as the sensory properties of the concentrates, the concentrates were reconstituted into the initial total soluble solids (16.4) as found in the fresh juice.

Preparation of black mulberry nectars:

To prepare black mulberry nectar, 40% from the single strength clarified reconstituted juice was used as participation ratio. 0.3% of citric acid was added to the formula. The final formula was adjusted to 16 Brix by adding sucrose solution. The nectar was packed into 200 ml. glass bottles and subjected to the pasteurization process at 90 C for 1 minute, directly cooled with running tap water and subjected to organoleptic evaluation.

Analytical methods:

Chemical constituents

Moisture, total soluble solids (T.S.S.) as Brix, total acidity, pH value, L ascorbic acid, crude fibers, ash and protein contents were determined according to A.O.A.C. (2012). Total sugars and reducing sugars were determined according to the method described by Somogy (1952) and Nelson (1974). Non-reducing sugars was calculated by difference

Physico-chemicals method

Viscosity was measured using Brookfield viscometer Model DV-III Rheometer, at 25°C using Spindle No. (21) LV and rotating speed of 50 rpm for one min. in both single strength turbid juice and in the case of clarified juice.

Browning index was measured according to the method reported by A.O.A.C. (2012), using (Shimadzu Spectrophotometer UV-120-02) at wave length of 420 nm. Color values were determined at (Food science

departs. Faculty of Agric., Cairo Univ., Giza, Egypt), using Hunter Lab model D-25 color and color difference meter according to the method of Cano *et al* (1995).

Pectin content was determined as described by Lees (1975).

Clarity was measured at 660 nm according to the methods of Krop and Pilnik (1974).

Antioxidant method

The total carotenoids were determined according to the method described by Askar and Treptow (1993). The antioxidant activities of the aqueous juice extracts was measured on the basis of the scavenging activity of the stable 1, 1-diphenyl-2-picrylhydrazyl (DPPH) free radical following the method described by Braca *et al.*, (2001). Total anthocyanin content of black mulberry juice was measured according the method described by Cevallos-casals and Cisneros-Zevallos, 2004)

Total flavonoids were measured using colorimetric assay developed by Zhishen *et al.*, (1999). The total phenolic compounds were determined with Folin-Ciocalteu reagent according to the method described by Danial and George (1979). The results were expressed as milligrams of gallic acid equivalent per gram of dry extract weight.

Phenolic compounds

Phenolic compounds were fractionated and identified by HPLC according to the method of Goupy *et al.* (1999).The fractionation was conducted at Food Tech. Res. Inst. A.R.C., Giza, Egypt.

Microbiological methods

Total viable bacterial count, yeast and mold were determined according to APHA (1992) .The counts were calculated per one ml of sample as colonies formed unit (CFU).

Sensory evaluation

Sensory properties (color, taste, aroma and overall acceptability) of black mulberry juice and the nectar prepared from it were evaluated by the staff at Food Tech. Res.

Chemical and technological studies on black mulberry juice and its concentrate

Inst., A.R.C. according to the method of Lindly *et al.* (1993).

Statistical analysis

The Statistical analysis of organoleptic evaluation was carried out using ANOVA under significant level of 0.05 for the analyzed data using SPSS (Ver. 11) .

RESULTS AND DISCUSSION

Physico-chemical parameter of fresh Black mulberry juice:

Based on the resulted data of Table (1), the juice yield of fresh juice was 62.80 %. The result was agreed with Amin and Attia (2003). Ercisli and Orhan (2008) who found that the fruit pulp were 64.03% and 68.58% of black mulberry genotypes.

Moisture content represented about 81.20 %. In this trend, Abo zaid (2014),

found that the moisture content in fresh black mulberry juice was 80.18%.

Also, total soluble solids, pH value and tetratable acidity, (as malice acid), were 16.4%, 5.49 and 0.341% as malic acid, respectively, Table (1). These results are in accordance with Amin and Attia (2003).

Browning index measured by spectrophotometer at (420 nm) recorded 1.259, while the color value measured by Hunter lab recorded (L 27, a 8 and b 18), and the related parameter of juice clarity measured at (660 nm) was 35.9% respectively, Aramwit *et al* (2010) reported that the Color parameters of fresh black mulberry juice were (L 11.94 to 27.75, a 13.9 to 20.62 and the b value was 2.12 to 9.28) respectively. Also in Table (1), the viscosity of juice recorded 89.1 cp. The results are in accordance with obtained by Ercisli and Orhan (2007) and Editor *et al* (2013).

Table (1) Physico-chemical attributes of fresh Black mulberry juice.

Physico-chemical attributes	
Yield(%)	62.8
Moister (%)	81.20
Total soluble solids (%)	16.4
pH value	5.49
Color index at (420nm)	1.259
*Viscosity(cp)	89.1
Clarity at (660 nm) %	35.9
*Colour parameters L	27
a	8
b	18
Total sugars (%)	12.80
Reducing sugars (%)	11.20
Non-reducing sugars (%)	1.60
Protein (%)	1.62
Fibers (%)	2.20
L-Ascorbic acid (mg/100g)	28.22
Titrateable acidity(% as malice cid)	0.341
Total antioxidant capacity%	53.72
Total flavonoids (mg/100g)	52.32
Total anthocyanin (mg/100g)	212.77
**Total phenol (mg GAE /100g)	434.61

*Colour parameters L- (lightness darkness) a- (yellowness bluness) b- (redness greeness)

**GAU- gallic acid equivalents

Concerning total sugars, reducing sugars and non-reducing sugars of fresh black mulberry juice, the analytical data represented 12.80 %, 11.20% and 1.6 % respectively, table (1). Neva and Memnune (2008) reported that the main sugars in fresh black mulberry juice were reducing sugars, which ranged between (9.42-15.46%) and non-reducing sugars ranged from 1.57- to 4.36% on fresh weight basis respectively. The determined protein and fiber of fresh black mulberry juice represented 1.62 % and 2.20 % respectively. These results are in accordance with obtained by Singhal *et al* (2010).

The percent of L-ascorbic acid of the fresh black mulberry juice was 28.22 mg/100g. This agrees with Amin and Attia (2003), Ercisli and Orhan (2007) and Yildiz (2013).

As natural phytochemical components, for instance, total antioxidant capacity (%),

total flavonoids, total anthocyanin (mg/100g) and total phenols, found of fresh black mulberry juice represented 53.72 %, 52.32 (mg/100g), 212.77 (mg/100g) and 434.61 (mg GAE /100g) respectively. These results are in agreement with Shih *et al* (2010) and Contessa *et al* (2013).

Identification of phenolic compounds using HPLC:

From Table (2) and Fig. (1), it could be noticed that fourteen phenolic compounds among the seventeen authentic samples of standard were detected on mulberry juice. The detected compounds were, 3-hydroxytyrosol, oleuropein, catechol, benzoic, chlorogenic acid, salicylic acid, catechin, ellagic, caffeine, ferulic, caffeic, gallic, vanilic and cinnamic acid, which counted (9.414 mg/100g, 3.513, 2.878, 2.789, 23.75, 1.729, 18.20, 1.605, 4.41, 4.30, 4.14, 2.68, 2.37 and 1.49 mg/100g respectively).

Table (2): Fractionations of phenolic compounds using HPLC method of fresh black mulberry juice.

Phenolic Compounds	(%)
Cinnamic acid	1.494
Ellagic	1.6056
Ferulic	4.3006
Caffeic	4.4133
Catechin	18.200
Caffeic	4.1480
Salicylic	1.72910
Chlorogenic	23.756
Gallic acid	2.688
Catechol	2.87810
3-hydroxy-tyrosol	9.414
Oleuropein	3.5130
Vanilic	2.376
Benzoic	2.789




Fig (1): HPLC chromatogram of phenolic compounds of fresh mulberry juice.

Clarification processes of fresh juice:

Clarified mulberry juice as affected by using gelatin agent and UF precise

The main determined parameters used to know the efficiency of clarification process, clarity and viscosity. The percent of clarity increased from 35.9% of turbid juice to 75.5% in clarified juice, as a result of using gelatin agent, (Table 3). On the contrary, Table (3) also shows the decreasing of viscosity from 70 cp of turbid juice to 55 cp in clarified mulberry juice. These results were in agreement with those reported by (Siebert and Lynn., 1997) and (Kwang-Sup *et al.*, 2004).

To explore the efficiency of UF process on resultant mulberry juice, TSS%, pH value, clarity and viscosity parameters were evaluated, Table (3). They were 16.1, 5.40, 97.5 and 10 cp in UF implemented juice comparing with turbid juice, which was 16.4, 5.49, 35.9 and 89.1 cp respectively, table (3). The results were agreement with (De-Barrod and peres. 2003), Rai *et al* (2005) and (Alvarez *et al.*, 1998).

Effect of concentration process on some physico-chemical attributes of reconstituted clarified mulberry juice

As implemented in the production line of some factories, after clarification processes, concentration process follows it. Also for analyzing the physico-chemical constituents as well as the sensory properties of the concentrates, it is a must to reconstitute the concentrates into the initial total soluble solids as found in the fresh juice, without treatment.

Dealing with the percent of clarity determined of clarified mulberry juice (by UF and gelatin), concentrated by rotary evaporator, it reached at zero time 69.8% compared with the fresh juice which was 35.9% and reached 93.7% in the case of UF applied juice. When freezing concentration used, slightly increasing was also observed of UF and gelatin implemented mulberry

juice, (Table 4). The results of freezing concentration met with obtained by Askar *et al.*, (1981).

Color index of fresh mulberry juice was also affected by concentration type used as well as the clarification type. In the case of freezing concentration, remarkable decrease of color index was observed than that concentrated by rotary evaporator. Indicating that the UF applied juice surpassed the gelatin treated juice of both rotary evaporator and freezing concentration method, (Table 4).

Small reduction of total sugars was observed during concentration conducted by freezing concentration followed by rotary evaporator and also when UF clarification and gelatin agent used. Dealing with reducing sugars of all application, the same trend was observed, Table (4).

Incubation time, temperature, chemical agent, nature of UF membrane, Concentration method and, pasteurization affect ascorbic acid stability. As shown in Table (4), when concentration applied by rotary evaporator, ascorbic acid decreased from 28.22 mg/100g of fresh turbid juice to 10.33 mg/100g of gelatin treated and 11.11 mg/100g of UF treated juice. While when concentration applied by freezing, ascorbic acid decreased from 28.22 mg/100g of fresh turbid juice to 15.18 mg/100g of gelatin treated and 17.22 mg/100g of UF treated juice, table (4). The results were in accordance with those reported by, Butera *et al* (2002), and Cassano *et al* (2003).

Total flavonoids (mg/100g), total anthocyanin (mg/100g), total phenol mg GAE/g and total antioxidant capacity of mulberry juices were listed in table (4). Generally there was observing decreasing of previous components as affected by clarification and concentration method used. The UF applied juice surpassed gelatin treated, and also freezing concentration surpassed the concentration by rotary evaporator. Based on the previous data discussed, freezing concentration was chosen, and used during storage period.

Chemical and technological studies on black mulberry juice and its concentrate

Table (3): Effect of clarification by gelatin agent and UF on some parameters of clarified black mulberry juice

Parameter	fresh	Gelatin	(UF)
Total soluble solids (%)	16.4	16.6	16.1
PH value	5.49	5.45	5.40
Clarity at (660 nm)	35.9	75.5	97.5
*Viscosity(cp)	89.1	55	10

*Cp- centipoise

Table (4): Some physio-chemical attributes of clarified black mulberry juice as affected by concentration using rotary evaporator

Juice properties	fresh	Black mulberry concentration process by rotary and freezing concentration			
		rotary		freezing concentration	
		gelatin	UF treated	gelatin	UF treated
Total soluble solids (%)	16.4	16.4	16.4	16.4	16.4
PH value	5.49	4.64	4.60	4.84	4.80
Colour index at (420nm)	1.259	1.440	1.411	1.380	1.355
Clarity at (660 nm) %	35.9	69.8	93.7	74.30	96.90
Total sugars (%)	12.80	11.97	11.75	12.11	12.55
Reducing sugars (%)	11.20	10.33	10.19	10.44	11.31
Non-reducing sugars (%)	1.60	1.64	1.56	1.67	1.24
Titrateable acidity % (as citric acid)	0.341	0.444	0.431	0.450	0.443
L-Ascorbic acid (mg/100g)	28.22	10.29	11.11	15.18	17.22
Total antioxidant capacity %	53.72	42.31	45.39	48.11	49.25
Total flavonoids (mg/g)	52.32	41.79	40.92	47.23	48.17
Total anthocyanin (ug/100g)	212.77	135.19	148.20	162.29	168.88
*Total phenol (mg GAE /g 100 g)	434.61	395.1	409.3	411.15	422.22

* GAU- gallic acid equivalents

Sensory properties of reconstituted juice at zero time:

Sensory properties of all reconstituted clarified juices at zero time were illustrated of Table (5). From the analyzed data, it can

be concluded that the reconstituted clarified juices at zero time were accepted in their sensory attribute such as color, taste, aroma and overall acceptability, which ranged between 7.7 and 9.8 degrees.

Table (5): Organoleptic evaluation of reconstituted mulberry juice.

Parameter	Gelatin concentration basket centrifuge	Ultrafiltration concentration basket centrifuge	Gelatin concentration rotary	Ultrafiltration concentration rotary
Color	8.6 ^c	9.5 ^a	8.1 ^{de}	9.1 ^{ab}
Taste	9.6 ^a	8.8 ^{bc}	8.6 ^c	7.7 ^d
Aroma	9.8 ^a	8.8 ^{bc}	9.1 ^b	7.9 ^c
Overall-acceptability	9.6 ^a	9.4 ^{ab}	8.5 ^b	7.8 ^c

Means in the same row with different letters are significantly different (pp≤0.05)

Effect of storage on Physico-chemical attributes of clarified reconstituted mulberry juice.

The effect of storage conditions on color index at 420 nm of reconstituted clarified mulberry juice by gelatin agent was observed of Table (6), and the data revealed that the color index increased from 1.38 at zero time to 1.52 at the end of storage period. While in the case of UF, the color index reached 1.512 at the end of storage period, Table (7).

Concerning, clarity at 660 nm determined, which shows of table (6 and 7), gradually decreasing during storage period was observed of both gelatin treated and also UF treated juice.

Also, there were some what changed of acidity, which recorded little increasing during storage period. On contrary pH value was decreased gradually during storage period, of gelatin treated and also UF treated juice table, (6 and 7).

Dealing with total sugars and reducing sugars of reconstituted black mulberry juice, there was little decreasing during storage period of gelatin treated and also UF treated juice Table, (6 and 7).

The effect of storage conditions on color index at 420 nm of reconstituted clarified mulberry juice by gelatin agent and UF was

illustrated in table (6 and 7), and the results show that the reconstituted juice at zero time contained 15.18 mg/100g and 17.22 mg/100g, whereas, they were 6.52 mg/100g and 8.36 mg/100g of gelatin agent and UF treated juice. The results were in agreement with (Davey *et al.*, 2000 and Franke *et al.*, 2004). Total antioxidant capacity as mg/100g , total flavonoids mg/100g, total anthocyanin as µg/100g and the total phenol GAE/g as natural antioxidant agents of reconstituted black mulberry juice were determined and the data revealed that, there were gradually decreasing of them during storage period at room temperature. This is agree with Polydera *et al* (2004), and Kalt *et al* (1999).

Microbiological aspects of black mulberry juice:

Some of basic microbial analysis of fresh, zero time and after 4 months of storage period such as(total bacterial count and yeast & mold counts) of all mulberry juice samples were analyzed, (Table 8) .

The percent of total counts resulted from analyzed data of gelatin and UF treated juice ranged from 1.0×10² cfu/g to 3.2×10² cfu/g. whereas the yeast & mold counts ranged from 2.8×10² cfu/g to 4.6×10² cfu/g. This means the reconstituted mulberry juice at zero time and after 4 months of storage period are safe.

Chemical and technological studies on black mulberry juice and its concentrate

This could be attributed mainly to the very low pH observed (3.94- 4.04). In addition, conditions under which the juice was processed, stored, and/or served might have contributed to the betterment of the product. In fact, its low pH did not inhibit the growth of acid tolerant yeasts and molds, and this allowed their proliferation to level as high as 3.5×10^4 and 4.0×10^4 log CFU/ml, respectively Olorunjuwon *et al.*, (2014).

Organoleptic characteristics of mulberry nectar:

Mulberry nectar prepared from its concentrate was sensory evaluated. The statistical analyzed sensory data of Table (9), declared that the whole parameters were accepted and ranged from 7.5 to 8.8 degree. The sensory properties of mulberry nectar indicated that the prepared nectar from its concentrate after the end of storage is accepted and met with the previous evaluated analysis.

Table (6): Effect of storage of concentrated clarified (by gelatin agent) mulberry juice, on physico-chemical attributes.

Parameter	Black mulberry					
	fresh	Zero time	1 month	2 month	3 month	4 month
pH value	5.49	4.84	4.78	4.71	4.66	4.58
Color index at (420nm)	1.259	1.380	1.410	1.460	1.511	1.522
Clarity at (660 nm)	35.9	74.30	74.0	73.6	69.1	60.36
Total sugars (%)	12.80	12.11	12.98	12.78	12.66	12.61
Reducing sugars (%)	11.20	10.44	11.17	11.02	10.87	10.79
Non-reducing sugars (%)	1.6	1.67	1.28	1.76	1.79	1.85
Titrateable acidity% (as citric acid)	0.341	0.450	0.462	0.477	0.484	0.492
L-Ascorbic acid (mg/100g)	28.22	15.18	13.22	10.39	8.96	6.52
Total antioxidant capacity %	53.73	48.11	46.33	44.21	38.36	35.49
Total flavonoids (mg/g)	52.32	47.23	45.66	42.39	38.11	35.33
Total anthocyanin (ug/100g)	212.77	162.29	148.39	132.11	121.21	118.85
*Total phenol (mg GAE /g)	434.61	411.15	398.33	381.02	364.65	357.25

*GAU- gallic acid equivalents

Table (7): Effect of storage of concentrated mulberry juice, clarified by UF on physico-chemical constituents.

Parameter	Black mulberry					
	fresh	Zero time	1 month	2 month	3 month	4 month
pH value	5.49	4.80	4.72	4.64	4.58	4.44
Colour index at (420nm)	1.259	1.355	1.411	1.489	1.496	1.512
Clarity at (660 nm)	35.9	96.9	94.36	91.96	89.63	87.98
Total sugars (%)	12.80	12.55	12.10	11.88	11.80	11.71
Reducing sugars (%)	11.20	11.31	11.14	11.01	10.96	10.84
Non-reducing sugars (%)	1.6	1.24	0.96	0.78	0.84	0.86
Titrateable acidity % (as citric acid)	0.341	0.443	0.471	0.510	0.533	0.541
L-Ascorbic acid (mg/100g)	28.22	17.22	14.20	12.11	10.91	8.36
Total antioxidant capacity %	53.73	49.25	47.33	45.10	41.32	39.39
Total flavonoids (mg/g)	52.32	48.17	47.36	45.30	41.44	40.05
Total anthocyanin (ug/100g)	212.77	168.88	144.36	139.38	131.27	128.28
*Total phenol (mg GAE /g)	434.61	422.22	414.33	401.12	384.92	371.25

Table (8): Microbiological counts of reconstituted clarified(by gelatin) black mulberry juice as affected by storage conditions.

UF treated Black mulberry juice storage period	Counts *(CFU)			
	gelatin		UF	
	Total count	Yeast & mold	Total count	Yeast & mold
Fresh	3.2×10 ²	4.6×10 ²	3.2×10 ²	4.6×10 ²
Zero time	1.3×10 ²	2.8×10 ²	1.1×10 ²	3.1×10 ²
After 4 month	1.1×10 ²	3.0×10 ²	1.0×10 ²	3.3×10 ²

Chemical and technological studies on black mulberry juice and its concentrate

Table (9): Sensory properties of mulberry nectar.

Parameter	Gelatin concentrated by basket centrifuge	UF concentrated by basket centrifuge
Color	8.3 ^b	8.7 ^a
Taste	8.5 ^a	8.1 ^b
Aroma	8.7 ^a	8.2 ^b
Overall- Acceptability	8.4 ^a	8.0 ^b

REFERENCE

- Abou-Zaid, M.S.M.A. (2014). Technological and biochemical studies on processing some untraditional vegetable and fruit varieties. Ph.D thesis (Food Science and technology) Fac. of Agric. Cairo. Al-Azhar University.
- Alvarez, S., F. A. Riera, R. A´lvarez and J. Coca (1998). Permeation of apple aroma compounds in reverse osmosis. Separation and Purification Technology, 14, 209–220.
- Amin, W.A. and R. S. Attia (2003). Characterization of the Egyptian Mulberry (TUT) Fruit for preparing processed products and Natural colors. J. Agric. Sci. Mansoura Univ., 28(3): 1895-1909.
- APHA (1992). Standard methods for the examination of water and wastewater, 18th edition. American Public Health Association, Washington, D. C.
- Aramwit, P., N. Bang and T. Srichana (2010). The properties and stability of anthocyanins in mulberry fruits. Food Res. Int... 43:1093-1097.
- Askar, A. and H. Treptow (1993). Amines. In Encyclopedia of food science, food technology and nutrition. New York: Academic Press.
- A.O.A.C. (2012). Official Methods of Analysis. Association of Official Analytical Chemists. Published by the AOAC. International 19th ed., MARYLAND, USA.
- Askar, A.; El-samahy, S.K.; Abd-El-Baki, M.M; Ibrahim, S.S. and Abd-El-Fadeel, M.G. (1981): Production of lime juice concentrates using the serum-pulp method. Alimenta. 20(5): 121-128.
- Awasthi, A.K., G.M. Nagaraja, G.V. Naik, S. Kanginakudru, K. Thangavelu and J. Nagaraja (2004). Genetic diversity in mulberry (Genus *morus*) as revealed by RAPD and ISSR marker assays, BMC Genetics. 5: 1.
- Braca, A., N.D. Tommasi, L.D. Bari, C. Pizza, M. Politi and I. Morelli (2001). Antioxidant principles from *Bauhinia terapotensis*. J. Nat. Prod. 64: 892-895.
- Butera, D., L. Tesoriere, F. Di Gaudio, A. Bongiorno, M. Allegra, A.M. Pintaudi, R. Kohen and M.A. Livrea (2002). Antioxidant activities of Sicilian prickly pear (*Opuntia ficus-indica*) fruit extracts and reducing properties of its betalains: betanin and indicaxanthin. J. Agricultural and Food Chemistry 50 (23): 6895–6901.
- Cano, E. and L.C. Mahadevan (1995). Parallel signal processing among MAPKs Trends Biochem. Sci., 20 (1995), pp. 117–122.
- Cassano, A., E. Drioli, G. Galaverna, R. Marchelli, G. Di Silvestro and P. Cagnasso (2003). Clarification and concentration of citrus and carrot juices by integrated membrane processes. Journal of Food Engineering, 57(2): 153–163.
- Cevallos-Casals, B.A. and L. Cisneros-Zevallos (2004). Stability of anthocyanin-based aqueous extracts of Andean purple corn and red-fleshed sweet potato compared to synthetic and natural colorants. Food Chemistry 86: 69–77.

- Contessa, A.K., G. M. Nagaraja, G. V. Naik, S. Kanginakudru, S. K. Thangavelu and J. Nagaraju (2013). Genetic diversity and relationships in mulberry (genus *Morus*) as revealed by RAPD and ISSR marker assays. *BMC Genetics*, 5: 1–9.
- Daniel, H.D. and C.M. George (1979). Peach seed dormancy in relation to endogenous inhibitor and applied growth Substances. *J. Amer. Soc. Hort. Sci.*; 97: 651-655.
- Darias-Martin, J., G. Lobo-Rodrigo, J. Hernandez-Cordero, E. Diaz-Diaz and C. Diaz-Romero (2003). Alcoholic beverages obtained from black mulberry. *Food Technol Biotechnol.* 41(2):173–176.
- Davey, M.W., M. Van-Montagu, D. Inze, M. Sanmartin, A. Kanellis and N. Smirnoff (2000). Plant L-ascorbic acid: Chemistry, function, metabolism, bioavailability and effects of processing. *J. Sci. Food Agric.* 80: 825-860.
- De-Barros, A.E.V., G.P. Manfio, V. Ribeiro-Maitan, L.A. Mendes-Bataus, S. B. Kim, L.A. Maldonado and M. Goodfellow (2003). *Nocardia cerradoensis* sp. nov., a novel isolate from Cerrado soil in Brazil. *Int J Syst Evol Microbiol* 53, 29–33.
- Editor, N.T.D., T.H. Le, P.M. Nguyen and T.A.D. Dong (2013). Application of natural fermentation to ferment mulberry juice into alcoholic beverage. *Journal of Advances in Biology.* 1(1): 244-247.
- Ercisli, S. and E. Orhan (2007). Chemical composition of white (*Morus alba*), red (*Morus rubra*) and black (*Morus nigra*) mulberry fruits. *Food Chemistry*, 103: 1380–1384.
- Ercisli, S. and E. Orhan (2008). Some physico-chemical characteristics of black mulberry (*Morus nigra* L.) genotypes from Northeast Anatolia region of Turkey. *Sci. Hortic.*, 116: 41–46.
- Fazaeli, M., Z. Emam-Djomeh, A. Kalbasi Ashtari and M. Omid (2011). Effect of spray drying conditions and feed composition on the physical properties of black mulberry juice powder. *Food and Bioproducts Processing*, 90 (4): 667-675.
- Franke, A.A., L.J. Custer, C. Arakaki and S.P. Murphy (2004). Vitamin C and flavonoid levels of fruits P.S. and vegetables consumed in Hawaii. *J. Food. Compos. Anal.* 17: 1-35.
- Goupy, P., M. Hugues, P. Boivin and M.J. Amiot (1999). Antioxidant compounds of barley (*Hordeum vulgare*) and malt extracts. *J. Science of Food and Agriculture*, 79: 1625–1634.
- Kalt, W., C.F. Forney, A. Martin and R.L. Prior (1999). Antioxidant Capacity, Vitamin C, Phenolics, and Anthocyanins after Fresh Storage of Small Fruits. *J. Agric. Food Chem.* 1999, 47, 4638–4644.
- Kirk, D.E., M.W. Moutgomery and Kortekaas (1983). Clarification of pear juice by hollow filter ultrafiltration. *J. Food Sci.* 48: 1663.
- Koyuncu, F. (2004). Morphological and agronomical characterization of native black mulberry (*Morus nigra* L.) in Su'ç'u'ler, Turkey. *IPGRI News Lett.* 138: 32–35.
- Krop, J.J.P. and W. Pilnik (1974). Effect of pectic acid and bivalent cation on cloud loss of citrus juice. *Lebensm Wiss Technol.*, 7: 62-63.
- Kwang-Sup, Y., J.H. Hong, D.H. Bae, S.J. Kim and S.D. Kim (2004). "Effective clarifying process of reconstituted apple juice using membrane filtration with filter-aid pretreatment," *J. Membrane Sci.*, 228: 179-186 .
- Lee, C.Y., T.E. Acree and R.M. Butts (1975). Determination of methyl alcohol in wine by gas chromatography. *Anal. Chem.* 47: 747–748.
- Lindley, R., A. Williams and H.G. Hunt (1993). Anomalous seasonal cycles of decapod crustacean larvae in the North Sea in an exceptionally warm year. *J. Exp. Mar. Biol. Ecol.* 172: 47–66.
- Matta, N.E., N. Bastos, R. Gutiérrez, O. A. Rodríguez and E. C. Greiner (2004). Prevalence of blood para sites in Tyrannidae (flycatchers) in the eastern plains of Colombia. *Mem. Inst. Oswaldo Cruz.* 99: 271-274.
- Megalla, S.E., N.E.M. El-Keltawi and S.A. Ross (1997). A study of antimicrobial action of some essential oil constituents. *Herba Polonica*, 26: 180-186.

Chemical and technological studies on black mulberry juice and its concentrate

- Nelson, N. (1974). A photometric adaptation of the Somogyi methods for the determination of glucose. *J. Biol. Chem.*, 153 : 375-380.
- Neva, G. and S. Memnune (2008). Antioxidant Activity, total phenolic content and selected physicochemical properties of white mulberry (*Morus Alba L.*) fruits. *Int. J. of Food Properties*, 11 :(1): 44 – 52.
- Olorunjuwon, B.O., B.K. Temitope, F.O. Muibat and O. Afolabi (2014). Microbiological quality of some locally produced fruit juices in Ogun state, South Western Nigeria. *J. Microbiology Research*, 2(1): 1-8.
- Ough, C.S. and E.A. Crowell (1979). Pectic-enzyme treatment of white grapes: Temperature, variety and skin-contact time factors. *Am. J. Enol. Viticult.*, 30: 22-27.
- Ozgen, M., S. Sedat and K. Cemal (2009). Phytochemical and antioxidant properties of anthocyanin-rich *Morus nigra* and *Morus rubra* fruits. *Scientia Horticulturae*. 119(3): 275-279.
- Polydera, A.C., N.G. Stoforos and P.S. Taoukis (2004). The effect of storage on the antioxidant activity of reconstituted orange juice which had been pasteurized by high pressure or heat. *International J. Food Science & Technology*. 39(7): 783–791.
- Rai, P., G.C. Majumdar, S. Das-Gupta and S. De (2004). Prediction of the viscosity of clarified fruit juice using artificial neural network: a combined effect of concentration and temperature. *Journal of Food Engineering* 68 (2005) 527–533.
- Shih, P., Y. Chan, J. Liao, M. Wang and G. Yen (2010). Antioxidant and cognitive promotion effects of anthocyanin-rich mulberry (*Morus atropurpurea L.*) on senescence-accelerated mice and prevention of Alzheimer's disease. *J. Nutritional Biochemistry* 21: 598–605.
- Siebert, K.J. and P.Y. Lynn (1997). Haze-Active Protein and Polyphenols in Apple Juice Assessed by Turbidimetry. *J. Food Science* 62(1): 79–84.
- Singhal, B.K., M.A. Khan, A. Dhar, F.M. Baqual and B.B. Bindroo (2010). Approaches to industrial exploitation of mulberry (*mulberry sp.*) fruits, *J. Fruit and Ornamental Plant Research* 18(1): 83-99.
- Somogyi, M. (1952). Notes on sugar determination. *J. Biol. Chem.*, 195: 19-23.
- Stocké, R. (1998). “The 3-component stabilization with bentonite, gelatin and silica sol,” *Fruit Processing* 1: 6–10 .
- Yildiz, O. (2013). Physicochemical and sensory properties of mulberry products: Gümüşhane pestil and köme. *Turk. J. Agric. For.*, 37: 762-771.
- Zhishen, J., T. Mengcheng and W. Jianming (1999). The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals, *Food Chem*, 64: 555-559.

دراسات كيميائية و تكنولوجيا على عصير و مركز التوت

أبو الفتح عبد القادر البديوي^(١) ، السيد حلمي رحمه^(١) ، السيد شريف عبدالوهاب^(٢) ،
نجلاء متولى على أحمد^(٢)

^(١) قسم علوم الاغذية- كلية الزراعة-جامعة المنوفية-شبين الكوم- مصر
^(٢) معهد بحوث تكنولوجيا الاغذية- مركز البحوث الزراعية- الجيزة - مصر

الملخص العربي

تعتبر عصائر الخضر والفاكهه سواء كانت فى صوره رائقه او غير رائقه والمنتجات التى تصنع من خلالها مثل المركزات و الشراب المركز والنكتار والمشروبات المتعدده التركيز ذات قيمة غذائيه و صحية وإقتصاديه بالغه الاهمية .
فيما يخص التركيب الطبيعى والكيمائى لعصير التوت الاسود، فمن خلال النتائج وجد انه يحتوى على نسبة تصافى وصلت الى ٦٢.٨% ومحتوى من الرطوبة وصل الى ٨١.٢٠%. كما سجلت قيم المواد الصلبة الكلية الذائبة و القم الهيدروجينى و الحموضة الكلية (١٦.٤% و ٥.٤٩ و ٣.٤١% على التوالى. اما فيما يخص التلون البنى فقد كان ١.٢٥٩ نانوميتر. أما الشفافية فقد كانت ٣٥.٩%. ومن خلال النتائج ايضا فقد سجلت السكريات الكلية اعلى القيم بالنسبة لباقي المركبات ، حيث كانت ١٢.٨% فى حين وصلت السكريات المختزلة إلى ١١.٢٠%. كما احتوى العصير الطازج على ٢٨.٢٢ ملجم/١٠٠ جم من حمض الاسكوربيك و ٥٣.٧٢ ملجم/١٠٠ جم من مضادات الاكسدة الكلية. كما احتوى العصير على ٥٢.٣٢ ملجم/١٠٠ جم من الفلافونويدات و ٢١٢.٧٧ ملجم/١٠٠ جم من الانثوثيانين ، علاوة على إحتوائه نسبة مرتفعة من الفينولات الكلية ، حيث وصلت إلى ٤٣٤.٦١ ملجم من حامض الجاليك/ جم من المستخلص الجاف.
اما فيما يخص المركبات الفينولية المقدره بالفصل الكروماتوجرافى ، تم الحصول على ١٤ مركبا فينوليا. كما أكدت النتائج ان الترويق بالترشيح الدقيق قد تميز عن الترشيح باستخدام الجيلاتين.
و فيما يتعلق بنتائج تأثير عمليه التركيز على الخواص الطبيعيه والكيمائيه، أكدت ان التركيز بالتجميد اعطى خواص جيدة للعصير الناتج، كما اكد ذلك اختبار التقييم الحسى عقب التركيز مباشره. ومن خلال النتائج اثناء تخزين العصير المركز بالتجميد على درجه حراره الغرفه ٢٥+٥ م ، وجد ان مكونات العصير الرائق بالترشيح الدقيق كانت اكثر تميزا وخاصة فى خصائص الترويق عنه فى العصير الناتج بالترشيح بالجيلاتين، علاوة على ان العصير عامة أمن ميكروبيا.
وفى النهايه تم اجراء تقييم حسى للنكتار المصنع من العصير المركز بعد انتهاء فتره التخزين وهى اربع شهور حيث اكدت النتائج ان النكتار الناتج فى كلتا الحالتين كان مقبولا لدى المحكمين وتتماشى نتائجه مع النتائج السابقه للتحليل الطبيعى والكيمائى.

