

## INTERACTION BETWEEN HUMIC ACIDS AND NEUTRAL SALTS ON PLANT GROWTH

Manal F. Tantawy<sup>(1)</sup>, R. A. Khalil<sup>(2)</sup>, E. A. Abou Hussien<sup>(2)</sup> and Asmaa M. M. Soliman<sup>(2)</sup>

<sup>(1)</sup> Soil, Water and Environment Int Res Arc Giza

<sup>(2)</sup> Soil Sci. Dept. Fac. Of Agric. Minufia Univ.

(Received: Sep. 23, 2014)

**ABSTRACT:** *This study was carried out evaluate the combined interaction between two humic acids varied in their sources and chemical compositions and three neutral salts. i.e. NaCl, CaCl<sub>2</sub> and FeCl<sub>2</sub> on barley variety Giza 123 (*Hordium vulgare* L. ) growth and its content of Na, Ca and Fe elements. This investigation was conducted on pots experiment in a completely block design with three replicates using sandy culture. The used humic acids were extracted from alluvial soil (HAS) and compost of clover straw (HAC) and added to sandy culture at application rates of 0, 10, 20, 40 and 100 mgkg<sup>-1</sup> sand. The application rates of neutral salts were 0, 250, 500 and 1000 mgkg<sup>-1</sup> sand. After 42 days of planting, the plants of each pot were harvesting. The dry matter yield of either of shoots or roots of the harvested plants were weighted and statically analyzed for LSD value at 0.05.*

*The dry weights of both shoots and roots of barley plants increased significantly with increasing added humic acids. The found increases of dry weights in the plants treated with HAC were higher relatively than those associated the treatments of HAS. With different treatments of humic acids, the found dry weights of shoots were higher than those of roots. The response of barley dry weight for the tested treatments of neutral salts were varied widely according to the used neutral salts and its application rates. Agronomical efficiency of humic acids was decreased with the increasing rate of added NaCl, but it increased with added CaCl<sub>2</sub> up to 500 mg/kg and also with the increase of added FeCl<sub>2</sub> up to 1000 mg/kg. Shoots and roots of barley plants content of Na, Ca or Fe increased with the increasing rates of added NaCl, CaCl<sub>2</sub> or FeCl<sub>2</sub>. Application of humic acids played a major role in the decrease of harmful effects of salinity and its effect on both plant growth and elements uptake.*

**Key words:** *Barley, Humic acids, Neutral salts, Agronomical efficiency, Chemical composition and Elements uptake.*

### INTRODUCTION

Humic acids are a commercial product contains many elements which improve the soil fertility and increasing the phyto-availability of nutrient elements and consequently affected plant growth and yield. Humic acid particularly is used to remove or decrease the negative effects of mineral fertilizers and some chemicals forms in the soil. Humic substances have many beneficial effects on soil and consequently on plant growth and are shown highly hormonal activity. These materials not only increase macronutrients contents and ions uptake but also enhance micronutrients of the plant organs (Brunetti *et al.*, 2005 ).

In other study, Liu (1998) found that the application of humic acids during salinity stress did not increase the uptake of N, P, K or Ca. Also, in their study; foliar application with 0.1% humic acid treatment increased the dry weight, N, P, K, Ca, Mg, Na, Fe, Zn, and Mn amounts in plants with 60mM NaCl treatment when compared with the control and 0.2% humic acid treatment.

El-Gundy (2005) ; Emam (2011) Nada and Tantawy (2013) showed that, increasing added HA and salinity level of irrigation water resulted in an increase of soil content of available Ca. Also Aydin *et al.* (2012) showed that shoot growth was more inhibited by NaCl than root growth. Humic acid (HA) application to the soil was

ameliorated to the adverse effects of salinity on the shoot and root dry matter. The highest salt doses (120 mM) of NaCl, CaCl<sub>2</sub>, MgCl<sub>2</sub> and KCl<sub>2</sub> without HA applications caused plant death, but no plant death was obtained when applied HA (0.05 and 0.1%) doses of all the salt types and doses with exception for CaCl<sub>2</sub>. Soil salinity is characterized by high amounts of Na<sup>+</sup>, Mg<sup>+2</sup>, Ca<sup>+2</sup>, Cl<sup>-</sup>, HCO<sub>3</sub><sup>-</sup>, SO<sub>4</sub><sup>-2</sup> and B ions which have negative effects on the plant growth. Generally, NaCl causes salt stress in nature. Aydin *et al.* (2012) found that salinity negatively affected the growth of corn; it also decreased the dry weight and the uptake of nutrient elements except for Na and Mn. Humus application of soil increased N uptake by corn while foliar application of humic acids increased the uptake of P, K, Mg, Na, Cu and Zn. Although the effect of interaction between salt and soil humus application was found statistically significant. The interaction effect between salt and foliar humic acids treatments were not found significant. Under salt stress, the first doses of both soil and foliar application of humic substances increased the uptake of nutrients. Atiyeh *et al.* (2002) found that, the root to shoot ratios of tomato seedlings increased significantly with increasing concentrations of humic acids in the soils container medium, indicating greater resource allocation towards the roots than the shoots.

This study was carried out to:- 1- Study the effect of some neutral salts i.e., NaCl, CaCl<sub>2</sub> and FeCl<sub>2</sub> applied at different rates on plant growth and its chemical composition, .2- Study the effect of humic acids different in their chemical composition on plant growth and its chemical composition and .3- Study the interaction of both neutral salts and humic acids on plant growth and its chemical composition.

## **MATERIALS AND METHODS**

This study was conducted on Soil Science Department, Faculty of Agriculture, Minufia University to study the combined interaction between two humic acids extracted from two different sources and

three neutral salts varied in their cationic valences on barley variety Giza 123 (*Hordium vulgare L* ) growth and elements uptake content and their uptake.

The first humic acid (HAS) used in this study was extracted from the alluvial soil collected from the Experimental Farm, Faculty of Agriculture, Minufia University were as the second one (HAC) was extracted from the composted clover straw. These humic acids were extracted, fractionated and purified according to the methods described by Kononova (1966), Posner (1966), Chen *et al.* (1978) and Schnitzer & Khan (1978). The purified humic acids content of C, N, P and H was determined according to Cottenie *et al.* (1982) for total organic-C; Bremner & Mulvaney (1982) for total-N; Olsen and Sommers (1982); Mann and Sounders (1966) (1966) for H-content respectively. Humic acids content of oxygen (O) was calculated by subtracting the content (%) of C, N, P and H from the total of 100 % Ash content (%) of these humic acids was estimated by burning the oven dry humic acid at 750 °C for 24 hrs (Holder and Griffith, 1983). The obtained results of the elemental composition and the calculated atomic ratios for the two humic acids were recorded in Table (1-a). Also, the studied humic acids contents of total acidity and some functional groups. i.e. carboxyl (COOH), total-OH, phenolic-OH and alcoholic -OH were determined according to the methods described by Dragunova (1958); Kukhareko (1937) and Brooks *et al.* (1958) and the obtained data were recorded in Table (1-b).

## **Sandy culture preparation.**

Sand used in this study was taken from desert part of Quessna region, Minufia Governorate. Sand was sieved through a 2 mm sieve, washed by tap water, treated with diluted HCl (6%) and H<sub>2</sub>O<sub>2</sub> (30%) to remove the carbonate and oxidize the organic matter, respectively. The treated sand was washed several times with tap water followed by distilled water. The refined sand was air-dried kept for using.

## ***Interaction between humic acids and neutral salts on plant growth***

**Table (1-a): Elemental composition (%), atomic ratios and ash content (%) of the studied humic acids.**

Humic acids	Elemental composition (%)					Atomic ratios				Ash content (%)
	C	H	N	P	O	C/H	C/O	C/N	C/P	
HAS	46.54	6.15	2.25	0.85	44.21	7.57	1.05	20.68	54.75	1.85
HAC	43.85	5.28	2.70	0.63	47.44	8.30	0.92	16.24	69.60	1.70

**Table (1-b): The tested humic acids content (meq / 100g HA ) of total acidity and some functional groups.**

Humic acid source	Total acidity	COOH	Total - OH	Phenolic - OH	Alcoholic - OH
HAS	580.4	270.1	445.8	310.3	135.5
HAC	710.50	330.4	527.6	380.1	182.5

### **Stocks of Hoagland solution were prepared as:**

#### **a- Macronutrients:-**

Solutions of the macronutrients were prepared by dissolving each salt in one liter solution, namely. 236 g of  $\text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$ , 101 g of  $\text{KNO}_3$ , 136 g of  $\text{KH}_2\text{PO}_4$  and 246 g of  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ .

#### **b- Micronutrients:-**

Solutions of the micronutrients were prepared by dissolving each salt in one liter solution, namely 2.86 g of  $\text{H}_3\text{BO}_3$ , 1.81 g of  $\text{MnCl}_2 \cdot \text{H}_2\text{O}$ , 0.22 g of  $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ , 0.08 g of  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$  and 0.02 g of  $\text{H}_2\text{MoO}_4 \cdot 4\text{MnO}$ , Iron citrate in 100 ml distilled water.

#### **Prepared Hoagland solution:-**

Hoagland solution was prepared by mixing 5ml of  $\text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$ , 5 ml of  $\text{KNO}_3$ , 1 ml of  $\text{KH}_2\text{PO}_4$ , 2 ml of  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$  and 1 ml from all micronutrient solution stocks and completed with distilled water to one liter volume.

#### **Expermental greenhouse setup.**

Their study was conducted on soil

Sciences Department . Faculty of Agriculture , Minufia University.

A 360 plastic pots with 20 cm inter diameter and 18 cm depth were used in this study. Each pot was filled by 1 kg clean and dried prepared sand. Each pot was planted by 12 grains of barley plants (*Hordium vulgare* L.) and irrigated every three days using Hoagland solution alternated with tap water to maintain the moisture content of the sandy culture 60 % of water holding capacity of sand. After 10 days of planting, the plants of each pot were thinned at 8 plants. After 21 days of planting, the pots were divided into two main groups (180 pot /main group ) representing the main factor or humic acids (HAS and HAC ) treatments. The pots of each main group were divided into equal five subgroups (36 pot for each sub group ) which treated by one application rate of humic acid (0, 10, 20, 40 and  $100 \text{ mg kg}^{-1}$  ). At the same time, the pots of each subgroup were divided into three sub subgroups representing the treatment of neutral salts ( $\text{NaCl}$ ,  $\text{CaCl}_2$ , and  $\text{FeCl}_2$ ). Finally, the pots of each sub sub group were divided into equal four groups ( 9 pot for each group ), where

the pots of each final group were treated by one concentration of the used neutral salts i.e. 0, 250, 500 and 1000 mg kg<sup>-1</sup>. The studied treatments were arranged in completely block design with three replicates. After 42 days of planting, the plants of each pot were taken as a whole, cleaned gently from sandy particles using current tap water, divided into shoots and roots, air-dried and oven-dried at 70 °C for 24 hrs and weighted to record the dry weights (g/pot) for both shoots and roots. The dried plant materials were finely ground and kept in glass bottles for its chemical analysis. The statistical design analysis for the dry matter yield carried out according to Gomez and Gomez (1984). The significant differences among means were tested using the least significant differences (L. S. D. ) at 5 % level of significance.

#### **Plant Analysis: -**

A 0.5 g of oven-dried plant sample was digested separately using 5 ml of mixture of conc. H<sub>2</sub>SO<sub>4</sub> and conc. HClO<sub>4</sub> at ratio of 3:1 on sandy hot plate up to become colorless( Chapman and Pratt, 1961 ). Then the digested product was diluted using distilled water and complete the volume up to 100ml. The final solution was kept in clean glass bottles for the following chemical analysis

- Sodium was determined using flame photometer as described by Cottenie *et al.* (1982).
- Calcium was determined by titration method with EDTA standard solution and ammonium murexide as indicator according to Lanyon and Heald (1982) as reported by Page *et al.*, (1982) .
- Iron was determined using atomic absorption according to the methods of described by Olsen and Ellis (1982) as reported by Page *et al.*, (1982) .

## **RESULTS AND DISCUSSION**

### **Effect of Humic Acid and Neutral Salts Application on Plant Growth.**

The present data in Table (2) show the effect of both source and application rate of humic acid individually or in presence of one chloride salts, i.e., Na, Ca and Fe at four application rates on dry weight (DW) of both shoots and roots of barley plants as g/pot. These data reveals that, increasing rate of added humic acids individually was associated by an increase of DW of both shoots and roots of barley plants. This trend was found under different application rates of the tested chloride salts. Such increases were related to the elemental composition and functional content of the tested humic acids. So, the highest values of dry weight of barley (shoots and roots) plants were found with the plants treated by HAC which characterized by low ratios of C/N and C/O. The enhanced effect of humic acids on plant growth was attributed to its content of many essential nutrients and improving growth media conditions. These results are in agreement with those obtained by Abou Hussien (1997) ; Atiyeh *et al.* (2002) ; Veronica *et al.* (2010) and Sadek and Sallam (2011).

With studied humic acids at different application rates under all treatments of NaCl, CaCl<sub>2</sub> and FeCl<sub>3</sub>, the found DW of shoots were higher than those of roots. These increases were significant for both shoots and roots and with the two humic acids. With different application rates of each humic acid, the obtained DW of barley plants varied widely according to the added salt and its application rates ( Table,2 ) this table show that, individual applications of NaCl, CaCl<sub>2</sub> and FeCl<sub>2</sub> appeared a wide effects on DW of shoots and roots. For example, with shoots and roots, increasing rates of added NaCl were associated by decrease of DW compared with that found with the control treatment. Such decreases may be resulted from the hazard effect of either of Na<sup>+</sup> or Cl<sup>-</sup> on plant growth and many metabolic processes within plant tissues. In this respect, similar decrease effect of NaCl on plant growth was found by El-Gundy (2005) and Nada and Tantawy (2013).

***Interaction between humic acids and neutral salts on plant growth***

**Table (2): The combined effect of both humic acids and neutral salts on shoots and roots dry weights (g/pot ) of barley plants.**

Humic acids treatment		Shoots					Roots				
Source	Added	Add neutral salt mg/Kg				Means	Add neutral salt mg/Kg				Means
		0	250	500	1000		0	250	500	1000	
<b>NaCl</b>											
HAS	0	1.40	1.20	1.04	0.963	1.155	0.662	0.646	0.620	0.500	0.607
	10	1.56	1.31	1.09	0.997	1.245	0.680	0.669	0.654	0.592	0.649
	20	1.65	1.41	1.10	1.025	1.300	0.731	0.698	0.673	0.601	0.676
	40	1.77	1.48	1.12	1.096	1.369	0.752	0.712	0.699	0.650	0.703
	100	1.89	1.51	1.32	1.101	1.456	0.760	0.723	0.701	0.699	0.721
	Mean	1.66	1.38	1.13	1.036	1.305	0.717	0.690	0.669	0.608	0.671
LSD(0.05)	0.63	0.43	0.35	0.371		0.497	0.278	0.193	0.115		
HAC	0	1.40	1.20	1.04	0.963	1.155	0.662	0.646	0.620	0.500	0.607
	10	1.58	1.46	1.29	1.201	1.385	0.989	0.901	0.819	0.796	0.876
	20	1.61	1.53	1.48	1.376	1.501	1.001	0.966	0.867	0.801	0.909
	40	1.86	1.81	1.77	1.573	1.756	1.630	1.136	1.000	0.899	1.166
	100	2.00	1.99	1.98	1.667	1.913	1.920	1.165	1.240	1.018	1.336
	Mean	1.69	1.60	1.51	1.356	1.542	1.240	0.963	0.909	0.803	0.979
LSD(0.05)	0.74	0.51	0.40	0.444		0.566	0.338	0.232	0.164		
<b>CaCl<sub>2</sub></b>											
HAS	0	1.409	1.940	1.825	1.610	1.696	0.662	0.730	0.630	0.621	0.661
	10	1.569	2.187	2.023	1.876	1.914	0.680	0.930	0.720	0.698	0.757
	20	1.657	2.358	2.245	2.102	2.091	0.731	1.112	0.966	0.745	0.889
	40	1.775	2.669	2.920	2.540	2.476	0.752	1.365	1.516	1.110	1.186
	100	1.896	2.879	3.089	2.830	2.674	0.760	1.430	1.621	1.356	1.292
	Mean	1.661	2.407	2.420	2.192	2.170	0.717	1.113	1.091	0.906	0.957
LSD(0.05)	0.274	0.643	0.982	0.913		0.211	0.417	0.771	0.603		
HAC	0	1.409	1.940	1.825	1.610	1.696	0.662	0.730	0.630	0.621	0.661
	10	1.582	2.410	2.354	2.214	2.140	0.989	1.230	1.031	0.953	1.051
	20	1.613	2.920	2.731	2.464	2.432	1.001	1.985	1.552	1.310	1.462
	40	1.866	3.462	3.654	3.365	3.087	1.630	1.996	2.113	1.985	1.931
	100	2.000	3.950	4.120	3.984	3.514	1.920	2.263	2.326	2.122	2.158
	Mean	1.694	2.936	2.937	2.727	2.574	1.240	1.641	1.530	1.398	1.452
LSD(0.05)	0.384	0.747	1.124	1.102		0.561	0.549	0.804	0.737		
<b>FeCl<sub>2</sub></b>											
HAS	0	1.41	2.19	2.59	2.50	2.172	0.66	0.93	1.00	0.95	0.886
	10	1.57	2.37	2.69	2.55	2.295	0.68	0.95	1.14	1.01	0.946
	20	1.66	2.56	2.80	2.65	2.415	0.73	1.08	1.21	1.19	1.050
	40	1.78	2.65	3.14	2.99	2.638	0.75	1.15	1.24	1.20	1.084
	100	1.90	2.87	3.28	3.44	2.872	0.76	1.19	1.34	1.38	1.167
	Mean	1.66	2.53	2.90	2.83	2.478	0.72	1.06	1.18	1.15	1.026
LSD(0.05)	0.289	0.392	0.699	0.823		0.532	0.804	0.737	0.813		
HAC	0	1.41	2.19	2.59	2.50	2.172	0.66	0.93	1.00	0.95	0.886
	10	1.58	2.51	2.66	2.65	2.352	0.99	1.38	1.62	1.47	1.365
	20	1.61	2.63	3.04	2.80	2.520	1.00	1.58	2.48	2.23	1.821
	40	1.87	2.69	3.59	3.38	2.882	1.63	2.59	2.86	2.70	2.445
	100	2.00	3.03	3.94	4.04	3.253	1.92	2.71	3.00	3.12	2.687
	Mean	1.69	2.61	3.16	3.07	2.636	1.24	1.84	2.19	2.09	1.841
LSD(0.05)	0.332	0.424	0.734	0.871		0.586	0.817	0.783	0.890		

The obtained DW of barley (shoots and roots ) plants in relation to added rates of  $\text{CaCl}_2$  individually as presented data in Table (2) show that, these weights were increased up to rate of 500 mg  $\text{CaCl}_2$  / kg compared control treatment and decreased at application rate of 1000 mg  $\text{CaCl}_2$  / kg compared with that found at low rates of added  $\text{CaCl}_2$ . These results were attributed to beneficial and promote effects of Ca on plant growth at low and medium rates of added  $\text{CaCl}_2$  in the growth media, but at added rate of 1000 mg / kg may be resulted in decrease of some metabolic processes especially in presence high concentration of  $\text{Cl}^-$  in growth media. These results are in agreement with these obtained by Hammad and Abou El-Khir (2005) and Fayed (2009). In addition, the presented data in Table (2) show that, individual application of  $\text{FeCl}_2$  at all application rates were associated by an increase of DW of barley (shoots and roots ) plants. These increases are related with positive and important role of Fe on plant growth and activity rates of metabolic processes (Alloway, 2008 ). These results are in agreement with these obtained by Abou Hussien (1997) ; Katkat *et al.* (2009) and El-Noamany (2013).

The data of interaction between different application rates of humic acids isolated from different sources and have varies chemical components and the three neutral salts i.e.,  $\text{NaCl}$ ,  $\text{CaCl}_2$  and  $\text{FeCl}_2$  which added at four application rates effects on DW of barley (shoots and roots ) plants as presented in Table (2) show that, decrease effect of  $\text{NaCl}$  on DW of barley shoots and roots was decreased as a result of plants treated by humic acids. In addition the increase effect of either of  $\text{CaCl}_2$  or  $\text{FeCl}_2$  on the obtained DW of shoots and roots were become more greater when these salts applied in combination with the humic acids. These increases were increased with the increase added rate of humic acids and varied from one to another. Under different treatments of the tested neutral salts, the highest values of DW of shoots and roots were associated the treatments of HAC.

These findings were in clear relations with the used humic acids elemental composition, atomic ratios and the content of functional groups (Abou Hussien, 1997 and Nada and Tantawy, 2013).

Data of the statistical (LSD at 0.05 ) of DW of barley (shoots and roots ) plants in relation with the studied treatments of humic acids and neutral salts individually or in combination are listed in Table (2). These data show that, individual application of humic acids resulted in a significant increase of DW, but there are a significant different between the used two humic acids effect on barley plants yield. The same data, also show that, the significant effect of individual applications of  $\text{NaCl}$ ,  $\text{CaCl}_2$  or  $\text{FeCl}_2$  was varied from one to another, where the high negative effect was associated the treatments of  $\text{NaCl}$  and the lowest one was found with  $\text{FeCl}_2$  treatments. The significant effects of neutral salts were become more positive when its applied in combination with humic acids. The latter effect was more clear with the plants treated by HAC. These findings are in harmony with used humic acids chemical composition and its content of functional groups. These results are in agreement with these obtained by Aydin *et al.* (2012) and Abd El-Kader *et al.* (2013).

The presented data in Table (3) show the relative change (RC) as a percent (%) of the obtained DW of both shoots and roots of barley plants in relation with the used humic acids under different types and application rates of some neutral salts. This table indicated that, at each rate of  $\text{NaCl}$ ,  $\text{CaCl}_2$  or  $\text{FeCl}_2$  RC values of DW with either of shoots or roots of barley plants were varied from acid to another. These values were increased with the increase of added HA. According to the found values of RC (%), the tested humic acids takes the order  $\text{HAC} > \text{HAS}$ . This trend was attributed to the humic acid content of functional groups and also its content of N and other nutrients (Abou Hussien, 1997 ). These results are in agreement with those obtained by Hussein and Hassan (2011) and Nada and Tantawy (2013).

***Interaction between humic acids and neutral salts on plant growth***

**Table (3): Relative change \*‘‘RC’’ (%) shoots and roots dry weights of barley plants planted in sandy culture as affected by different additives of both humic acids and neutral salts.**

Humic acids treatment		Added neutral salt (mg/kg)							
Source	Added (mg/kg)	Shoots (g / pot )				Roots ( g / pot )			
		0	250	500	1000	0	250s	500	1000
NaCl									
HAS	10	11.350	9.136	5.470	3.530	2.71	3.56	5.48	18.40
	20	17.600	17.690	5.660	6.438	10.45	8.05	8.55	20.20
	40	25.970	23.250	7.580	13.810	13.59	10.21	12.74	30.00
	100	34.560	24.660	27.150	14.330	14.84	11.91	13.06	39.80
HAC	10	12.270	18.330	24.470	24.710	49.54	39.47	32.09	59.20
	20	14.470	27.150	42.410	42.880	51.66	49.53	39.83	60.20
	40	32.430	50.330	70.440	63.340	146.90	75.85	61.29	79.80
	100	41.940	65.940	90.490	73.100	190.90	80.30	100.00	103.60
CaCl <sub>2</sub>									
HAS	10	11.35	12.73	11.12	16.52	2.71	27.39	14.28	12.39
	20	17.60	21.54	23.01	30.55	10.45	52.32	53.33	19.96
	40	25.97	37.57	60.00	57.76	13.59	86.98	140.60	78.74
	100	34.56	48.40	69.26	75.77	14.84	95.89	157.30	118.30
HAC	10	12.27	24.22	28.98	37.51	49.54	68.49	63.65	53.46
	20	14.47	50.51	49.64	53.04	51.66	171.90	146.30	110.90
	40	32.43	78.45	100.00	109.00	146.90	173.40	235.30	219.60
	100	41.94	103.60	177.90	147.40	190.90	210.10	269.20	241.70
FeCl <sub>2</sub>									
HAS	10	11.35	8.18	3.94	2.00	2.71	2.08	14.41	6.44
	20	17.60	17.01	7.99	5.80	10.45	15.46	20.72	25.15
	40	25.97	21.17	21.06	19.55	13.59	23.52	23.72	26.00
	100	34.56	31.13	26.54	37.62	14.84	27.38	34.03	45.47
HAC	10	12.27	14.86	2.70	5.96	49.54	48.33	62.16	54.73
	20	14.47	20.11	17.36	11.91	51.66	69.28	148.04	134.70
	40	32.43	23.04	38.46	35.18	146.90	178.40	185.78	184.30
	100	41.94	38.04	51.81	61.57	190.90	191.40	199.70	228.20

Dry matter yield of treated plants – Dry matter yield of untreated plants.  
 \* RC=  $\frac{\text{Dry matter yield of treated plants} - \text{Dry matter yield of untreated plants}}{\text{Dry matter yield of untreated plants}} \times 100$ .

In addition the values of RC (%) of DW of shoots and roots varied from low to high values in the treatments of neutral salts according to added salt and its application rate (Table,3). The low values of RC of DW were found with the plants treated by NaCl and become more lowest at high application rate of NaCl especially with low application rate of humic acids. Also, with CaCl<sub>2</sub> treatments, the data indicated that, for both shoots and roots, the highest RC (%) values of DW were found with application rate at 1000 mg. These findings were found with the tested humic acids at different application rates. In addition, RC values of the plants treated with FeCl<sub>2</sub> takes the reversal trend reported with CaCl<sub>2</sub> at different application rates under different treatments of humic acid. These findings were observed with shoots and roots for DW. These findings showed that, humic acids additives with neutral salts decreased its stress or its hazard effects on plant growth. This beneficial effect attributed to the improve effect of humic acids on growing media and its as a good source for many essential nutrients. Moreover presence humic acids in growing media increased water availability and uptake by plants, (Hussein and Hassan, 2011 and Nada and Tantawy, 2013).

Also, the obtained values of RC indicated that, NaCl additions were associated by high stress on plant growth, where the lowest one was associated the treatments of NaCl. This trend may be attributed to the type and strong complexes formed between NaCl, CaCl<sub>2</sub> or FeCl<sub>2</sub> with humic acids, where these complexes strong takes the order: CaCl<sub>2</sub> > FeCl<sub>2</sub> > NaCl. Many authors showed that, ion humic acid complexes become more stable and strong with the valence ion increase (Stevenson 1994 and Abou Hussien *et al.*,2002 ).

The presented data in Table (4) show, the calculated values of agronomical

efficiency (AE) of humic acids as mg dry plant materials / mg humic acid in relation with source and application rates of humic acid individually or in combination with three neutral salts, i.e., NaCl, CaCl<sub>2</sub> and FeCl<sub>2</sub> used at rates of 250, 500 and 1000 mg / kg with, AE values calculated with the humic acids for both shoots and roots of barley plants were decreased with the increase rate of added humic acids and varied from acid to another. With the same rate of added humic acids and according to AE values, these acids may be arranged in the following order HAC > HAS.

This order in harmony with these humic acids content of total acidity, functional groups and essential nutrients, i.e., C, N, H, O and others. Also, the same data showed that, the values of AE for the humic acids with shoots were higher than these found with roots. These results are in agreement with these obtained by Tonder (2008) ; Celik *et al.* (2008) ; Katkat *et al.* (2009 ) and Aydin *et al.* (2012).

In addition, the AE values of humic acids for DW of barley plants as affected by different additives of humic acids in combination with neutral salts appeared a wide variations depending on neutral salt type and its application rate ( Table,4). For example, with the humic acids, AE values were decreased with the increase rate of added NaCl and increased with the increase of added CaCl<sub>2</sub> and FeCl<sub>2</sub> up to 500 mg/ kg and decreased at application rate of 1000 mg / kg. These findings were found with DW for shoots and roots, mostly. These findings also reveals that NaCl have a greater stress on plant growth compared with that associated the treatments of either of CaCl<sub>2</sub> or FeCl<sub>2</sub>. These results means that humic acids additives resulted in a decrease of salinity stress and its effect on plant growth. In this respect El-Gundy (2005) ; Emam (2011) and Nada and Tantawy (2013) obtained on similar results.



***Interaction between humic acids and neutral salts on plant growth***

**Table (4): Agronomical efficiency \*\*“AE” of shoots and roots of barley plants (mg/mg HA) planted in sandy culture as affected by different additives of humic acids and neutral salts.**

Humic acids treatment		Shoots					Roots				
Source	Added	Add neutral salt mg/Kg				Means	Add neutral salt mg/Kg				Means
		0	250	500	1000		0	250	500	1000	
NaCl											
HAS	10	16.00	11.00	5.70	3.40	9.03	1.80	2.30	3.40	9.20	4.18
	20	12.40	10.65	2.95	3.10	7.28	3.45	2.60	2.65	5.05	3.44
	40	9.15	7.00	1.97	3.32	5.36	2.25	1.65	1.98	3.75	2.41
	100	4.87	2.97	2.83	1.38	3.01	0.98	0.77	0.81	1.99	1.14
	Mean	8.48	6.32	2.69	2.24	4.93	1.70	1.46	1.77	4.00	2.23
HAC	10	17.20	25.70	25.50	23.80	23.05	32.70	25.50	19.90	29.60	26.93
	20	10.20	16.35	22.10	20.65	17.33	16.95	16.00	12.35	15.05	15.09
	40	11.42	15.15	18.35	15.25	15.04	24.20	12.25	9.50	9.97	13.98
	100	5.91	7.94	9.43	7.04	7.58	12.58	5.19	6.20	5.18	7.29
	Mean	8.95	13.03	15.08	13.35	12.60	17.29	11.79	9.59	11.96	12.66
CaCl <sub>2</sub>											
HAS	10	16.00	24.70	19.80	26.60	21.78	1.80	20.00	9.00	7.70	9.63
	20	12.40	20.90	21.00	24.60	19.73	3.45	19.10	16.80	6.20	11.39
	40	9.15	18.22	27.37	23.25	19.50	2.25	15.87	22.15	12.22	13.12
	100	4.87	9.39	12.64	12.20	9.78	0.98	7.00	9.91	7.35	6.31
	Mean	8.48	14.64	16.16	17.33	14.15	1.70	12.39	11.57	6.69	8.09
HAC	10	17.30	47.00	52.90	60.40	44.40	32.70	50.00	40.10	33.20	39.00
	20	10.20	49.00	45.30	42.70	36.80	16.95	62.75	46.10	34.45	40.06
	40	11.42	38.05	45.72	43.87	34.77	24.20	31.57	37.07	34.10	31.74
	100	5.91	20.10	22.95	23.74	18.18	12.58	5.33	16.96	15.01	12.47
	Mean	11.18	30.83	33.37	34.14	26.83	17.29	29.93	28.05	23.35	24.65
FeCl <sub>2</sub>											
HAS	10	16.00	17.90	10.20	5.00	12.28	1.80	2.00	14.40	5.80	6.00
	20	12.40	18.60	10.35	7.25	12.15	3.45	7.20	10.35	11.95	8.24
	40	9.15	11.57	13.65	12.22	11.65	2.25	5.48	5.93	6.18	4.96
	100	4.87	6.81	6.88	9.41	6.99	0.98	2.55	3.40	4.32	2.81
	Mean	8.48	10.98	8.22	6.78	8.61	1.70	3.45	6.82	5.65	4.40
HAC	10	17.20	32.50	7.00	14.90	17.90	32.70	45.00	62.10	52.00	47.95
	20	10.20	22.00	22.50	14.90	17.40	16.95	32.25	73.95	64.00	46.79
	40	11.42	12.60	24.92	22.00	17.74	24.20	41.52	46.40	43.77	38.97
	100	5.90	8.47	12.73	15.40	10.63	12.58	17.82	19.96	21.68	18.01
	Mean	8.94	15.11	13.43	13.44	12.73	17.29	27.32	40.48	36.29	30.34

Dry matter yield of treated plants– Dry matter yield of untreated plants.

$$*AE = \frac{\text{Dry matter yield of treated plants} - \text{Dry matter yield of untreated plants}}{\text{Added humic acid (mg kg}^{-1}\text{)}}$$

## **Effect of Humic Acids and Neutral Salts on Plant Chemical Composition.**

### **a. Sodium (Na) content.**

The presented data in Table (5) show barley plants (shoots and roots) concentration (mg/kg) and uptake (mg / pot ) of Na in relation with both humic acids isolated from different sources and NaCl at different application rates. This table shows that, with both shoots and roots of barley plants Na concentration was increased with the increase of added NaCl individually.

Also, at the same individual application rate of NaCl, Na concentration of shoots was higher than that in roots. On the other hand, individual NaCl additions at 250 mg/kg was associated by increase of Na uptake, but at high application rate, i.e., 500 and 1000 mg/kg resulted in a decrease of Na uptake. These findings were found in shoots, while Na uptake in roots was increased with increase rate of added NaCl. The latter results were attributed to the reductions found in the dry matter yield of shoots and roots at high rates of added NaCl. In this respect, similar results were obtained by Hammad and Abo El-Khir (2005) and Nada and Tantawy (2013).

The presented data in Tables (6) show that, increasing of added rates individually of the tested humic acids was associated by a decrease of Na concentration (mg/kg) of both shoots and roots of barley plants. This decrease was attributed to the found increase of barley plants growth associated with the treatments of humic acids. This effect namely by dilution effect (Marschner, 1998). So, most individual treatments of humic acids resulted in a decrease of Na uptake by both shoots and roots. Such this decrease was become more high at high rates of added humic acids. At the individual application rate of humic acids, the found decrease of Na concentration and its uptake by either of shoots or roots was varied widely from acid to another depending on the chemical composition of the tested humic acids and its effect on plant growth and elements uptake. So, the high Na content was found in the plants treated by HAC. These results are in agreement with those obtained by, Abou Hussien (1997);

Abou Hussien *et al.* (2002) ; El-Desuki (2004) and Shaaban *et al.* (2009).

In addition application humic acids and NaCl at different rates in combination appeared a wide variations in their effect on Na concentration and uptake by shoots and roots of barley plants (Tables, 6). Humic acids application reduced Na concentration and uptake by shoots and roots compared with those found in the individual treatments of NaCl but this content was higher than associated with the individual treatments of humic acids. These results mean that, Na may be weakly retained by humic acids and become less available for uptake by plant. Meloni *et al.* (2001 and 2004) ; Turan and Aydin (2005) ; El-Gundy (2005) and Aydin *et al.* (2012).

### **b. Calcium (Ca) content.**

The presented data in Table (6) show individual and combined effect of both humic acids isolated from different sources and CaCl<sub>2</sub> at different application rates on barley plants concentration (mg/kg) of Ca and its uptake (mg/pot). These data show that, Ca concentration and uptake by both shoots and roots was increased with the increase of added CaCl<sub>2</sub> as alone. This may be considered as natural results which attributed to the high concentration of Ca in growth media. With the same rate of CaCl<sub>2</sub> individual application Ca concentration and uptake by shoots were higher than those found with roots. In this respect Hammad and Abou El-Khir (2005) and Nada and Tantawy (2013) obtained similar results. In addition the data reveals that with, both Ca concentration (mg kg<sup>-1</sup>) of shoots and roots was decreased with the increase of added humic acids as alone. Such this decrease was resulted from the high dry matter yield of shoots and roots associated with the high rates of added humic acids as common by dilution effect (Marschner, 1998). The rate of this decrease was decreased with the increase of humic acid application rate. Also, Ca concentration in both shoots and roots was varied with from humic acid to another, where high Ca concentration of shoots and roots was recorded with different application rates of humic acid isolated from soil (HAS).

***Interaction between humic acids and neutral salts on plant growth***

**Table (5): Sodium concentration (mg/kg) and uptake (mg/pot ) in shoots and roots of barley plants of as affected by source and application rates of humic acids under different application rates of NaCl**

Humic acids treatment		Conc. mg-Na/kg					Uptake mg- Na/ pot				
Source	Added (mg/kg)	Add NaCl. mg/kg				Means	Add NaCl mg/kg				Means
		0	250	500	1000		0	250	500	1000	
<b>Shoots</b>											
HAS	0	1026	11596	12960	14070	9913.0	1.45	13.96	13.50	13.54	10.61
	10	1020	11520	12825	14000	9841.2	1.60	15.13	14.09	13.95	11.19
	20	840	10560	12150	12600	9037.5	1.39	14.96	13.37	12.91	10.66
	40	720	8960	10800	11900	8095.0	1.28	13.29	12.10	13.04	9.93
	100	600	8320	9450	10500	7217.5	1.14	12.48	12.52	11.56	9.42
	Mean	841	10191	11637	12614	8820.8	1.37	13.96	13.12	13.00	10.36
HAC	0	1026	11596	12960	14070	9913.0	1.45	13.96	13.50	13.54	10.61
	10	858	10880	12150	13300	9297.0	1.58	15.89	15.75	15.97	12.30
	20	686	9600	11643	12460	8597.2	1.11	14.69	17.27	17.14	12.55
	40	429	8000	9450	10500	7094.7	0.80	14.48	16.78	16.51	12.14
	100	384	6880	8775	10150	6547.2	0.77	13.74	17.41	16.92	12.21
	Mean	677	9391	10996	12096	8289.8	1.14	14.55	16.14	16.02	11.96
<b>Roots</b>											
HAS	0	516	5824.0	7155.0	8120	5403.750	0.341	3.762	4.436	4.807	3.34
	10	510	5760	7087	7980	5334.250	0.346	3.853	4.634	4.724	3.39
	20	480	54400	6615	7000	4883.750	0.350	3.797	4.451	4.207	3.20
	40	456	5120	6210.0	6720	4626.500	0.342	3.645	4.340	4.368	3.17
	100	408	4736	5535.0	6300	4244.750	0.310	3.424	3.880	4.403	3.00
	Mean	474	5376	6520.4	7224	4898.60	0.338	3.696	4.348	4.502	3.22
HAC	0	516.0	5824.00	7155.0	8120.00	5403.750	0.341	3.762	4.436	4.807	3.34
	10	492	5760	7425	8400	5519.250	0.486	5.189	6.081	6.686	4.61
	20	468	5376	6750	7840	5108.500	0.468	5.193	5.852	6.279	4.45
	40	420	4864	6210	7000	4623.500	0.684	5.525	6.21	6.293	4.68
	100	300	3712	4995	6300	3826.750	0.576	4.324	6.193	6.413	4.38
	Mean	439.2	5107.2	6507.0	7532.0	4896.350	0.511	4.799	5.754	6.096	4.29

**Table (6): Calcium concentration (mg/kg) and uptake (mg/pot ) and its relative change (RC ) percent (%) in shoots and roots of barley plants of as affected by source and application rates of humic acids under different application rates of CaCl<sub>2</sub>.**

Humic acids treatment		Conc. mg-Na/kg					Uptake mg- Na/ pot				
		Add CaCl <sub>2</sub> . mg/kg				Means	Add CaCl <sub>2</sub> mg/kg				Means
Source	Added (mg/kg)	0	250	500	1000		0	250	500	1000	
<b>Shoots</b>											
HAS	0	855.0	9060.0	9600.0	10050.0	7391.3	1.204	17.570	17.520	16.180	13.119
	10	850.0	9000.0	9500.0	10000.0	7337.5	1.333	19.680	19.210	18.760	14.746
	20	700.0	8250.0	9000.0	9000.0	6737.5	1.159	19.450	20.200	18.910	14.930
	40	600.0	7000.0	8000.0	8500.0	6025.0	1.065	18.680	23.360	21.590	16.174
	100	500.0	6500.0	7000.0	7500.0	5375.0	0.948	18.710	21.620	21.220	15.625
	Mean	701.0	7962.0	8620.0	9010.0	6573.3	1.142	18.818	20.382	19.332	14.918
HAC	0	855.0	9060.0	9600.0	10050.0	7391.3	1.204	17.570	17.520	16.180	13.119
	10	715	8500	9000	9500	6928.8	1.131	20.48	21.18	21.03	15.955
	20	572	7500	8625	8900	6399.3	0.922	21.9	23.55	21.92	17.073
	40	358	6250	7000	7250	5214.5	0.668	21.63	25.57	25.23	18.275
	100	320	5375	6500	6000	4548.8	0.64	21.23	26.78	28.88	19.383
	Mean	564.0	7337.0	8145.0	8340.0	6096.5	0.913	20.562	22.920	22.648	16.761
<b>Roots</b>											
HAS	0	430.0	4550.0	5300.0	5800.0	4020.0	0.284	3.320	3.339	3.601	2.636
	10	425.0	4500.0	5250.0	5700.0	3968.8	0.289	4.185	3.780	3.978	3.058
	20	400.0	4250.0	4900.0	5000.0	3637.5	0.292	4.726	4.730	3.725	3.368
	40	380.0	4000.0	4600.0	4800.0	3445.0	0.285	5.460	6.973	5.328	4.512
	100	340.0	3700.0	4100.0	4500.0	3160.0	0.258	5.290	6.646	6.102	4.574
	Mean	395.0	4200.0	4830.0	5160.0	3646.3	0.282	4.596	5.094	4.547	3.630
HAC	0	430.0	4550.0	5300.0	5800.0	C114:F114	0.284	3.320	3.339	3.601	2.636
	10	410	4500	5500	6000	4102.5	0.405	5.535	5.67	5.718	4.332
	20	390	4200	5000	5600	3797.5	0.39	8.337	7.76	7.336	5.956
	40	350	3800	4600	5000	3437.5	0.389	7.584	9.719	9.925	6.904
	100	250	2900	3700	4500	2837.5	0.48	6.562	8.606	9.549	6.299
	Mean	366.0	3990.0	4820.0	5380.0	3543.8	0.390	6.268	7.019	7.226	5.225

## ***Interaction between humic acids and neutral salts on plant growth***

---

These results takes the the reversible trend for the effect of these humic acids on obtained dry matter yield of barley plants. On the other hand, with individual additives of humic acids, Ca uptake (mg/ pot ) for both shoots and roots of barley plants was decreased with the increase rate of added humic acid (Table, 6 ) in mostly. This decrease effect was varied from humic acid to another. The highest uptake of Ca uptake by shoots and roots was found in the plants treated by HAC. These findings were found with all tested rates of the humic acids. Such this increase was related with found dry matter yield of shoots of barley plants. These results are in agreement with those obtained by Hussein and Hassan (2011) and Aydin *et al.* (2012).

Regarding to the results of combined treatments of humic acids and  $\text{CaCl}_2$  at different application rates on Ca concentration ( $\text{mg kg}^{-1}$ ) and uptake ( $\text{mg pot}^{-1}$ ) by shoots and roots of barley plants as listed in Table (6) may be observed that, humic acids additives in combination with  $\text{CaCl}_2$  reduced Ca concentration and uptake at the same rate of added  $\text{CaCl}_2$  compared with that found in the plants untreated by humic acids. This decrease was become more clear at high application rate of humic acids. The rate of this decrease was varied from humic acid to another depending on its content of total acidity and functional groups. The lowest one was found in the plants treated by HAS at low application rate. This trend was found with all application rates of  $\text{CaCl}_2$ . These findings of decrease of Ca concentration with humic acids additives was attributed to chelating action for these humic acids to Ca as Ca - humate and complex which become less available to uptake by plants (Stevenson, 1994 ). Chelating action or reducing Ca solubility was varied from humic acid to another, where this effect was increased with the increase of humic acid content of total acidity and functional groups. So, at the same application rate of the used humic

acids the high decrease of Ca concentration was found in both shoots and roots of barley plants treated by HAC.

### **c. Iron (Fe ) content.**

The presented data in Table (7) show the effect of individual and combined treatments of humic acids and  $\text{FeCl}_3$  at different application rates of them on Fe concentration ( $\text{mg kg}^{-1}$ ) and uptake ( $\text{mg pot}^{-1}$ ) by shoots and roots of barley plants. These data show that, Fe concentration and uptake were increased with the increase of added  $\text{FeCl}_2$  as alone. This trend was found with both shoots and roots. Under the same individual treatment of  $\text{FeCl}_2$ , Fe concentration of shoots was higher than that of roots. Nearly similar trend of Fe uptake was found with the individual treatment of  $\text{FeCl}_2$ . These findings attributed to the enhanced effect of Fe on plant growth and enzymes activity. In this respect, Abou Hussien (1997) and El-Noamany (2013) obtained on similar results.

The effect of individual treatments of humic acids on Fe concentration as presented in Table (7) show that, increasing rate of added humic acids was associated by decrease of Fe concentration in both shoots and roots. The rate of this decrease was become more clear at high application rates of added humic acids. Also this effect was varied from humic acid to another. The found decrease of Fe concentration attributed to the found increase of dry matter yield of barley plants associated humic acids treatments. This effect normally named by dilution effect ( Marschner, 1998 ). So, the high concentration was found in the plants treated by HAS. This trend was observed with both shoots and roots. With all combined treatments of humic acids and  $\text{FeCl}_2$  at different application rates on Fe concentration of shoots was higher than that of roots. In this respect, Abou Hussien (1997) and Abou Hussien *et al.* ( 2002 ) obtained on similar results.

**Table (7): Iron concentration (mg/kg) and uptake (mg/pot in shoots and roots of barley plants of as affected by source and application rates of humic acids under different application rates of FeCl<sub>2</sub>.**

Humic acids treatment		Conc. mg-Na/kg					Uptake mg- Na/ pot				
		Add FeCl <sub>2</sub> . mg/kg				Means	Add FeCl <sub>2</sub> mg/kg				Means
Source	Added (mg/kg)	0	250	500	1000		0	250	500	1000	
<b>Shoots</b>											
HAS	0	1150.0	5900.0	6500.0	7000.0	5137.5	1.620	12.900	16.840	17.500	12.215
	10	1102.0	5850.0	5900.0	6100.0	4738.0	1.729	13.840	15.890	15.560	11.755
	20	975.0	5500.0	5800.0	6000.0	4568.8	1.615	14.070	16.230	15.870	11.946
	40	967.0	5300.0	5600.0	5900.0	4441.8	1.716	14.040	17.570	17.640	12.742
	100	890.0	5100.0	5400.0	5700.0	4272.5	1.687	14.620	17.710	19.610	13.407
	Mean	1016.8	5530.0	5840.0	6140.0	4631.7	1.673	13.894	16.848	17.236	12.413
HAC	0	1150.0	5900.0	6500.0	7000.0	5137.5	1.620	12.900	16.840	17.500	12.215
	10	1080	5750	5850	6000	4670.0	1.708	14.44	15.57	15.9	11.905
	20	965	5200	5600	5800	4391.3	1.556	13.66	17.03	16.23	12.119
	40	940	5000	5400	5700	4260.0	1.754	13.45	19.38	19.27	13.464
	100	880	4950	5200	5500	4132.5	1.76	15.01	20.46	22.22	14.863
	Mean	1003.0	5360.0	5710.0	6000.0	4518.3	1.680	13.892	17.856	18.224	12.913
<b>Roots</b>											
HAS	0	900.0	4900.0	5200.0	6000.0	4250.0	0.5950	4.5610	5.1940	5.7000	4.0125
	10	880.0	4800.0	5100.0	5900.0	4170.0	0.5980	4.5640	5.8290	5.9470	4.2345
	20	846.0	4400.0	4900.0	5700.0	3961.5	0.6180	4.7300	5.9090	6.7770	4.5085
	40	805.0	4150.0	4600.0	5400.0	3738.8	0.6050	4.7720	5.6850	6.4600	4.3805
	100	770.0	4000.0	4750.0	5100.0	3655.0	0.5850	4.7440	6.3600	7.0480	4.6843
	Mean	840.2	4450.0	4910.0	5620.0	3955.1	0.6002	4.6742	5.7954	6.3864	4.3641
HAC	0	900.0	4900.0	5200.0	6000.0	4250.0	0.5950	4.5610	5.1940	5.7000	4.0125
	10	850.0	4650.0	4950.0	5700.0	4037.5	0.84	6.42	8.019	8.379	5.9145
	20	810.0	4300.0	4800.0	5550.0	3865.0	0.81	6.776	11.89	12.37	7.9615
	40	790.0	4050.0	4550.0	5300.0	3672.5	1.287	10.49	12.99	14.31	9.7693
	100	740.0	3800.0	4300.0	4900.0	3435.0	1.42	10.3	12.87	15.27	9.9650
	Mean	818.0	4340.0	4760.0	5490.0	3852.0	0.9904	7.7094	10.1926	11.2058	7.5246

## ***Interaction between humic acids and neutral salts on plant growth***

The presented data in Table (7) show the effect of combined treatments of humic acids and FeCl<sub>2</sub> at different application rates of them on Fe content in shoots and roots of barley plants. These data show that, at the same rate of FeCl<sub>2</sub> application, increasing rate of added humic acids was associated by decrease of Fe concentration by shoots and roots of barley plants while the Fe uptake was increased. The rate of this effect was increased with the increase rate of added humic acids and varied from acid to another. With different application rates of FeCl<sub>2</sub>, barley plants treated by HAS characterized by high concentration of Fe. This trend was in harmony with the named by dilution effect. At the same rate of each humic acid application, increasing application rates of FeCl<sub>2</sub> was associated by increase of shoots and roots of barley plants content of Fe. This increase resulted from increase of soluble Fe in growth media, but the found decrease of this content which found with the increase of added humic acids together with FeCl<sub>2</sub> attributed to chelation effect of these acids for Fe and converted to insoluble form followed by decrease Fe uptake. These results are in agreement with those obtained by Abou Hussien *et al.* (2002)

### **REFERENCES**

- Aakl, A. A. (1998). Interaction of humic substance with trace elements. M. Sc. Thesis, Fac. of Agri. Minufia Univ., Egypt.
- Abd El- Kader, M.G. and H.M.G. El-Shaboury (2013). Evaluation of soaking and foliar methods with compost tea, humic acid and bio-fertilizer on soil fertility and faba bean yield productivity and quality under saline soil conditions. *Minufia J. Agric. Res.*, Vol. 38 No. 6(3): 1663-1675.
- Abou Hussien, E. A. (1997). Evaluation of humic acids as iron carriers for plant nutrition. *Minufia J. Agric. Res.*, 22 (2): 581-611.
- Abou Hussien, E. A., R. A. Khalil and A. A. Aakl (2002). Adsorption of Fe and Zn by clay – humic complex. *Egypt J. Soil Sci.*, 42 (1): 127-138.
- Alloway, B.J. (2008). Copper and zinc in soils: Too little or too much in NZ Trace Elements Group Conference. Waikato University, Hamilton, Oct. 13-15.
- Atiyeh, R. M., S. Lee, C. A. Edwards, N. Q. Arancon and J. D. Metzger (2002). The influence of humic acids derived from earthworm processed organic wastes on plant growth. Soil Ecology laboratory, 105 Botany and zoology Building. The Ohio State University, 1735 Neil Avenue, Columbus, O.H 43210, USA.
- Aydin, A., K. Canan and T. Metin (2012). Humic acid application alleviate salinity stress of *Phaseolus vulgaris* plants decreasing membrane leakage. *African Journal of Agricultural Res.* 7(7): pp. 1037-1086.
- Brooks, J. D., R. A. Durie and S. Sternhell (1958). Chemistry of brown coal: 3-pyrolytic reaction. *Aust. J. Appl. Sci.*, 9: 303-320.
- Bremmes, J. M. and C. S. Mulvaney (1982). Nitrogen- Total. Pages 595- 624 in A.L. page., R. M. Miller and D.R. Keeney (eds.) *Methods of Soil Analysis. Part (2<sup>nd</sup> Ed.) : Chemical and Microbiological Properties.* ASA, Inc., SSSA, Inc. publishes, Madison, Wisconsin, USA.
- Brunetti, G., C. Plaza and N. Seneri (2005). Oliva pomace amendment in mediterranean conditions. Effect on soil and humic acid properties and (wheat *Triticum aestivum* L ) yield. *J. Agric. Food Chem.*, 53 (17): 6730 – 6737.
- Celik, H., A. V. Katkat, B. B. Asik and M. A. Turan (2008). Effects of soil application of humus on dry weight and mineral nutrients uptake of maize under calcareous soil conditions. *Archives of Agronomy and Soil Sci.*, 54 (6): 605-614.
- Chapman, H. D. and P. F. Pratt (1961). "Methods of Analysis for Soils, Plants and Water". Univ. of California Agric. Sci. Preiced publication. 4034, p 50.
- Chen, Y., N. Senesi and M. Schnitzer (1978). Chemical and physical characteristics of humic acids extracted from soils of Mediterranean region. *Geoderma*, 20: 87- 104.
- Cottenie, A., M. Verloo, L. Kikens, G. Velghe and R. Camerlynck (1982). "Analytics

- Problems and Methods in Chemical Plant and Soil Analysis". Handbook, Ed. A. Cottenie. Ghent. Belgium.
- Dragunova, A. E. (1958). "A Rapid Method for Determining Functional Groups in Humic Acids". Nauch. Trydy. Mosk. In. Zh-a Konon. Inst. ser. Khi. Droizvod. C. F. Kononova (1966) p. 410.
- El-Desuki, M. (2004). Response of onion plants to humic acid and mineral fertilizers application. *Annals of Agric. Sci. Moshtohor*, 42 (4): 1955- 1964.
- El-Gundy, A. G. (2005). Effect of different amendments on sodic soil properties and plant growth. M. Sc. Thesis, Fac. of Agric. Minufia Univ., Egypt.
- El-Noamany, Naglaa A. E. (2013). Studies on diazotrophy in soils under different conditions. M. Sc. Thesis, Fac. of Agric. Minufia Univ., Egypt.
- Emam, A. A. M. (2011). Effect of irrigation water salinity on organic matter in soil and status of some nutrients. M. Sc. Thesis, Fac. Of Agric., Minufia Univ., Egypt.
- Fayed, S. A. S. (2009). Effect of irrigation with sewage and drainage water on properties of some new reclaimed soils and plant Growth. M. Sc. Thesis, Fac. of Agric., Minufia Univ., Egypt.
- Gomez, K. A. and A. A. Gomez (1984). "Statistical Procedures for Agricultural Research". 2<sup>nd</sup> Ed., Wiley, New York.
- Hammad, Salwa A. R. and A. M. Y. Abou El- Khir (2005). Effect of total salinity and salinity types on soybean growth and its chemical composition. *Menufia. J. of Agric. Res.*, 30 (3): 835-852.
- Holder, M. P. and S. M. Griffith (1983). Some characterization of humic materials in caribbean vertisols. *Can. J. Soil Sci.* 63: 151- 159.
- Hussein, K. H. and A. F. Hassan (2011). Effect of different humic acids on the nutrient content, plant growth and soil properties under conditions of salinity. *Soil and Water Res.*, 6 (1): 21-29.
- Katkat, A.V., H. Celik, M. A. Turan and B. B. Asik (2009). Effect of soil and foliar applications of humic substances on dry weight and mineral nutrients uptake of wheat plant under calcareous soil conditions. *Australian J. of Basic and Applied Sciences*, 3(2): 1266-1273.
- Kononova, M. M. (1966). "Soil Organic Matter Pergamon", Oxford, pp. 190-199.
- Kukhareko, T. A. (1937). "Reactions of Humic Acids With Neutral Salts. part 1. Khin. Tverd. Tool. 8 (9) & part.1 hid.8 (12) (C.F. Kononova 1966, pp. 411-412 ).
- Liu, C. (1998). Effects of humic substances on creeping bentgrass growth and stress tolerance. Ph. D. Thesis, philosophy Department of Crop Science, North Carolina State University, Raleigh.
- Mann, F. G. and B. C. Sounders (1966). *Practical organic matter 4 th Ed.* Western printing services L.T.D Bristol.
- Marschner, H. (1998). "Mineral Nutrition of Higher Plants". Harcourt Broce &Comp. Publishers. London, New York.
- Meloni, D. A., M. A. Oliva, H. A. Ruiz and C. A. Martinez (2001). Contribution of proline and inorganic solutes to osmotic adjustment in cotton under salt stress. *J. Plant Nutri.*, 24: 599-612.
- Meloni, D. A., M. R. Gulotta, C. A. Martinez and M. A. Oliva (2004). The effects of salt stress on growth, nitrate reduction and proline and glycinebetaine accumulation in *prosopis alba*. *Braz. j. Plant. Physiolo.*, 16:39-46.
- Nada, W. M. and Manal F. Tantawy (2013). Effect of humic acid applications on soil properties and plant tolerance for high salinity levels of irrigation water. *Egypt. J. Soil Sci.* (Under print).
- Nelson, D. W. and L.S. Sommers (1982). Total carbon, organic carbon and organic matter . pages 539-579. In A.L. page., R. M. Miller and D.R. Keeny (eds.) *Methods of Soil Analysis. Part (2<sup>nd</sup> Ed. ) : Chemical and Microbiological Properties.* ASA, Inc., SSSA, Inc. publishes, Madison, Wisconsin, USA.
- Olsen, S. R. and L. E. Sommers (1982). Phosphorus. Pages 403 – 430. In A.L. page., R. M. Miller and D.R. Keency (eds.) *Methods of Soil Analysis. Part (2<sup>nd</sup> Ed. ) : Chemical and Microbiological Properties.* ASA, Inc., SSSA, Inc. publishes, Madison, Wisconsin, USA.
- Page, A. L., R. H. Miller and D. R. Keeney (1982). "Methods of Soil Analysis". Part



## ***Interaction between humic acids and neutral salts on plant growth***

2. Chemical and Microbiological Properties. 2<sup>nd</sup> Edition, American Society of Agronomy. Inc, Soil Sci. Soci. of America, Madison, Wisconsin USA. pp. 149-985.
- Posner, A. M. (1966). The humic acid extracted by various reagents from a soil. Part 1. Yield in organic components and titration curves. J. Soil Sci., 17: 65-78.
- Sadek, Jacklin G. and Amany M. Sallam (2011). Effect of grains soaking with humic acid and micronutrients foliar spray on quality and productivity of rice plant under saline soil conditions. Minufia J. Agric. Res., 36 (1): 177- 196.
- Schnitzer, M. and S. U. Khan (1978). "Soil Organic Matter". El-Sevier Scientific Publishing Company Amsterdam, Oxford, New York.
- Shaaban, S. H. A., Manal F. M. and M. H. M. Afifi (2009). Humic acid foliar application to minimize soil applied fertilization of surface irrigated wheat. World journal of Agricultural Sciences 5 (2): 2007 – 2010.
- Stevenson, F.J. (1994). "Humus Chemistry: Genesis, Composition, reaction". 2<sup>nd</sup> Ed. John Wiley & Sons. New York.
- Stevenson, F.J. (1982). "Humus Chemistry: Genesis, Composition". John Wiley & Sons, New York.
- Tonder, J.T.V. (2008). Effect of potassium humate on soil properties and growth of wheat. Department Soil, Crop and Climate Sciences Faculty of Natural and Agricultural Sciences, University of the free State Bloemfontein.
- Turan, M. and A. Aydin (2005). Effects of different salt sources on growth, inorganic ions and proline accumulation in corn ( *Zea mays* L. ) Europ. J. Hortic. Sci., 70: 149 – 155.
- Veronica, M., E. Bacaicoa, M. Z. Angel, A. Elena, G. Maria, E. Marta and M. G. M. Jose (2010). Action of humic acid on promotion of cucumber shoot growth involves nitrate – related changes associated with the root – to – shoot distribution of cytokinins, polyamines and mineral nutrients. J. of Plant physiology, 167 (8): 633 – 642.

### **التأثير المشترك لأحماض الهيوميك والأملاح المتعادلة على نمو النبات منال فتحي طنطاوى<sup>(1)</sup> ، رفعت أحمد خليل<sup>(2)</sup> ، الحسيني عبد الغفار أبو حسين<sup>(2)</sup> ، أسماء مختار متولي سليمان<sup>(2)</sup>**

<sup>(1)</sup> معهد بحوث الأراضي والمياه والبيئة – مركز البحوث الزراعية – الجيزة.

<sup>(2)</sup> قسم علوم الأراضي – كلية الزراعة – جامعة المنوفية.

#### **الملخص العربي:**

أجريت هذه الدراسة لتقييم التأثير المشترك لاثنتين من أحماض الهيوميك مختلفي في مصدرهما وكذلك تركيبهما الكيميائي مع ثلاث من الأملاح المتعادلة على صورة كلوريد صوديوم وكلوريد كالسيوم وكلوريد حديدوز على نمو نبات الشعير (جيزة 123) ومحتواه من الصوديوم والكالسيوم والحديد وأجريت الدراسة كتجربة أصص في نظام قطع كاملة العشوائية في ثلاث مكررات باستخدام مزرعة رملية. تم استخلاص حامض الهيوميك من الأرض الرسوبية ومن كمبوست قش البرسيم وتم إضافة كل منهما عند معدلات إضافة صفر، 10، 20، 40 و 100 ملليجرام / كجم. أما عن الأملاح المتعادلة الثلاثة فقد أضيفت عند معدلات صفر، 250، 500 و 1000 ملليجرام / كجم تم حصاد النباتات من كل أصيص بعد 42 يوم من الزراعة وقد قدر الوزن الجاف لكل من المجموع الخضري والمجموع الجذري للنباتات المحصودة كما أجرى لها التحليل الإحصائي عند مستوى معنوية 0.05.

ازداد الوزن الجاف لكل من المجموع الخضري والمجموع الجذري لنباتات الشعير زيادة معنوية بزيادة المضاف من أحماض الهيوميك وكانت الزيادة المتحصل عليها في الأوزان الجافة في النباتات المعاملة بحامض الهيوميك المستخلص من الكمبوست أعلى نسبيا من تلك المتحصل عليها في معاملات حامض الهيوميك المستخلص من الأرض ومع جميع معاملات أحماض الهيوميك كان الوزن الجاف للسوق المتحصل عليه أعلى من مثيله للجذور. كانت استجابة الوزن الجاف لنبات الشعير قد اختلفت باختلاف الملح المتعادل وكذلك معدل إضافته.

تناقصت الكفاءة المحصولية لأحماض الهيوميك بزيادة معدل إضافة كلوريد الصوديوم ولكنها تزداد بزيادة المضاف من كلوريد الكالسيوم حتى معدل إضافة 500 ملجرام / كجم في حين استمرت هذه الزيادة مع كلوريد الحديدوز حتى معدل إضافة 1000 ملجرام / كجم. ازداد محتوى سوق وجذور نباتات الشعير من الصوديوم والكالسيوم والحديد بزيادة المضاف من كلوريد الصوديوم وكلوريد الكالسيوم وكلوريد الحديدوز، على الترتيب. زيادة المضاف من أحماض الهيوميك المستخلصة لعبت دورا كبيرا في إنقاص التأثير المثبط للملوحة على نمو النبات وكذلك امتصاصه للعناصر.