



Official Journal Issued by  
Faculty of  
Veterinary Medicine

## Benha Veterinary Medical Journal

Journal homepage: <https://bvmj.journals.ekb.eg/>



Since 1990

### Original Paper

## Dual functional properties of enzymatically hydrolyzed dromedary milk proteins with improved antioxidant and antibacterial activity

Dina A. B. Awad<sup>1,\*</sup>, Osama Ahmed<sup>2</sup>, Samar H. Baloza<sup>3</sup> and Maha Mamdouh<sup>4</sup>

<sup>1</sup> Department of Food Hygiene and Control, Faculty of Veterinary Medicine, Benha University, Qalyubia, Egypt.

<sup>2</sup> Anatomy and embryology department, faculty of veterinary medicine, Benha University, Egypt.

<sup>3</sup> Genetic and Genetic Engineering, Department of Animal Wealth Development, Faculty of Veterinary Medicine, Benha University, Egypt.

<sup>4</sup> Department of Physiology, Faculty of Veterinary Medicine, Benha University, Egypt.

### ARTICLE INFO

#### Keywords

Digestive enzymes  
Enzymatic hydrolysates  
Gram-negative bacteria  
SCM hydrolysates

Received 11/08/2023

Accepted 05/09/2023

Available On-Line  
01/10/2023

### ABSTRACT

Dromedary milk has a proven outstanding history of being a nutrient-rich food. Camel milk possesses both medicinal and technological advantages, including antioxidant, antibacterial, and antiparasitic activities. The present investigation is proposed to assess the antibacterial ability and free radical scavenging capacity against Gram-negative bacteria of various enzyme-digested skimmed camel's milk (SCM) using papain, pepsin, and trypsin at four hydrolysis times at one-hour intervals. An antibacterial growth assay was conducted to detect the antibacterial capacity of the generated various enzymatic camel milk hydrolysates. The results showed that the antibacterial activity of unhydrolyzed intact SCM displayed mild effect against *Escherichia coli* and no effect against *Salmonella enteritidis*. Pepsin-treated SCM hydrolysates after 3h hydrolysis significantly enhanced SCM antimicrobial activity against *E. coli*, and after 2h, hydrolysis produces SCM hydrolysates against *Salmonella enteritidis*. Pepsin and trypsin SCM-treated hydrolysates after 2h hydrolysis could completely abolish the survival growth rate of *Salmonella enteritidis*. Additionally, hydrolysates treated with papain SCM after 3 and 4 hours of hydrolysis showed a 100% ability to kill *Salmonella enteritidis*. Both DPPH method and ABTS assays were employed to evaluate antioxidants. Our findings demonstrated that all SCM hydrolysates gradually acquired strong antioxidant capacity after four hours of hydrolysis, with trypsin-treated hydrolysates having the highest values. The current work offered a great role of skimmed camel's milk hydrolysates and the possibilities for using these hydrolysates as food natural preservatives.

## 1. INTRODUCTION

Microbial contamination is a critical food safety issue as it leads to the spread of food poisoning. Many foodborne diseases and outbreaks are documented due to pathogens, viruses, and protozoa contamination (Bintsis, 2018). Bacterial foodborne illnesses continue to seriously impact public health, the global economy, and society. World Health Organization emphasizing the need to measure the burden and incidence of foodborne illnesses and promote policies that control and limit foodborne diseases through prioritizing food safety measures as a public health issue (WHO, 2015; Pires *et al.*, 2021).

Salmonellosis is one of the most reported foodborne diseases leading to gastroenteritis symptoms, abdominal cramps, headache, bloody diarrhea, myalgia, fever, nausea, and vomiting (Ehuwa *et al.*, 2021). Food ingestion items associated with salmonella spp, primarily from livestock such as chickens, pigs, egg products, and milk products, are associated with salmonellosis food poisoning. Among the avenues for transmission pathways are the bad hygienic practice as improper hand sanitation and exposure to infected animals (Munck *et al.*, 2020). According to cross

sectional study, showing high prevalence of *Salmonella enterica*, it has antibiotic resistance from food items, water and human (Sabeq *et al.*, 2022). Additionally, some pathogenic strains of *Escherichia coli*, considered serious foodborne pathogens, can infect the gastrointestinal tract and cause life-threatening sickness and acute diarrhea (Croxen *et al.*, 2013; Yang *et al.*, 2017). Therefore, novel natural antimicrobials are urgently needed to reduce bacterial infections in food. The ability of food's derived components to reduce the chance of human food poisoning is widely acknowledged. In place of synthetic chemical preservatives, functional foods, and nutraceuticals are now used, particularly in preventing foodborne illness and acting as natural preservatives (Kris-Etherton *et al.*, 2002).

Dietary proteins can modulate the body's physiological processes and act as good sources of amino acids. Some of the peptides that are encoded in original parent food protein sequences have acquired bioactive biological properties. These bioactive peptides serve as principles for nutraceuticals or functional foods because of their potential to have positive health effects (Chakrabartiet *et al.*, 2018). It is possible to designate various food products or components as functional smart foods or nutraceuticals with health

\* Correspondence to: dina.awad@fvvm.bu.edu.eg

benefits by adding various bioactive peptides derived from food (Hartmann and Meisel, 2007).

Many food proteins are hydrolyzed to produce bioactive peptides, which have a variety of biological effects, including antibacterial, antihypertensive, antioxidant, antiviral and antiparasitic activities. Several bioactive peptides can be generated from a variety of protein forms, such as meat, milk, and egg products, as well as marine, animal, and plant biomasses. Enzymatic hydrolysis is the most prevalent technique being researched for the production of bioactive peptides due to its generally recognized as safe (GRAS) (Marciniak *et al.*, 2018). Utilizing commercial enzymes derived from various sources is known as "enzymatic hydrolysis" (Toldrá *et al.*, 2020). Protein enzymatic hydrolysis uses mild temperature and pH settings, commercial selective enzymes, and a dearth of secondary by-products (Cruz-Casas *et al.*, 2021). Furthermore, the hydrolysis process using enzymes is quick and simple to inactivate the reaction, and if optimized, it can generate large quantities of high-quality bioactive peptides. Peptide structural characteristics, including amino acid arrangement, sequence, hydrophobicity, chain length, and net charge, play a significant impact in determining the peptide's biological activity (Maestri *et al.*, 2019).

Dromedary milk protein possesses a variety of bioactive qualities, including impacts on the immune system, antioxidant protection, cancer-fighting abilities, and antibacterial, antiviral, and hypoglycemic, ovidal capabilities (Gader and Alhaider, 2016; Behrouz *et al.*, 2022; Awad *et al.*, 2023). Additionally, camel's milk bioactive peptides availability for technologically functional uses in food systems has been recently increased (Al-Shamsi *et al.*, 2018).

Camel's milk proteins are diverse types of proteins with varied properties in structure and characteristics. They are divided into two main categories according to their solubility whey protein and caseins, which comprise 26.9% and 73.1% of total proteins, respectively (Lajnaf *et al.*, 2020). Recently, Camel milk hydrolysates derived from skimmed camel milk (SCM) has made by enzymatic procedures display more noticeable functional bioactive properties as compared to intact camel milk proteins (AliRedha *et al.*, 2022).

The antioxidant activity of camel milk proteins is due to their functional properties. Reactive oxygen species is a component of oxidative stress that has been related to several diseases, (Collins, 2005). Many research papers claimed camel milk has antioxidant properties, but these investigations were focused on bioactive components derived from fermented camel's milk products (Ayyash *et al.*, 2018).

So, there is an urgent need for more information on the comprehensive examination of camel protein hydrolysates' bioactive functional properties against various foodborne pathogens. In addition, it would be highly fruitful to assess camel milk protein hydrolysates from enzymes (papain, pepsin, and trypsin) for their antioxidant capacities and antibacterial against foodborne pathogens (*Salmonella enteritidis*, and *Escherichia coli*).

## 2. MATERIALS AND METHODS

### 2.1 Materials

Raw camel's milk (estimated 4.55% fat and 3.42% total protein) was procured from the open grazing land in Ras Sedr, South Sinai Governorate, Egypt. After collecting two litters from healthy female *Camelus dromedarius*, a local breed, samples were kept in sterile plastic bottles and on the ice bag till reach to the laboratory. Papain, pepsin, and

trypsin were received from Yanaye Bio-Technology Co., Ltd. (Shanghai, China). Mueller Hinton broth (MHB) and Tris-HCl buffer were purchased from Sigma-Aldrich (St. Louis, MO, USA). 2, 2'-azinobis (3- ethylbenzthiazoline-6-sulphonon acid) (ABTS) and 2, 2'-diphenyl-1-picrylhydrazyl (DPPH) were obtained from Sigma-Aldrich Chemical Co., India.

### 2.2 Skimmed camel milk hydrolysates

Raw camel's milk was skimmed (SCM) via centrifugation at 4700 x g for 10 min, at 4°C, using a refrigerated centrifuge (Thermo Scientific USA, central laboratory, faculty of veterinary medicine) till reached fat 1% after skimming. Enzymatic hydrolysis will be performed using digestive enzymes papain, pepsin, and trypsin at their optimal conditions at pH 6.0, 3.0, and 8.0, respectively, as shown in flow chart fig. (1). The protein substrate was dissolved in the appropriate buffer at 10 mg/ml. The enzyme had been dissolved at an enzyme-to-substrate ratio of 1:200 (w/w). After a 4-hour reaction, samples were heated in a boiling water bath for 10 minutes to inactivate the enzyme. The various SCM hydrolysis products were centrifuged at 4000 g for 15 minutes, and the supernatant was dialyzed, lyophilized, and then stored at 4 °C (Al-Shamsi *et al.*, 2018).




Fig. 1 Flowchart of skimmed camel milk hydrolysate

### 2.3 Antibacterial activity

Skimmed camel milk protein hydrolysates were tested for antibacterial activity as reported by Shavandi *et al.* (2017) and Esmaeilpour *et al.* (2017) method with slight modifications by bacterial count. Two indicator Gram-negative common pathogenic bacterial strains, *Salmonella enteritidis*, and *Escherichia coli*, were grown overnight in Mueller Hinton broth (MHB) medium at 37°C and working cultures were generated by diluting to roughly 4.0 log<sub>10</sub>cfu/ml using sterile MHB medium. The antimicrobial activity was assessed by monitoring the growth of indicator microorganisms in the presence or absence of 1000 µg/ml of different SCM hydrolysates after 2h of incubation at 37°C. The bacterial or colony forming unit (cfu) on nutrient agar was obtained by incubation at 37°C for 18 h. The bacterial growth rate is represented as log<sub>10</sub>cfu/ml, and killing power was also calculated. All assays were performed in triplicate. Killing power = (log<sub>10</sub> control - log<sub>10</sub>test).

### 2.4 Antioxidant activity

#### 2.4.1 DPPH radical scavenging activity

The radical scavenging activity against DPPH was carried out in accordance El-Hadary and Ramadan (2019) with some changes using specific dissolving buffer. Briefly, DPPH stock solution prepared (0.004% w/v) in ethyl alcohol was used for the assay. The ethanolic group of DPPH was used as a control, and samples of various skimmed milk hydrolysates produced from different enzyme digestion were dissolved in the alcohol. One milliliter of samples was mixed with DPPH solution (1ml). After vigorously shaking the mixture and storing it at room temperature for 30 minutes in the dark, the absorbance (Abs) of the DPPH radical was

measured by spectrophotometry at 517 nm (UV-1800 spectrophotometer, TOMOS, Italy).

According to the equation to measure the free radical scavenging activity.

$$\text{RSA \%} = \frac{[\text{Abs Blank} - \text{Abs Test}]}{\text{Abs Blank}} \times 100$$

#### 2.4.2 ABTS radical scavenging activity

Following the method outlined by Hernández-Ledesma et al. (2007), radical scavenging activity was measured against 2, 2'-azinobis (3-ethylbenzthiazoline-6-sulphonic acid) (ABTS). The stock was prepared and stored in the dark at room temperature for 16h before use. After that, ethanol was added to the solution to dilute it and produce a wavelength absorbance of 0.706 at 734 nm. Samples of various skim milk hydrolysates produced from different enzyme digestion were mixed well with (0.9 mL) of the ABTS solution and then measured at 734 nm after 15 minutes of incubation at room temperature. In the blank solution, distilled water was utilized in place of the sample. The reduction % was calculated using the reaction absorbance at 734 nm by the following equation:

$$\text{Inhibition \%} = \frac{[\text{Absorbance Control} - \text{Absorbance Sample}]}{\text{Absorbance Control}} \times 100$$

#### 2.5 Statistical analysis

The statistical analysis results were provided as mean value of three repeated trials, standard error ( $\pm$ ), and a P value of 0.05 was utilized for statistical evaluation using the MS Excel program (Spss, 2007).

### 3. RESULTS

The SCM antimicrobial efficacy and its different enzymatic hydrolysates, at various hydrolysis times, on the bacterial growth survival and inhibition rate against Gram-negative bacteria *E. coli* and *Salmonella enteritidis* were observed, and the mean values were recorded in  $\log_{10}$  cfu/ml as illustrated in figures (2-5) and table (1).

#### 3.1 Antibacterial activity of SCM against tested Gram-negative foodborne pathogens

Intact undigested SCM at a concentration of 1000  $\mu\text{g/ml}$  displayed minor antibacterial activity on *E. coli* as in fig. (2A), while intact undigested SCM showed no inhibitory antibacterial activity against *Salmonella enteritidis* in comparison to the control groups (Ctrl). As indicated in Figures (2 to 5A, B). It was observed that intact unhydrolyzed SCM could inhibit the *E. coli* growth rate to  $6.04 \pm 0.16 \log_{10}\text{cfu/ml}$ , whereas the Ctrl group reached  $8.48 \pm 0.17 \log_{10}\text{cfu/ml}$ , with killing power  $2.43 \pm 0.16 \log_{10}\text{cfu/ml}$  as showed in Fig. (5. A). On the other hand, in the case of *Salmonella enteritidis*, intact SCM showed antibacterial resistance with no killing capacity during the antibacterial assay against intact SCM, as demonstrated in Fig. (5. B).

#### 3.2 Antibacterial activity of papain camel milk hydrolysates

The effect of papain skim camel milk hydrolysates (HPaP) at different hydrolysis times (1-4h) at protein content 1000 $\mu\text{g/ml}$  against tested indicator Gram-negative bacteria

was shown in Fig. (2). We discovered the antibacterial efficacy of HPaP was hydrolysis time rate dependent as illustrated in fig. 2 (A, B). We observed that the inhibitory growth rate at the first hour of hydrolysis was  $6.41 \pm 0.19 \log_{10}\text{cfu/ml}$ , and it increased with increasing hydrolysis time rate at 4h with a mean value of  $4.76 \pm 0.17 \log_{10}\text{cfu/ml}$ . Furthermore, the hydrolysis time rate showed a great significant difference against *Salmonella enteritidis*, with skimmed camel milk displaying 100% killing power with a killing capacity of  $7.22 \pm 0.13 \log_{10}\text{cfu/ml}$  after enzymatic hydrolysis with papain (3 and 4h), as shown in Figs. (2B) and (5B).

#### 3.3 Antibacterial activity of pepsin camel milk hydrolysates

Figure (3A) has been showed the efficacy of pepsin hydrolysis concerning the antibacterial activity of SCM. We discovered pepsin hydrolysate (HPeP) after 3h hydrolysis performing a higher bactericidal activity against *E. coli* with a growth survival rate of  $3.31 \pm 0.16 \log_{10}\text{cfu/ml}$  in comparison to the Ctrl group, which reached  $8.48 \pm 0.17 \log_{10}\text{cfu/ml}$  during the antibacterial assay. Conversely, pepsin hydrolysate (HPeP) after 2h hydrolysis could completely abolish *Salmonella enteritidis* survival growth rate with killing power 100%, as illustrated in Fig. (3B).

#### 3.4 Antibacterial activity of trypsin camel milk hydrolysates

Figure (4) depicts the impact of trypsin hydrolysis for four hours (HTrY) on the antibacterial efficacy of skim camel milk. It was worth noting that HTrY, after 2h hydrolysis, acquired potent antibacterial activity compared to intact SCM and Ctrl group. HTrY after 2h hydrolysis could inhibit *E. coli* survival growth rate to  $5.07 \pm 0.16 \log_{10}\text{cfu/ml}$ . In contrast, it reached  $6.4 \pm 0.16$  and  $8.4 \pm 0.17 \log_{10}\text{cfu/ml}$  in SCM and Ctrl groups, respectively, as reported in fig. (4A). At the same time, HTrY after 2h hydrolysis could completely inhibit *Salmonella enteritidis* survival growth rate with 100% killing capacity as recorded in fig. (4B).

#### 3.5 Antioxidant activity of camel milk hydrolysates

The skim milk hydrolysates using different digestive enzymes (papain, pepsin, and trypsin) were estimated for the radical scavenging antioxidant activity using DPPH and ABTS assay (Fig. 6). It was observed that all various enzymatic digestion for SCM using papain, pepsin, and trypsin exhibited higher antioxidant activity than intact unhydrolyzed SCM. By increasing hydrolysis time, we found that the antioxidant activity gradually improved, as illustrated in Fig. (6, red arrow) in DPPH and ABTS assay. Skim camel milk hydrolysates generated by trypsin showed higher antioxidant properties in the DPPH and ABTS assays, followed by skim camel milk hydrolysates produced by papain and pepsin, respectively. As a result, the DPPH activity in HTrY after 1h hydrolysis increased 3 times the DPPH activity of intact SCM with mean values of 32.33% and 10.33%, respectively, and with prolonged hydrolysis rate; we found HTrY after 4h hydrolysis, we reported that DPPH radical scavenging activity enhanced six times the value of SCM antioxidant activity with mean 61.76%. Similarly, HTrY acquired the highest antioxidant activity in the ABTS assay than intact SCM.

Table (1) The antibacterial of camel milk hydrolysates using (papain, pepsin and trypsin) enzymes at different hydrolysis rates.

Strain/Enzyme Hydrolysis rate (hours)	<i>Escherichia coli</i>				<i>Salmonella enteritidis</i>			
	1h	2h	3h	4h	1h	2h	3h	4h
Papain hydrolysates (HPaP)	6.41 $\pm$ 0.19a	5.20 $\pm$ 0.19b	5.70 $\pm$ 0.16b	4.67 $\pm$ 0.17c	6.50 $\pm$ 0.20a	6.43 $\pm$ 0.23a	ND	ND
Pepsin hydrolysates (HPeP)	5.47 $\pm$ 0.16a	5.42 $\pm$ 0.12a	3.31 $\pm$ 0.06b	5.43 $\pm$ 0.12a	3.88 $\pm$ 0.07b	ND	3.54 $\pm$ 0.13a	3.41 $\pm$ 0.07a
Trypsin hydrolysates (HTrY)	5.53 $\pm$ 0.17b	5.07 $\pm$ 0.16b	6.71 $\pm$ 0.16a	6.20 $\pm$ 0.12a	3.65 $\pm$ 0.18c	ND	3.43 $\pm$ 0.17c	3.44 $\pm$ 0.18c

The antibacterial activity estimated against indicators Gram negative food borne pathogens were the camel milk hydrolysates at specific concentration (1000  $\mu\text{g/ml}$ ) incubated with indicators bacteria at 37  $^{\circ}\text{C}/2\text{h}$ . The control *E. coli* reach  $8.48 \pm 0.17$  and control *Sal. enteritidis* reach  $7.21 \pm 0.12$  after incubation. The antibacterial activity estimated as mean  $\log_{10}$  cfu/ml.

ND: Non detected, no bacterial colonies appear during the antibacterial assay.




Fig. (2) The antibacterial of skim camel milk hydrolysates using papain enzyme (HPaP) at different hydrolysis rate at specific concentration (1000 µg/ml). The antibacterial activity estimated against indicators Gram negative food borne pathogens where the skim camel milk hydrolysates by papain at specific concentration (1000 µg/ml) incubated with indicators bacteria at 37 °C/2h in comparison with intact skim camel milk (SCM) . The antibacterial activity is estimated as mean log<sub>10</sub> cfu/ml.




Fig. (3) The antibacterial of skim camel milk hydrolysates using pepsin enzyme (HPeP) at different hydrolysis rate at specific concentration (1000 µg/ml). The antibacterial activity estimated against indicators Gram negative food borne pathogens where the skim camel milk hydrolysates by pepsin at specific concentration (1000 µg/ml) incubated with indicators bacteria at 37 °C/2h in comparison with intact skim camel milk (SCM) . The antibacterial activity is estimated as mean log<sub>10</sub> cfu/ml.




Fig. (4) The antibacterial of skim camel milk hydrolysates using trypsin enzyme (HTrY) at different hydrolysis rate at specific concentration (1000 µg/ml). The antibacterial activity estimated against indicators Gram negative food borne pathogens where the skim camel milk hydrolysates by trypsin at specific concentration (1000 µg/ml) incubated with indicators bacteria at 37 °C/2h in comparison with intact skim camel milk (SCM) . The antibacterial activity is estimated as mean log<sub>10</sub> cfu/ml.




Fig. (5) The killing power of skim camel milk hydrolysates using different digestive enzymes enzyme (papain, pepsin, trypsin) at different hydrolysis rate at specific concentration (1000 µg/ml). The antibacterial activity estimated against indicators Gram negative food borne pathogens where the skim camel milk hydrolysates at specific concentration (1000 µg/ml) incubated with indicators bacteria at 37 °C/2h in comparison with intact skim camel milk (SCM) . The killing power activity estimated as mean log cfu<sub>10</sub>/ml.




Fig. (6) The antioxidant activity of skim camel milk hydrolysates using different digestive enzymes enzyme (papain, pepsin, trypsin) at different hydrolysis rate at specific concentration (1000 µg/ml). The antioxidant activity is estimated using DPPH assay and ABTS assay.

#### 4. DISCUSSION

Through consuming contaminated foods, pathogenic microorganisms, particularly bacteria, are the primary common cause of various food poisonings that carry a moderate to high risk of illness and a higher mortality rate (Fu *et al.*, 2016). Synthetic preservatives like synthetic chemicals and antimicrobials have largely been used to limit food deterioration and pathogenic bacteria (Calo *et al.*, 2015; Zheng *et al.*, 2013). However, these chemicals' use is restricted because of their unfavorable characteristics, such as teratogenicity, acute toxicity, and carcinogenicity (Calo *et al.*, 2015). Therefore, the current research aimed to find natural antibacterial compounds that might be extracted from camel milk and employed as natural preservatives. Compared to bovine milk, camel milk is characterized by more pronounced inhibitory systems (El Agamy *et al.*, 1992). Notably, dromedary milk contains twice as much Lysozyme and lactoferrin as bovine milk (Kappeler *et al.*, 1999). Additionally, Bioactive functional peptides and proteins included in camel's milk have various biological effects that could account for the antibacterial impact of skimmed camel milk against the growth rate of *E. coli* (Korhonen and Pihlanto, 2003)

On the contrary, the Gram-negative bacteria in the form of *Salmonella enteritidis*, which could enter the colon and translocate in the gastrointestinal epithelium (Eng *et al.*, 2015). *Escherichia coli* has a high pathogenicity level for the host it infects. The current results showed that intact SCM had slight antibacterial activity against *E. coli* and *Salmonella enteritidis*, which came in agreement with that of Benkerroum *et al.* (2004), who claimed that camel's milk has bacteriostatic properties against *E. coli* and *Listeria monocytogenes*. High levels of antimicrobial proteins, like Lysozyme, lactoferrin, lacto-peroxidase, and short peptidoglycan recognition protein, present in camel milk may cause this inhibitory effect (Benkerroum *et al.*, 2004). Although Gram-negative bacteria acquired an extra barrier outer membrane covering the peptidoglycan (Ramachandran, 2014).

The antibacterial activity and antioxidant capacity of the peptide fragments produced by the hydrolysis of dromedary skim milk proteins were evaluated in the current study using a variety of digestive enzymes. Generally, the various SCM hydrolysates exhibited more potent antibacterial activity than intact SCM, This result agreed with Kumar *et al.* (2016), who claimed that camel milk's antibacterial and antioxidant properties were enhanced by enzymatic hydrolysis.

Current result showed that intact SCM has no killing power, where the SCM generated hydrolysates at different hydrolysis times acquired potent antibacterial activity as in Fig. (5). It was suggested that the hydrophobicity was acquired for peptides generated from papain than from pepsin and trypsin at different hours of enzymatic hydrolysis. Due to the cationic charge of the produced SCM hydrophobic peptides and their initial contact with the microbial membrane, these peptides primarily target anionic microbial membranes (Teixeira *et al.*, 2012; Oñate-Garzón *et al.*, 2017). The hydrophobic amino acids are then introduced into the hydrophobic core of the membrane, causing the phospholipids to become more disordered and the barrier membrane function to be lost (Teixeira *et al.*, 2012). In order to achieve insertion into the microbial membrane, various peptide structural characteristics, such as hydrophobicity, amphipathicity, and secondary structure, are crucial (Lee *et al.*, 2016). Such structural qualities can alter the peptide's physicochemical properties and antibacterial effectiveness (Giangaspero *et al.*, 2001; Torrent *et al.*, 2011). During the 4h of hydrolysis, the antioxidant capacity was increased, indicating that SCM contains various cleavage and cutting sites for the used enzymes; the same increase was observed during the first 2h of hydrolysis (Oussaief *et al.*, 2020). As the hydrophobic areas buried within the parent protein are revealed during enzymatic proteolysis, alterations in protein globular shape may result. It is understood that more hydrophobic amino acids are exposed following proteolytic digestion due to the catalytic selectivity of the enzymes that cleave the peptide bonds produced by these amino acid residues. Due to enzyme differences specificity, the various digestive enzymes utilized in this study were able to have a variety of effects on the surface hydrophobicity, leading to diverse proteolytic patterns and the exposure of more hydrophobic amino groups, including phenylalanine, tyrosine, tryptophan, and leucine amino acids (Al-Shamsi *et al.*, 2018). Trypsin-treated SCM hydrolysates showed the highest antioxidant values, reflecting the highest degree of hydrophobicity, the main reason for higher antioxidant activity as in Fig. (6). However, pepsin-treated SCM after 3h hydrolysis could inhibit bacterial growth. Papain SCM hydrolysates could completely inhibit *Salmonella enteritidis* after 3 and 4h hydrolysis, pepsin SCM hydrolysates after 2h hydrolysis, and trypsin-treated SCM after 2h hydrolysis as in Fig. (6). Conversely, all SCM enzymatic-treated hydrolysates gradually acquired potent antioxidant activity with prolonged hydrolysis time as in Fig. (6). It was suggested for future studies to apply camel milk enzymatic hydrolysates generated by papain, pepsin, and trypsin as a natural preservative in food.

## 5. CONCLUSIONS

Most of the medicinal value of camel milk is attributed to its biological qualities, such as its antibacterial activity. In this study, after being in vitro hydrolyzed by papain, pepsin, and trypsin, skim camel's antibacterial activity was detected against *E. coli* and *Salmonella enteritidis*. These preliminary results confirmed the inhibitory activity of powerful camel's milk protein hydrolysates produced against various Gram-negative bacteria (*E. coli* and *Salmonella enteritidis*). The papain hydrolysates demonstrated pronounced antibacterial activity at different times of hydrolysis compared to that produced from pepsin and trypsin hydrolysis. The SCM hydrolysates showed a gradual increase in radical scavenging activity after 4 hours of hydrolysis. The results demonstrated that skimmed camel milk protein hydrolysates

can be tuned and manipulated to achieve specific functionalities, opening up a wide range of potential applications as new antimicrobial agents in food. Therefore, additional research is required to identify the antibacterial peptides and explore the potential for their use as natural preservative in food models.

## 6. REFERENCES

1. El Agamy, E.I., Ruppner, R., Ismail, A., Champagne, C.P. and Assaf, R. (1992) Antibacterial and antiviral activity of camel milk protective proteins. *The Journal of dairy research*. 59 (2), 169–175.
2. Al-Shamsi, K.A., Mudgil, P., Hassan, H.M. and Maqsood, S. (2018) Camel milk protein hydrolysates with improved technofunctional properties and enhanced antioxidant potential in in vitro and in food model systems. *Journal of dairy science*. 101 (1), 47–60.
3. Ali Redha, A., Valizadenia, H., Siddiqui, S.A. and Maqsood, S. (2022) A state-of-art review on camel milk proteins as an emerging source of bioactive peptides with diverse nutraceutical properties. *Food Chemistry*. 373, 131444.
4. Awad, D.A.B., Eldiarby, A.S., Abdallah, M., Hamad, A. and Gawad, S.M.A. (2023) Brief report first report of the in vitro ovidical activity of camel milk and its fractions on zoonotic-liver fluke (*Fasciola gigantica*) eggs. *Veterinary Research Communications*. doi:10.1007/s11259-023-10144-8.
5. Ayyash, M., Al-Nuaimi, A.K., Al-Mahadin, S. and Liu, S.-Q. (2018) In vitro investigation of anticancer and ACE-inhibiting activity,  $\alpha$ -amylase and  $\alpha$ -glucosidase inhibition, and antioxidant activity of camel milk fermented with camel milk probiotic: A comparative study with fermented bovine milk. *Food Chemistry*. 239, 588–597.
6. Behrouz, S., Saadat, S., Memarzia, A., Sarir, H., Folkerts, G. and Boskabady, M.H. (2022) The Antioxidant, Anti-Inflammatory and Immunomodulatory Effects of Camel Milk. *Front Immunol*. 12:13:855342. doi: 10.3389/fimmu.2022.855342.
7. Benkerroum, N., Mekkaoui, M., Bennani, N. and Hidane, K. (2004) Antimicrobial activity of camel's milk against pathogenic strains of *Escherichia coli* and *Listeria monocytogenes*. *International Journal of Dairy Technology*. 57, 39–43.
8. Bintsis, T. (2018) Microbial pollution and food safety. *AIMS microbiology*. 4 (3), 377–396. doi:10.3934/microbiol.2018.3.377.
9. Calo, J.R., Crandall, P.G., O'Bryan, C.A. and Ricke, S.C. (2015) Essential oils as antimicrobials in food systems – A review. *Food Control*. 54, 111–119.
10. Chakrabarti, S., Guha, S. and Majumder, K. (2018) Food-Derived Bioactive Peptides in Human Health: Challenges and Opportunities. *Nutrients*. 10 (11). doi:10.3390/nu10111738.
11. Collins, A.R. (2005) Antioxidant intervention as a route to cancer prevention. *European Journal of Cancer*. 41 (13), 1923–1930.
12. Croxen, M.A., Law, R.J., Scholz, R., Keeney, K.M., Wlodarska, M. and Finlay, B.B. (2013) Recent advances in understanding enteric pathogenic *Escherichia coli*. *Clinical microbiology reviews*. 26 (4), 822–880.
13. Ehuwa, O., Jaiswal, A.K. and Jaiswal, S. (2021) Salmonella, Food Safety and Food Handling Practices. *Foods*. 10 (5). doi:10.3390/foods10050907.
14. El-Hadary, A.E. and Ramadan, M.F. (2019) Phenolic profiles, antihyperglycemic, antihyperlipidemic, and antioxidant properties of pomegranate (*Punica granatum*) peel extract. *Journal of food biochemistry*. 43 (4), e12803.
15. Eng, S.-K., Pusparajah, P., Ab Mutalib, N.-S., Ser, H.-L., Chan, K.-G. and Lee, L.-H. (2015) Salmonella: A review on pathogenesis, epidemiology and antibiotic resistance. *Frontiers in Life Science*. 8 (3), 284–293.
16. Esmailpour, M., Ehsani, M., Aminlari, M., Shekarfroush, S. and Hoseini, E. (2017) Antimicrobial Peptides Derived from Goat's Milk Whey Proteins Obtained by Enzymatic Hydrolysis. 65–72.
17. Fu, Y., Sarkar, P., Bhunia, A.K. and Yao, Y. (2016) Delivery systems of antimicrobial compounds to food. *Trends in Food Science and Technology*. 57, 165–177.
18. Gader, A.G.M.A. and Alhaider, A.A. (2016) The unique medicinal properties of camel products: A review of the scientific

- evidence. Journal of taibah university medical sciences. 11 (2), 98–103.
19. Giangaspero, A., Sandri, L. and Tossi, A. (2001) Amphipathic alpha helical antimicrobial peptides. European journal of biochemistry. 268 (21), 5589–5600.
  20. Hartmann, R. and Meisel, H. (2007) Food-derived peptides with biological activity: from research to food applications. Current Opinion in Biotechnology. 18 (2), 163–169.
  21. Hernández-Ledesma, B., Quirós, A., Amigo, L. and Recio, I. (2007) Identification of bioactive peptides after digestion of human milk and infant formula with pepsin and pancreatin. International Dairy Journal. 17 (1), 42–49.
  22. Kappeler, S., Farah, Z. and Puhan, Z. (1999) Alternative Splicing of Lactophorin mRNA from Lactating Mammary Gland of the Camel (*Camelus dromedarius*). Journal of Dairy Science. 82 (10), 2084–2093.
  23. Korhonen, H. and Pihlanto, A. (2003) Food-derived bioactive peptides--opportunities for designing future foods. Current pharmaceutical design. 9 (16), 1297–1308.
  24. Kris-Etherton, P.M., Hecker, K.D., Bonanome, A., Coval, S.M., Binkoski, A.E., Hilpert, K.F., Griel, A.E. and Etherton, T.D. (2002) Bioactive compounds in foods: their role in the prevention of cardiovascular disease and cancer. The American journal of medicine. 113 Suppl, 71S-88S.
  25. Kumar, D., Chatli, M.K., Singh, R., Mehta, N. and Kumar, P. (2016) Antioxidant and antimicrobial activity of camel milk casein hydrolysates and its fractions. Small Ruminant Research. 139, 20–25.
  26. Lajnaf, R., Zouari, A., Trigui, I., Attia, H. and Ayadi, M.A. (2020) Effect of different heating temperatures on foaming properties of camel milk proteins: A comparison with bovine milk proteins. International Dairy Journal. 104, 104643.
  27. Lee, T.-H., Hall, K.N. and Aguilar, M.-I. (2016) Antimicrobial Peptide Structure and Mechanism of Action: A Focus on the Role of Membrane Structure. Current topics in medicinal chemistry. 16 (1), 25–39.
  28. Maestri, E., Pavlicevic, M., Montorsi, M. and Marmiroli, N. (2019) Meta-Analysis for Correlating Structure of Bioactive Peptides in Foods of Animal Origin with Regard to Effect and Stability. Comprehensive Reviews in Food Science and Food Safety. 18 (1), 3–30.
  29. Marciniak, A., Suwal, S., Naderi, N., Pouliot, Y. and Doyen, A. (2018) Enhancing enzymatic hydrolysis of food proteins and production of bioactive peptides using high hydrostatic pressure technology. Trends in Food Science and Technology. 80, 187–198.
  30. Munck, N., Smith, J., Bates, J., Glass, K., Hald, T. and Kirk, M.D. (2020) Source Attribution of *Salmonella* in Macadamia Nuts to Animal and Environmental Reservoirs in Queensland, Australia. Foodborne pathogens and disease. 17 (5), 357–364.
  31. Oñate-Garzón, J., Ausili, A., Manrique-Moreno, M., Torrecillas, A., Aranda, F.J., Patiño, E. and Gomez-Fernández, J.C. (2017) The increase in positively charged residues in cecropin D-like *Galleria mellonella* favors its interaction with membrane models that imitate bacterial membranes. Archives of biochemistry and biophysics. 629, 54–62.
  32. Oussaief, O., Jrad, Z., Adt, I., Khorchani, T. and El-Hatmi, H. (2020) Dromedary Milk Protein Hydrolysates Show Enhanced Antioxidant and Functional Properties. Food technology and biotechnology. 58 (2), 147–158.
  33. Pires, S.M., Desta, B.N., Mughini-Gras, L., Mmbaga, B.T., Fayemi, O.E., Salvador, E.M., Gobena, T., Majowicz, S.E., Hald, T., Hoejskov, P.S., Minato, Y. and Devleeschauwer, B. (2021) Burden of foodborne diseases: think global, act local. Current Opinion in Food Science. 39, 152–159.
  34. Ramachandran, G. (2014) Gram-positive and gram-negative bacterial toxins in sepsis: a brief review. Virulence. 5 (1), 213–218.
  35. Sabeq, I., Awad, D., Hamad, A., Nabil, M., Aboubakr, M., Abaza, M., Fouad, M., Hussein, A., Shama, S., Ramadan, H. and Edris, S. (2022) Prevalence and molecular characterization of foodborne and human-derived *Salmonella* strains for resistance to critically important antibiotics. Transboundary and Emerging Diseases. 69 (5), e2153–e2163.
  36. Shavandi, A., Hu, Z., Teh, S., Zhao, J., Carne, A., Bekhit, A. and Bekhit, A.E.-D.A. (2017) Antioxidant and functional properties of protein hydrolysates obtained from squid pen chitosan extraction effluent. Food Chemistry. 227, 194–201.
  37. Spss, I. (2007). SPSS version 16.0. Chicago, 649 IL: SPSS Incorporated. (n.d.)
  38. Teixeira, V., Feio, M.J. and Bastos, M. (2012) Role of lipids in the interaction of antimicrobial peptides with membranes. Progress in lipid research. 51 (2), 149–177.
  39. Torrent, M., Andreu, D., Nogués, V.M. and Boix, E. (2011) Connecting peptide physicochemical and antimicrobial properties by a rational prediction model. PloS one. 6 (2), e16968.
  40. WHO, W. (2015) WHO estimates of the global burden of foodborne diseases: foodborne disease burden epidemiology reference group 2007–2015. Geneva, Switzerland.
  41. Yang, S.-C., Lin, C.-H., Aljuffali, I.A. and Fang, J.-Y. (2017) Current pathogenic *Escherichia coli* foodborne outbreak cases and therapy development. Archives of Microbiology. 199 (6), 811–825.
  42. Zheng, L., Bae, Y.-M., Jung, K.-S., Heu, S. and Lee, S.-Y. (2013) Antimicrobial activity of natural antimicrobial substances against spoilage bacteria isolated from fresh produce. Food Control. 32 (2), 665–672.