



Aflatoxins in Cereals: Prevalence, Health Implications and Management Strategies: A Review

Yusuf O. Oyebamiji⁽¹⁾, Noraziyah A. Shamsudin^{(1)#}, Ismail A. Adebayo⁽²⁾, Olayinka B. Umar⁽³⁾, Nurul A. Mohd Zaini⁽⁴⁾, Mohd N. Ismail⁽⁵⁾

⁽¹⁾Department of Biological Sciences and Biotechnology, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia;

⁽²⁾Department of Microbiology, Parasitology and Immunology, School of Medicine, Kabale University, Kabale, Uganda; ⁽³⁾Department of Plant Biology, Faculty of Life Sciences, University of Ilorin, Ilorin, Kwara State, Nigeria; ⁽⁴⁾Department of Food Sciences, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia; ⁽⁵⁾Analytical Biochemistry Research Centre, Universiti Sains Malaysia (USM), 11800, Pulau Pinang, Malaysia.

⁽¹⁾Department of Biological Sciences and Biotechnology, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia; ⁽²⁾Department of Microbiology, Parasitology and Immunology, School of Medicine, Kabale University, Kabale, Uganda; ⁽³⁾Department of Plant Biology, Faculty of Life Sciences, University of Ilorin, Ilorin, Kwara State, Nigeria; ⁽⁴⁾Department of Food Sciences, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia; ⁽⁵⁾Analytical Biochemistry Research Centre, Universiti Sains Malaysia (USM), 11800, Pulau Pinang, Malaysia.



CEREALS are consumed globally because of their nutritional values and potential to reduce malnutrition. In spite of their usefulness, cereals are prone to aflatoxin contamination. Aflatoxins are highly toxic and carcinogenic secondary metabolic products that contaminate agricultural products consumed by humans. Studies have shown that aflatoxins are found in cereals at a high level. Human exposure to aflatoxins through food and feed results in a wide range of health issues, including a weakened immune system and cancer. Worst, it can cause death depending on the level and extent of exposure. Several climate-induced factors, such as drought, can trigger aflatoxin production worldwide, especially in Africa, where the environment is conducive. Several precautions have been taken to mitigate human exposure to aflatoxins, including strict regulations, pre- and post-harvest contamination prevention, detoxification, and decontamination. In addition, good farm management and practices, and awareness creation and education can help to reduce aflatoxin contamination and exposure, if not eradicating it. The current review detailed the up-to-date information on aflatoxin occurrence, health implications, and control measures for aflatoxins in cereals to ensure food safety and human well-being. This review also illuminated the potential hazards of human exposure to aflatoxins beyond the permissible level, particularly in children.

Keywords: Aflatoxins, Carcinogenic, Food contamination, Health complications, Malnutrition.

Introduction

Cereals, such as maize, have gained popularity worldwide due to their nutritional value and potential to reduce malnutrition (Temba et al., 2017). Currently, the annual cereal production globally is pegged at 1513.0 million tons, which is below the expectations by 3 million tons based on the projected output (Anonymous, 2021). The 20th century has seen a massive rise in the global population due to an increased birth rate from 1.65 to 6 billions (Kobayashi et al., 2020). The increase in cereal demand and consumption has doubled due to the rapidly escalating population.

As the production capacity increases, proper and adequate resource management at various processing and production stages is required (Nazir et al., 2019). However, this process is undermined due to the susceptibility of cereals to aflatoxins contamination, thus threatening food safety.

Aflatoxin groups are highly toxic secondary metabolites produced by *Aspergillus* spp., principally *A. flavus*, *A. parasiticus*, and *A. nomius* (Aasa et al., 2023). They are derived chemically from difuranocoumarin with a lactone ring or a pentanone ring, and a coumarin nucleus-

#Corresponding author email: nora_aziz@ukm.edu.my;

Tel.: +6017311527672

Received 26/02/2023; Accepted 06/11/2023

DOI: 10.21608/ejbo.2023.196133.2265

Edited by: Prof. Dr. Neveen M. Khalil, Faculty of Science, Cairo University, Giza, Egypt.

©2024 National Information and Documentation Center (NIDOC)

based bifuran group (Nazhand et al., 2020; Roy et al., 2023). Aflatoxins are compounds that have mutagenic, estrogenic, and immunosuppressive effects on the well-being of humankind and animals (Peles et al., 2021). The most common types of aflatoxins include B1 (AFB1), B2 (AFB2), G1 (AFG1), and G2 (AFG2), among others (Aasa et al., 2023). These types of aflatoxins can intoxicate the human body system via several pathways, such as the cutaneous layers, mucous layers, and respiratory tract, subsequently promoting the activation of an inflammatory response. Aflatoxin M1 (AFM1) and M2 (AFM2) are found in milk and are the hydroxylated metabolites of AFB1 and AFB2. *A. flavus* produces AFB whereas *A. parasiticus* produces both AFB and AFG. The toxic nature of the common types of aflatoxins decreases in the order of B1, G1, B2, and G2 (Kumar et al., 2017; Wanniarachchi et al., 2023). AFB1 is considered the most poisonous of all mycotoxins, with a lethal dose (LD50) of 0.36 mg/kg, placing it among the highly hazardous group of toxins (Ndagijimana et al., 2020). Based on experimental results, AFB1 toxicity was ten times higher than potassium cyanide, sixty-eight times higher than arsenic, and seventy times more carcinogenic than dimethylnitrosamine (Yan et al., 2020). The International Agency for Research on Cancer (IARC) has categorized AFB1 as a group 1 carcinogen ("carcinogenic to humans") owing to its genotoxic and carcinogenic effects (Jallow et al., 2021; Zhang et al., 2023).

Aflatoxins exist everywhere in nature. Factors, such as high humidity, poor harvesting, climatic change, and storage methods trigger aflatoxin development in different regions. Compounded by these factors is the policymakers' inappropriate enforcement of aflatoxin safe limit, contributing to several outbreaks, especially in developing nations (Yard et al., 2013; Somda et al., 2023; Wanniarachchi et al., 2023). Aflatoxin contamination is common in cereals, dry fruits, spices, tree nuts, cottonseed, and cowpea, among other crops (Awuchi et al., 2021; Somda et al., 2023). Contamination of crops can occur before and after harvest. The occurrence of aflatoxins at the pre-harvest stage is not common compared to the storage stage because the associated fungi are frequently regarded as storage molds. Grain deterioration at the post-harvest stage is attributed to *A. flavus* and *A. parasiticus* contamination, while *A. flavus* is the primary fungus that contaminates crops in the field (Liu et al., 2006). Due to the heat resistance of these

fungi, the current food processing methods are insufficient to eliminate aflatoxins from contaminated agricultural commodities. The consumption of human aflatoxin-contaminated products has resulted in severe health problems and complications (Mahato et al., 2019), such as malaise, fever, anorexia, vomiting, acute hepatitis and liver problems, and in severe cases, death (Udomkun et al., 2017).

To control the amount of aflatoxin intake in food, the Codex Alimentarius Commission (an intergovernmental agency) created by the World Health Organization (WHO) in cooperation with the Food and Agriculture Organization (FAO) aims to safeguard consumer health and enhance trade by developing an international standard for food and feed (Sirma et al., 2018). Moreover, in 2007, the European Union (EU) has set a safe limit for total aflatoxins and AFB1 at 4g/kg and 2g/kg, respectively, for human intake. In 2010, the limit was revised and set at 5g/kg and 10g/kg for AFB1 and total aflatoxin, respectively. The United States and Canada set their limits at 20g/kg and 15g/kg, respectively. Bosnia and Herzegovina and Switzerland have the lowest limit for AFB1, which is 1g/kg (Ali, 2019). In Japan, any concentration of aflatoxin in crops is prohibited (Dadzie et al., 2019). In Malaysia, based on the 1985 Malaysia Food Regulation, the maximum limit of all mycotoxins was set at 35µg/kg initially. Later, it was reviewed and set at 5 g/kg for all mycotoxins, including aflatoxin (Sabran et al., 2013). Egypt set its limit at 4 to 15g/kg and 2 to 12g/kg for total aflatoxin and AFB1, respectively, while 0.01g/kg was set as the limit for AFB1 in processed cereal-based foods and baby foods for infants and young children (Marrez & Ayeshe, 2022). Furthermore, the Food and Agriculture Organization (FAO) has also set the safe limit for AFB1 to be 1–20g/kg in food (Ndagijimana et al., 2020).

Aflatoxin occurrence in cereals

Cereals, such as corn, rice, and wheat are vulnerable to aflatoxin contamination by aflatoxigenic fungi. According to the FAO survey, mycotoxins contaminate approximately one-quarter of the world's cereal crops (Jallow et al., 2021; Rudraboyina et al., 2023). A previous study revealed that out of 18,097 cereal samples tested for aflatoxin contamination, 36.7% of the samples were contaminated with a form of aflatoxin (Andrade & Caldas, 2015). The occurrence of aflatoxins in grains is not restricted to a particular geographical or climatic zone. It is prevalent in

tropical, subtropical, and a few temperate regions with conducive environmental factors that favor its production. Besides, these regions prone to aflatoxin contamination, the Mediterranean zones are now vulnerable to aflatoxin contamination due to a shift in local occurrence areas of AFs driven by the main changes in climate variables, consequently bringing about a rapid

increase in aflatoxin contamination of cereals globally (Mahato et al., 2019). Furthermore, most developing countries' suitable social and environmental conditions have triggered and aided the prevalence of mycotoxins in agricultural commodities. Table 1 shows the occurrence of aflatoxins in cereals in different countries worldwide.

TABLE 1. Worldwide occurrence of aflatoxins in cereals

Country	Food Matrix	Type of Aflatoxin	Range (µg/kg)	References
Ghana	Cereals based product	AFB1	0.18–23.27	Blankson et al. (2019)
Ivory Coast	Maize	Total Aflatoxin	0.3-173	Aasa et al. (2023)
Zambia	Maize	AFB1	16	Thakur et al. (2022)
Zimbabwe	Maize	AFB1	0.5 -26.6	Murashiki et al. (2017)
Vietnam	Maize	AFB1	1-34.8	Lee et al. (2017)
India	Maize	AFB1	62	Thakur et al. (2022)
Togo	Maize	AFB1	Max 256	Hanvi et al. (2019)
China	Maize	AFB1	4.4	Zhao et al. (2021)
Vietnam	Maize	AFB1	>5	Nguyen et al. (2018)
Pakistan	Rice Noodles	AFB1	3.60	Iqbal et al. (2016)
Indonesia	Rice products	AFB1	2.0 to 7	Ali (2019)
Korea	Rice	AFB1	1.8-7.3	Ali (2019)
China	Rice bran	AFB1	7.5	Zhao et al. (2021)
Ivory Coast	Rice	Total Aflatoxin	4.1	Aasa et al. (2023)
China	Rice	AFB1	0.1–136.80	Mahato et al. (2019)
Iran	Rice Flour	AFB1	0.46-10.16	Mottaghianpour et al. (2021)
Saudi Arabia	Rice	Total Aflatoxin	0.07-7.09	Elzupir et al. (2018)
Togo	Sorghum	AFB1	6–16	Hanvi et al. (2019)
Uganda	Sorghum	Total Aflatoxin	11.8 ± 1.8	Echodu et al. (2019)
Tunisia	Sorghum	Total Aflatoxin	0.4-25.8	Filazi & Tansel Sireli (2013)
Turkey	Wheat	AFB1	0.21–0.35	Turksoy & Kabak (2020)
Malaysia	Wheat	AFB1	0.55–5.07	Mahato et al. (2019)
Iran	Wheat	AFB1	Max 7.08	Mohadesch et al. (2016)
Algeria	Wheat	AFB1	0.22–13.96	Riba et al. (2010)
Lebanon	Wheat	AFB1	1.05 - 7.36	Joubrane et al. (2020)
Iran	Wheat Flour	AFB1	0.1-0.26	Mottaghianpour et al. (2021)

Rice

Globally, rice is one of the most important sources of calories for more than 50% of the world's population. Asian countries constitute the major rice producers and consumers; approximately 600 million tons are produced yearly on more than 150 million hectares of land (Mohd Ikmal et al., 2019). Rice cultivation is typically done in humid and warm climatic conditions (Lai et al., 2015). When stored under poor conditions, rice is easily contaminated by mycotoxin-producing fungi. Harvested rice in waterlogged areas and with high moisture levels is vulnerable to mold infection and mycotoxin contamination (Majeed et al., 2018). In certain parts of India, frequent and heavy waterfalls are witnessed during harvest, causing the crop to become damp and vulnerable to fungal attacks. Furthermore, farmers cannot sun-dry the grains during this period, which is the norm. As a result, the grain moisture content remains unchanged. Thus, transporting grains with a moisture content higher than the average level (>14%) into the storage system makes them susceptible to fungal attack. The detrimental consequences of such contamination include reduced seed quality, grain discoloration or husk, and losses in viability (Reddy et al., 2009). Several nations worldwide, including Pakistan, Brazil, Indonesia, China, India, Korea, and Austria, have reported the presence of aflatoxin in rice samples. A study in Pakistan showed that 72 out of 208 samples were contaminated with AFB1 at a concentration of 3.60g/kg (Iqbal et al., 2016). In Canada, AFB1 was reported in imported rice from USA and Asia, with the mean concentration ranging between 0.34g/kg and 0.39g/kg (Ndagijimana et al., 2020). A recent study conducted in Iran showed that all 24 samples of rice flour analyzed tested positive for AFB1, with the concentration ranging between 0.46g/kg and 10.16g/kg (Mottaghianpour et al., 2021). Despite the average aflatoxin contamination level being within the safe limit, a previous study indicated a positive correlation between daily consumption of aflatoxin-contaminated rice and hepatocellular carcinoma (HCC) incidences in some rice-consuming countries in Asia. Chronic exposure to aflatoxin explains this relationship (Elzupir et al., 2018).

Maize

Maize is considered a staple food by the world population. The United States is regarded as the world's largest producer of maize, with 370 million metric tons in 2017 while countries, such

as Ethiopia, Brazil, China, and the EU combined produce 436 million metric tons, making a total of more than 807 million metric tons of maize by the world's major maize producers (Block et al., 2018). Despite maize's global impact, the natural enemy's existence and invasion of maize have considerably impacted its production. Insects and other herbivores' invasion results in nearly 6 to 19% grain damage, with pathogen attacks, such as fungus, causing an additional 10% damage (Di Domenico et al., 2016). One of the significant challenges encountered by corn at the storage stage is fungal attack and contamination. This can be attributed to corn's high starch content, which serves as a good substrate, thus making it vulnerable to contamination by fungi, especially *Aspergillus* spp., and *Fusarium* spp. (Di Domenico et al., 2016). Maize can be contaminated with airborne fungi and mycotoxins at pre- and post-harvest stages. In maize, aflatoxin occurs during pre-harvest activities, such as cultivation (i.e., in the field); the harvesting period; and post-harvest stages, such as transportation, storage, and processing. Poor farm management during cultivation allows aflatoxin to enter the post-harvest stage. Inadequate and poor storage conditions result in fungal attacks and aflatoxin contamination (Singh et al., 2019). Other variables, including humidity and temperature, trigger fungal growth in grains (Mtega et al., 2020).

According to a previous study, maize was considered as one of the crops mainly contaminated by aflatoxin in sub-Saharan Africa. Moreover, other studies found that the level of aflatoxin in contaminated crops, such as maize was unacceptable, with samples having up to 1000ppb of aflatoxins (Nakavuma et al., 2020). Due to their socio-economic significance, investigations have been conducted worldwide to determine the occurrence and extent of aflatoxins in maize samples. A study in Vietnam revealed that 204 and 141 maize samples tested positive for AFB1 out of 378 samples at concentrations of >5 g/kg and >20g/kg, respectively (Nguyen et al., 2018). In South Africa, AFB1 was reported in six maize samples out of 29, with the concentration ranging from 1 to 149g/kg (Mngqawaa et al., 2015). The prevalence of aflatoxins has caused a considerable reduction in net yield and economic profits. For example, the total money spent on reducing aflatoxins in the US is around \$500 million yearly through aflatoxin on maize and other grown crops, in addition to that spent on animal health

maintenance (which is a small fraction of the cost) (Wu, 2015). Similarly, it was estimated that \$163 million was lost annually on average to US maize growers through the prevention of aflatoxin infection (Wu, 2015). Yearly, approximately \$1.2 billion is lost due to contaminated agricultural products, of which African countries account for 38% of the losses (\$450 million) (Gbashi et al., 2018).

Sorghum

Sorghum is one of the essential grains consumed by the world's populace. In sub-Saharan Africa, it is considered the second most cultivated and important grain after maize. Globally, it is fifth based on consumption and is significantly behind wheat, maize, rice, and barley. Based on 2013 data, 61.5 million metric tons were produced and cultivated on approximately 42.3 million hectares (Garba, 2019). Despite these huge global production capacities, mycotoxin production threatens sorghum production. The presence of mycotoxins in sorghum has been proven, thus making it one of the major sources of mycotoxin exposure for animals and humans (Garba, 2019). Contaminants of sorghum grains include fungal genera, occurring during the panicle and grain developmental stages. The most common grain mold pathogens that contaminate sorghum include *Aspergillus* spp., *Alternaria* spp., *Fusarium* spp., *Cladosporium* spp., *Curvularia* spp., and *Penicillium* spp. (Lahouar et al., 2016). In the first comprehensive study of aflatoxin contamination in sorghum, 70% of the samples exhibited aflatoxin levels greater than 10ppb. The great amount of aflatoxin level in the sorghum samples can be attributed to improper agricultural management, such as poor storage and processing techniques because the production of sorghum is mostly carried out at the subsistence level, at which there is no standard mechanism to inspect and regulate the quality of the produce (Lukwago et al., 2019). Other studies have also reported the presence of aflatoxin in sorghum. For instance, in Togo, AFB1 was reported in three sorghum samples out of 12, with a concentration between 6 g/kg and 16g/kg (Hanvi et al., 2019).

Climate-Induced factors affecting aflatoxin production

The frequent changes in climatic conditions are considered the main factor significantly contributing to food insecurity worldwide. Therefore, there is considerable concern about the possible implication of environmental

changes on the existence of mycotoxins in agricultural commodities (Battilani et al., 2016). Fungal proliferation and mycotoxin secretion can occur at any developmental stage during the plant life cycle depending on environmental factors, such as rainfall, temperature, humidity, and agricultural management methods. Plant immunocompromising factors, which include injury, water stress, poor fertilization, and pest infestation, are recognized enablers of aflatoxin progress in agricultural commodities (Jallow et al., 2021). These variables impact mycotoxigenic fungal occurrence, survival, frequency, distribution, and subsequent toxin accumulation (Daou et al., 2021).

Temperature, humidity, and water activity

Temperature and humidity have tremendous impacts on which fungi that attack crops, with warm climates favoring aflatoxin production (Petrović et al., 2023). The infection of crops by fungi has brought about a drastic decrease in the yield of agricultural commodities, especially cereals and their derivatives. Variation in the temperature of the immediate environment significantly impacts aflatoxin production and the levels of expression of regulatory genes (*aflR* and *aflS*). Previous research reported a strong relationship between an early structural gene (*aflD*) expression and AFB1. Temperature reacts with water activity (a_w) and affects the ratio of regulatory genes (*aflR/aflS*), which is directly proportional to the production of AFB1. *Aspergillus* spp. and the production of aflatoxin are greatly influenced by the interaction between temperature and water activity (a_w) (Kumar et al., 2017). Documentation from previous studies has shown that environmental stresses, including low a_w and temperature, play a crucial role in the regulation of *A. flavus* growth and trigger aflatoxin production. Aflatoxins' production in several grains differs in terms of sensitivity to a_w and temperature (Tai et al., 2020). Based on a previous study Mousa et al. (2013), the rapid growth of *A. flavus* and aflatoxin production were observed in brown rice between 25°C and 35°C at 0.82 a_w , but not in polished rice in a similar condition. However, brown and polished rice showed increased aflatoxin production at maximum a_w values (0.90 to 0.92) at a temperature of 20°C after 21 days of incubation). Even though it was recorded that progressive toxin production happened at 25 to 30°C within a wider range of 0.82 a_w values, it could be concluded that polished rice did not

seem to allow *A. flavus* growth and aflatoxin production as compared to brown rice. Another study done on the effects of environmental factors on aflatoxin production indicated that the growth of *A. parasiticus*, *A. flavus*, and *A. oryzae* was observed at 25 °C in 0.82 a_w and 0.81 a_w at 30°C and 37°C. The study did not assess the relationship between aflatoxin production and *A. flavus* and *A. parasiticus* (Milani, 2013). A study by Lv et al. (2019) revealed that optimal growth of fungi was observed between 28°C and 37°C with an a_w value of 0.92 to 0.96. A study by Battilani et al. (2016) predicted the occurrence of AFB1 in cereals was due to the changes in climatic conditions in European countries; for every 2°C rise in temperature, there is an increase in aflatoxin risk in countries such as Italy, Greece, Turkey, Cyprus, Spain, Bulgaria, and Albania. Naturally, the control of ambient temperature is impossible, but its implications on the later stages of plant development can be avoided by early planting. For instance, in North Carolina, maize planted in April was reported to experience low aflatoxin contamination compared to maize grown in May (Abbas et al., 2009).

Drought

Drought incidence during crop production makes them vulnerable to diseases and fungal attacks, increasing aflatoxin contamination and reducing grain yield, fertility, and livestock production. The 2016 El-Nino drought events experienced in South Africa caused losses of several million dollars, and it was identified as the hottest year in nearly a century (Gbashi et al., 2018). Yearly, climate change causes a significant economic loss of about US\$ 1.68 billion to the U.S. maize sector (Thakur, et al., 2022). In 2018 and 2019, a severe drought hit central Europe, with the continent experiencing all-time high summer temperatures. This trend is expected to continue and occur more often if greenhouse gas emissions increase, significantly impacting agricultural productivity (Valencia-Quintana et al., 2020). Water deficits induced by climate change weaken plant systems. It makes them more vulnerable to fungal attacks and aflatoxin production, which could be triggered if the crop is cultivated in the rainy season (Benkerroum, 2020). Water stress triggers aflatoxin biosynthesis, as witnessed in tropical nations, where acute aflatoxicosis cases are frequently reported upon ingesting contaminated crops (Valencia-Quintana et al., 2020). A study conducted in South Africa as early as 1965

established the relationship between drought and high aflatoxin contamination. This association was also reported in studies conducted on agricultural products in the US and Nigeria (Sanders et al., 1993). Moreover, research has been performed extensively to determine the linkage between pre-harvest aflatoxin contamination and drought. For instance, studies conducted by Wilson & Stansell (1983) reported the presence of aflatoxins in agricultural products under drought-stress conditions. Furthermore, research conducted in Niger demonstrated that a drought lasting less than ten days caused a significant increase in aflatoxin contamination in the field. However, the extent of aflatoxin contamination depends on the severity of the water stress, the stage at which the stress is induced, and the soil and/or ambient temperature. The effect of terminal drought on aflatoxin contamination has been well established (Hamidou et al., 2014). Abbas et al. (2009) observed a higher level of aflatoxin concentration in corn under drought stress compared to non-stress conditions. Although a significant amount of aflatoxin contamination was reported under drought conditions, using tolerant varieties under drought stress could help reduce aflatoxin contamination in the field.

Flooding/excessive rainfall

Flooding is considered one of the most severe environmental stresses affecting plant productivity. Yearly, several floods are experienced on many farmlands, causing the loss of millions of dollars worth of agricultural products (Mohd Ikmal et al., 2021). This situation is expected to get worse due to climate change. Simultaneous flooding and aflatoxin production may incur more economic losses and worsen food safety issues. Based on the author's knowledge, although no studies have specifically reported the effect of flooding on aflatoxin production, a study by Das et al. (2012) reported that under ambient conditions (submerged fermentation), rice straw could favor aflatoxin production. Since the increase in moisture is positively correlated with aflatoxin production, it can be safely concluded that aflatoxin growth is favored when plants are submerged, affecting plant metabolism and functionality. Moreover, a recent survey conducted in sub-Saharan African (SSA) countries on food contamination reported that crop samples collected during the rainy season recorded higher aflatoxin content than dry-season crop samples (Benkerroum, 2020). This report also indicated that flooding might promote

aflatoxin production. The increased aflatoxin level is also linked to delayed harvest, late rainfall, irrigation, and dew during warm weather conditions. Aflatoxin levels were observed to be higher in crops that received more than 50 mm of rain during boll opening. Furthermore, when high rainfall is experienced at the pre-flowering stage, the crop has a high amount of aflatoxin concentration (Benkerroum, 2020). Concerning this, research needs to be conducted on the impact of submergence on aflatoxin production, focusing on the extent to which aflatoxin contaminates plants at different stages of plant production, whether crop resilience to submergence can help reduce aflatoxin contamination, and to what extent. This research is significant to ensure food security due to increased flooding on farmland, especially paddy fields. Since the production of aflatoxin is dependent on climate, it has been suggested and established that climatic change can result in a drastic change in the fungal population consequently bringing about the emergence of new mycotoxigenic fungal strains, favoring the already existing mycotoxin productions in food (Gbashi et al., 2018).

Other factors

Apart from climate change factors, other factors such as pH, fungal strain, substrate, nature of the soil, and availability of nutrients like carbohydrates, phosphates, zinc, and nitrogen play an important role in aflatoxin production (Benkerroum, 2020; Daou et al., 2021; Somda et al., 2023). The immediate fungi environment and its pH value play significant roles in the fungal growth and development, and production of mycotoxin. Studies have shown that the presence of "hydrogen ion concentration" (pH) in the fungus' immediate surroundings directly impacts fungal development, either through its action on cell surfaces or through an indirect effect on nutrient availability (Daou et al., 2021; Abubakar et al., 2013). For instance, at pH 4.0 and 7.0, the growth of *A. carbonarius*, a fungus isolated from wine and table grapes, was enhanced compared to that at pH 2.6, irrespective of water activity (Abubakar et al., 2013).

Another important factor influencing fungal contamination is the type of soil. Aflatoxin occurrences differ considerably among crops depending on the soil type. Based on the report, light sandy soils promote fungi growth, especially when subjected to water stress, while lower fungal infection are experienced in heavier soils possibly

due to their water retention ability, which helps to maintain irrigation frequency and to decrease the water stress effect (Marrez & Ayeshe, 2022; Somda et al., 2023). Finally, to avoid aflatoxin contamination at the pre-harvest stage, sufficient nutrient availability to plants is required, especially nitrogen. Crops may be susceptible to aflatoxin contamination if the root zone lacks mineralized nitrogen. Inadequate levels of mineralized nitrogen may be caused by leaching due to high amounts of water droplets and water stress (Abbas et al., 2009). Based on experimental results, a low quantity of aflatoxin was observed in corn produced with higher nitrogen (120 kg/ha). In contrast, corn produced with a lower amount of nitrogen (80kg/ha) recorded high aflatoxin contamination (Abbas et al., 2009). This signifies that a proper mixture of macronutrients is essential in crop management.

Impact of aflatoxin consumption on human health

Food insecurity issues are predominant in undeveloped and developing nations and emerging and transitional economies as a significant portion of the population lacks access to safe, nutritious, and cheap agricultural products (Udomkun et al., 2017). Estimates suggest that approximately 1500 diarrheal cases are reported annually across the globe, and more than 70% are associated with biological contamination of food, causing about 3 million deaths (Yáñez et al., 2002). This may be attributed to several factors, including consuming contaminated agricultural commodities. In the future, many health-related problems may be encountered due to the emergence and prevalence of aflatoxins in agricultural commodities ingested by humans.

In the 1960s, hundreds of deaths were reported in Turkey because the peanut was severely infested by mold, leading to the discovery of aflatoxins (Khlangwiset et al., 2011; Petrović et al., 2023). Since then, Turkey has recorded a series of outbreaks that have resulted in morbidity and mortality (Atherstone et al., 2016). Diseases caused by the ingestion of aflatoxin are referred to as aflatoxicosis. AFB1 is absorbed in the small intestine and transported to the bloodstream, where red blood cells and plasma proteins are transferred to the liver. The toxin transported to the liver is broken down by an enzyme known as microsomal-mixed function oxidase (MFO), a member of the Cytochrome P450 (CYP450) superfamily (Janik et al., 2020). AFB1 is converted to reactive 8, 9-epoxide

formed by Cytochrome P450 enzymes which can bind to DNA and proteins. Mechanistically, it is understood that the reactive AFB1 epoxide binds to the guanines at the N7 position. Furthermore, TA to GC transversions may result from AFB1-DNA adducts. A reactive glutathione S-transferase system located in the cytosol and microsomes catalyzes the conjugation of activated AFB1 with decreased glutathione, resulting in aflatoxin being excreted (Bennett & Klich, 2003). Aflatoxins can be excreted through bile, feces, urine, semen, milk, and eggs. In humans, aflatoxin M1 (AFM1), aflatoxin P1 (AFP1), and free guanine residues (AFB1-N7-guanine) are excreted in urine while they are excreted via the bile in rats. In ruminant animals, AFB1 is excreted via feces and AFM1 predominantly through urine and milk (Thakur et al., 2022).

The previous study showed that disruption of the human immune system was caused by aflatoxin infection, making it vulnerable to other infectious diseases. Aflatoxins have also been associated with congenital disabilities and stunted growth in children exposed for a long time (Jallow et al., 2021). AFB1 intoxication is hazardous, especially in regions where the hepatitis B virus (HBV) is prevalent. A survey indicated that HBV-positive people are at greater risk of developing liver cancer than HBV-negative people. AFB1 was regarded as the primary causative agent of hepatocellular carcinoma (HCC), a common type of liver cancer (Janik et al., 2020). The metabolic implication of aflatoxin intoxication includes disruption of the synthesis of protein, RNA, and DNA; depletion of the activities of the miscellaneous enzymes; disruption of the synthesis of lipids as well as esters, phospholipids, and triglycerides; and depletion of glucose metabolism (Giray et al., 2007; Petrović et al., 2023). The occurrence of HCC is triggered by dominant-negative oncogenes and changes in the P53 tumor-suppressing gene (Giray et al., 2007).

Aflatoxins' unpleasant impacts on human health are highly dynamic based on the type of exposure to contaminated feed. The effects can either be acute, witnessed within a few days after consumption of a significant amount of aflatoxin-contaminated food, or chronic, showing after many months or years of ingestion. Acute effects experienced can be vomiting, jaundice, liver problems, abdominal pains and even death for the affected person while chronic consequences can be associated with weakened immunity, poor growth, cancer development, mutagenicity (Yard

et al., 2013; Barajas-Ramirez et al., 2021; Petrović et al., 2023), liver damage, and even death because of the accumulation of toxins within the body (Yan et al., 2020). Several studies on aflatoxin toxicity, especially AFB1, have been conducted with sufficient animal and human epidemiological findings that prove the teratogenicity, mutagenicity, and carcinogenicity of aflatoxins. Even more, studies have shown that aflatoxin can cause cancer, attacking different organs such as the stomach, lung, and liver (Ndagijimana et al., 2020).

The severity of an aflatoxin outbreak is influenced by several variables, including the contamination level of the mycotoxin, the individual's age and prior health condition, and the toxicity and possible impacts of other chemicals the person is exposed to (Majeed et al., 2018). In 2004, over 317 individuals were admitted to the hospital due to the ingestion of aflatoxin-contaminated food in Kenya, resulting in 125 fatalities. Recurrent incidences of this nature were reported in 1981 and 2005 in Kenya, with 12 and 16 deaths recorded, respectively (Agriopoulou et al., 2020). In 1975, an aflatoxin outbreak was witnessed in India among the Bhils Tribe, which had earlier fed on aflatoxin-contaminated maize. This led to ascites and portal hypertension, affecting about 400 individuals (Filazi & Tansel Sireli, 2013). Similarly, 100 people were reported dead in India due to a hepatitis outbreak that may have been attributed to the ingestion of contaminated maize (Bennett & Klich, 2003). In addition, 14.1 million new cancer cases and 8.2 million fatalities were recorded globally in 2012. Liver cancer is the second-leading cause of human death after lung cancer, accounting for approximately 745,000 deaths per year. In the same year, various parts of Africa recorded 847,000 cancer cases and 591,000 fatalities. Over 80% of the cases of HCC occur in poor countries due to the risks of dietary aflatoxin exposure and chronic hepatitis B and C (Lukwago et al., 2019). In Nigeria, ingestion of aflatoxin-contaminated agricultural products was linked to 7,761 liver cancer cases. Elsewhere, in Tanzania, approximately 3,334 cases of HCC were estimated, with 95% of the cases resulting in death (Gbashi et al., 2018). In the same country, another aflatoxicosis outbreak was reported recently. This outbreak caused the deaths of 20 out of the 68 individuals affected (Benkerroum, 2020). Mycotoxin contamination of agricultural products is considered a big threat to public health in sub-Saharan Africa, with approximately 250,000

HCC-related fatalities yearly caused by aflatoxins alone (Echodu et al., 2019). Worldwide, 4.5 billion people are affected by aflatoxin exposure, mostly in underdeveloped countries (Rudraboyina et al., 2023).

The implication of aflatoxin exposure on infants and children

Globally, most children are exposed to a significant amount of aflatoxins at an early stage of their development and throughout their entire life because most communities rely heavily on the subsistence agriculture system for their daily diet, and they are unaware of the existence of aflatoxins (Mupunga et al., 2017). Children and infants are exposed to aflatoxins at different growth and developmental stages through maternal food ingestion during pregnancy, breastfeeding, and post-weaning diets, particularly in areas where maize is the primary food source. After children are weaned from breastfeeding, their exposure to aflatoxin skyrockets; on the other hand, exposure during pregnancy also tremendously affects the infants (Khlangwiset et al., 2011). A study was reported on the effect of aflatoxin exposure on Gambian infants (6, 12, and 18 months). The result revealed that aflatoxins were the infants' leading cause of growth retardation (Watson et al., 2018). It was also found that aflatoxin exposure was associated with lifelong cognitive and physical deficits (Passarelli et al., 2020). In 1988, Malaysia experienced an aflatoxicosis outbreak in Perak state due to consumption of contaminated noodles with up to 3mg of aflatoxins, resulting in the deaths of 13 children (Sowley, 2016). Research carried out by Gong and his colleagues reported that out of 479 children and infants studied, aflatoxins and aflatoxin-albumin (AF-alb) were found in 99%, and the amount increased as they got older due to the consumption of complementary foods (Achaglinkame et al., 2017). At the early stage, exposure to aflatoxins and AF-alb was attributed to stunted growth in infants and children (Gong et al., 2003). aflatoxins were also found in the umbilical cords, signifying the presence of toxins around the placenta and beyond. Furthermore, maternal exposure was found to correlate positively with breast milk aflatoxin levels (Achaglinkame et al., 2017).

Stunted growth in children has become predominant in certain parts of the world, including in South and East Asia and sub-Saharan Africa, despite significant feeding and the adoption of other nutrition intervention schemes

in affected regions (Mitchell et al., 2017). In Bangladesh, childhood stunting is prevalent. Approximately 36% of children below the age of 5 years are shorter than the normally expected height of their age or are stunted, with 15% being severely stunted (Mahfuz et al., 2019). Based on the low height-for-age z-score (HAZ) recorded, stunting can be defined as height below two standard deviations (SD) of the standard average. The HAZ score is a metric that indicates how far a child is from the average height-for-age, with a HAZ of 2 indicating stunting growth in a child (more than two standard deviations below average height) and a HAZ of 3 indicating severe stunting in a child (Ahlberg et al., 2018). According to a review of nutritional interventions on child growth, the highest growth improvement provided by feeding and dietary programs is a 0.7 increase in HAZ (Mitchell et al., 2017). Stunting is a well-documented risk indicator of poor development in a child and chronic malnutrition, and it has been linked to aflatoxin exposure (Ahlberg et al., 2018). Impaired growth and stunting are considered significant problems because stunting has long-term consequences beyond infancy and childhood. The long-term impact may include reduced productivity, increased health complications, and lower academic achievements (Ahlberg et al., 2018). Chronic aflatoxin exposure has been associated with kwashiorkor. Research conducted in the last 30 years have shown that children with kwashiorkor have significant amounts of aflatoxin in their urine and blood samples compared to healthy children. Similarly, a study in Cameroon revealed the presence of AFB1 in the blood and urine of kwashiorkor patients (Mupunga et al., 2017). A survey conducted on Egyptian infants showed that aflatoxin is prevalent in the urine of children suffering from kwashiorkor, followed by marasmus patients. At the same time, no aflatoxins were detected in the urine samples of the control group (Hatem et al., 2005). In Nigeria, a post-humous autopsy study conducted on children with kwashiorkor and other miscellaneous diseases showed significant amounts of aflatoxins in their lungs due to the ingestion of infected maize (Oyelami et al., 1997; Gbashi et al., 2018). Out of the 20 children who suffered from kwashiorkor, 18 cases ended in death, while 13 out of the 20 children with miscellaneous diseases were reported dead. This study has shown that infants are exposed to a significant level of aflatoxins, which may be accumulated in the lungs (Oyelami et al., 1997).

Based on available and established data, aflatoxin prevalence is experienced in Africa and certain Asian countries. This can be attributed to the lack of strict safety regulations to curb the level of aflatoxins present in food commodities consumed by the population. This has resulted in major health consequences for people in this part of the world. In addition, the presence of suitable environmental conditions for aflatoxin development, technological hurdles, a high rate of illiteracy among farmers and consumers, lack of awareness, poor storage conditions and facilities, and an overall high rate of poverty may also be considered as possible reasons for a high level of aflatoxin in Africa and a certain part of Asia (Ismail et al., 2018). Thus, adequate detection techniques and control methods are essential to combat the problems of aflatoxin in food.

Possible strategies for mitigating aflatoxin exposure

Development of a tolerant cultivar

The development of resistant varieties has been established as one of the most effective methods for preventing aflatoxin contamination in crops. This can be done through molecular plant breeding and genetic engineering approaches. The molecular breeding technique, which includes the pyramiding of quantitative trait loci (QTLs) via marker-assisted selection, has extensively been used to develop tolerant varieties to abiotic stresses, such as submergence and drought. The pyramiding of QTLs via marker-assisted selection implies drought-tolerant cultivars which helps to reduce aflatoxin contamination (Rudraboyina et al., 2023). Since its establishment, many newly developed genotypes using molecular markers can be screened in the field for their tolerance against aflatoxin production. This method can help discover more genotypes and germplasm, which can be used for further breeding programs. The development of resistant varieties of crops such as maize has been realized through the screening of new tools during field and laboratory screening. The techniques (restriction fragment length polymorphism, RFLP analysis for corn populations) have shown that different resistant traits can be successfully developed into agronomically useful germplasm while proteomics has helped to identify proteins associated with resistance (RAPs). The pin-bar technique has been used to discover two resistant inbreds (Mp420 and Mp313E), which have passed field trials in various locations and been distributed as sources of resistant germplasm

(Brown et al., 2004). Furthermore, 36 inbred maize lines collected from West and Central Africa were screened and evaluated for aflatoxin resistance. The result showed that aflatoxin levels in over half of the inbred lines were lower than in the resistant US lines. The same research team registered six tropical maize germplasm lines resistant to aflatoxins (Xu et al., 2022).

A wide range of biotechnology techniques have been developed to understand better the resistance mechanisms and to identify genes, proteins, and pathways involved during host-pathogen interactions for aflatoxin contamination in the crop. These molecular techniques include RNA interference, microarray, whole genome sequencing, RNA-sequencing, proteomics, and metabolomics (Xu et al., 2022). The genetic engineering approach has been used to achieve over 80% decrease in groundnut content via the *RNAi* technique to silence aflatoxin-producing genes (*aflC*, *aflP*, *aflR*, and *aflS*). Furthermore, by silencing the genes (*aflM* and *aflP*) via host-induced gene silencing (HIGS) and overproducing genes responsible for plant defense (*MsDef1* and *MtDef4.2*), groundnuts with reduced aflatoxin content were developed (Pandey et al., 2019). Another study by Thakare et al. (2017) showed that host-induced gene silencing could efficiently eliminate aflatoxin content in transgenic maize. This research demonstrated that small interfering RNA molecules could be used to silence aflatoxin biosynthesis in maize. Regulation of enzymatic antioxidants has been proven to inhibit aflatoxin production. For instance, superoxide's intracellular accumulation helps to inhibit aflatoxin production by downregulating *aflR* expression, and the addition of Cu/ZnSOD externally decreased aflatoxin production (Furukawa & Sakuda, 2019). The use of genetic engineering as well as plant breeding techniques for the development of resistant crops is considered a sustainable and eco-friendly long-term strategy for pre-harvest interventions.

Biological methods

The biological detoxification of mycotoxins works in two major processes which are enzymatic degradation and sorption, both of which can be accomplished through biological systems (Aliabadi et al., 2013). Biological methods have been proven effective and promising in mitigating aflatoxin contamination. Several organisms have been tested for their ability to control aflatoxin contamination including yeasts, bacteria, and

some non-toxicogenic fungal strains of *A. flavus* and *A. parasiticus* (Somda et al., 2023). Application of non-toxicogenic strains of *A. flavus* and *A. parasiticus* in the maize field has significantly reduced aflatoxin contamination. The non-toxicogenic strains compete with aflatoxigenic strains in the field, occurring in the same niche. Hence, they displace the toxicogenic strains (Thakur et al., 2022; Rudraboyina et al., 2023). The use of lactic bacteria (LABs) such as *Lactococcus lactis* subsp. *lactis*, *Pediococcus acidilactici*, *Lactobacillus acidophilus*, and *Enterococcus avium* has been proven effective in preventing and eradicating aflatoxins in agricultural commodities (Peles et al., 2021; Roy et al., 2023). This technique has been established in several studies. For instance, an experiment by Asurmendi et al. (2014) showed that the LABs inhibited the activities of two *A. flavus* strains as well as the production of AFB1 in the brewer's grains. Similarly, another study by Saladino et al. (2016) reported 84.1%–99% reduction in the aflatoxin content of bread and an increase in the shelf-life due to the inhibitory activities of LABs on aflatoxins. Apart from LABs, non-lactic acid bacteria including *Pseudomonas* spp., *Myxococcus* spp., *Brachybacterium* spp., *Cellulosimicrobium* spp., *Nocardia* spp., *Escherichia* spp., *Stenotrophomonas* spp., and *Klebsiella* spp., also cause inhibitory activities on the aflatoxin growth and production of molds. A study by Peles et al. (2021) demonstrated that a *Bacillus subtilis* strain decreased the concentration of AFB1 by 60%–95% in contaminated agricultural commodities. Using yeast species such as *Debaryomyces* spp., *Aureobasidium pullulans*, *Zygosaccharomyces* spp., *Saccharomyces* spp., and *Schizosaccharomyces* spp. has been proven effective in reducing the aflatoxin production in food. However, partially, or wholly eradicating aflatoxins using yeast depends on the strain. The aflatoxin reduction varies from 15%–100% for AFB1 and 60%–90.3% for AFM1 (Pickova et al., 2021). Aflatoxin contamination can also be controlled by using mycotoxin adsorbents (activated carbon, kaolinite, clay, bentonites) and binders, which aim to prevent mycotoxins from entering the intestinal tract of animals by adsorbing toxins from the surface (Kamle et al., 2019; Wanniarachchi et al., 2023).

Pre-harvest practices

Pre-harvest strategies for preventing aflatoxins include good manufacturing practices (GMPs), appropriate environmental factors, good

agricultural practices (GAPs), and favorable storage practices. Good farming practices include the implementation of a crop rotation program; the use of registered fungicides, insecticides, and herbicides for control of insect damage, fungal infection, and weed eradication; the proper treatment of the seed bed; the soil analysis to determine the need to add fertilizers, and the improvements in genetic synthesis to reduce mycotoxin production (Agriopoulou et al., 2020; Marrez & Ayesh, 2022; Rudraboyina et al., 2023). For example, legume crop rotation with maize improves soil fertility and disrupts pest and disease cycles. Additionally, rotating maize with non-host crops can help to minimize plant residues that could allow the buildup of inoculum. Adopting and maintaining GAPs in the proper manner helps to enhance the safety and the quality of food and other agricultural commodities. Good agricultural practices (GAPs) also offer smallholder farmers the additional advantages of enhanced yield and decrease overall post-harvest losses (Xu et al., 2022).

Chemical methods

Hydrochloric acid (HCL) has been proven effective in reducing AFB1 concentration by 19% in 24 hours. Alkalinity also makes aflatoxins unstable. Ammonization has been shown to decrease aflatoxin content by more than 99%. The use of ammonia to destroy aflatoxins has been extensively investigated and proven to be a success both in the field and laboratory (Pickova et al., 2021; Zhang et al., 2023; Somda et al., 2023). Moreover, a study by Abubakar et al. (2013) revealed that certain alkaline media could help to prevent *A. parasiticus* growth and sporulation. Ozone treatment has been reported to reduce contamination by the degradation of mycotoxins. It can also be used in gaseous form to avoid increase in moisture. However, treatment using this method can take a long time to work and can result in the oxidation of fat components, thus reducing the quality of the food (Daou et al., 2021; Zhang et al., 2023). Ozone exposure inhibits the germination of conidia and brings about a change in the morphology of hyphae which consequently resulting to the death of hyphae and ROS accumulation of the fungal species from *Aspergillus*, *Fusarium*, and *Penicillium* genera (Xue et al., 2023). Using fungicides is also considered as one of the most effective methods in preventing fungal invasion before harvest and, subsequently, mycotoxin contamination (Roy et al., 2023). However, research on fungicide usage is debatable. While other literature found them

effective, some believe that in some cases, they enhance aflatoxin production and poses a threat to human and animal health (Daou et al., 2021).

Physical methods

The use of physical methods, including adequate drying and physical treatment, can assist in mitigating aflatoxin contamination at the post-harvest stage and reduces the effects of contamination and the subsequent accumulation of mycotoxins in crops (Somda et al., 2023). One of the prerequisites for completely eradicating aflatoxin content is the restriction of colonization by aflatoxin-producing fungi on the surface layers of grains. Dehulling techniques are used to remove the grain's outer layers, removing around 93% of aflatoxin (Pickova et al., 2021). Up to 50% of the aflatoxin content in corn is eliminated by milling. Extrusion can help to reduce aflatoxin content by 50%–80% depending on the temperature and the grain moisture (Karlovsky et al., 2016). Furthermore, 40% of aflatoxin content is reduced in maize by roasting, while temperatures beyond 160°C destroy AFB1 (Karlovsky et al., 2016).

Irradiation is a valuable method for inhibiting some mycotoxin's action. A previous study had shown that exposure of feed and food to γ -radiation, microwave heating, and solar radiation are effective ways to decontaminate any left residues of aflatoxins present in food. However, the efficiency of these methods depends on few factors including type of fungus, dosage applied, food composition, moisture content, and storage conditions (Adejumo & Adejoro, 2014). New technologies such as electron beam, gamma irradiation, microwave heating, electrolyzed water, cold plasma, UV, and pulsed light have been proven effective in mitigating the contamination of aflatoxins (Pankaj et al., 2017; Wanniarachchi et al., 2023). A previous study reported 59 to 88% aflatoxin reduction when 10kGy irradiation was induced while in another study, 11 to 21% aflatoxin reduction was observed when 15kGy irradiation was employed (Pickova et al., 2021). UV-A irradiation has also been reported to reduce AFB1 and AFM1 concentrations in pure water by 70% and 84%, respectively, at a dose of 1200mJ/cm (Stanley et al., 2020). Due to the effect of high temperatures on food, non-thermal approaches such as pulsed electric fields (PEF) have been adopted and proven effective in reducing aflatoxin concentration in agricultural commodities without losing their quality and nutritional value (Vijayalakshmi et al., 2018). Apart from PEF,

cold plasma has also been effective in controlling mycotoxin contamination by destroying the cell wall of fungi and their DNA, allowing the leakage of intercellular components. Other studies on the effect of cold plasma on mycotoxins have shown that they are either partially or entirely destroyed (Daou et al., 2021).

The use of nanobiotechnology

Recently, nanotechnology has drawn increasing attention from scientists around the world due to its extensive use in numerous fields including field management of agricultural pests and diseases (Wen et al., 2023). This method appears to be a novel, promising, less expensive technique that can provide environmentally friendly control of mycotoxigenic fungi and mycotoxins in the agriculture and food industries. Mycotoxins or pathogens in food and feed can be removed by the use of nanomaterials like nanosilver (AgNPs), zinc oxide nanoparticles (ZnO-NPs), selenium nanoparticles (SNPs), and copper nanoparticles (CuNPs) (Haque et al., 2020). A study by Gacem & Abd-Elsalam (2022) proved that treatment of maize with ZnO-NPs at the rate of 100g/g of ZnO-NPs, significantly suppress the development of *A. flavus*. Haque et al. (2020) reported that CuNPs demonstrate high antifungal activity against *Curvularia lunata*, *Alternaria alternata* and *Fusarium (culmorum, oxysporum, graminearum)* fungi. Recently, Abd El-Ghany et al. (2023) reported that the conidial germination of the tested mycotoxigenic fungi is significantly inhibited by bioinspired AgNPs. Increased DNA and protein leakage is a result of the bioinspired AgNPs, indicating that membrane permeability and integrity have been compromised. Interestingly, at concentrations lower than 8 μ g/mL, the biogenic AgNPs totally prevent the growth of total aflatoxins and ochratoxin A. Recent studies discovered that mycotoxin contamination in the food chain can be reduced, and toxigenic fungal formation can be inhibited using nanomaterials loaded with phytochemicals (Haque et al., 2020). The use of AgNPs at lower concentrations than the minimum inhibitory concentration has been reported to inhibit AFB1 production. This result suggests that AgNPs can be considered as an important weapon to mitigate aflatoxin contamination in vulnerable crops in the field (Mousavi & Pourtalebi, 2015).

The use of essential oils

The use of essential oils (EOs) is considered as a good substitute to synthetic chemicals because

of their antifungal effect against *Aspergillus* and other fungi, which inhibits the growth of mycelium, spore germination and production of aflatoxins (Nerilo et al., 2020). The use of EOs is less expensive and it does not have the same drawbacks inherent to synthetic chemical pesticides (Nerilo et al., 2020). In the food sector, EOs are frequently employed as natural food preservatives to increase product shelf life and as flavoring agents (Belasli et al., 2020). Essential oils (EOs) has been used to inhibit aflatoxin production. Miri et al. (2019) reported that at 4 μ L/mL of EO, *Cicuta virosa* L., EOs reduced the growth of *A. flavus* and the formation of AFB1. At 2 μ L/mL, the AFB1 was roughly halved in comparison to the control. Furthermore, AFB1 synthesis was completely reduced at the minimal fungicidal concentration value and was inhibited by EOs values between 0.25mg/mL (15% drop) and 1.50mg/mL (86% decrease). Additionally, study by Belasli et al. (2020) revealed that in a dose-dependent manner, the EOs demonstrate a broad antifungal spectrum against additional species.

Conclusion

Cereals are considered as staple foods and raw materials in the food industry. The contamination of cereals by aflatoxins is a global concern, posing a significant threat to food security and the well-being of humankind and thus, disrupting the world economy. The production of aflatoxins is triggered by several environmental factors such as temperature, humidity, pH, soil type and storage conditions. Consumption of aflatoxin contaminated foods causes serious health problems to human especially children including retarded growth and even death, depending on the severity. With the use of appropriate control measures can help to reduce aflatoxin production to certain extent if not completely. These approaches include breeding of resistant varieties via genetic engineering, biological, chemical, physical, and proper post-harvest handling of cereal crops. Furthermore, the adoption of new technologies such as nanotechnology, cold plasma and essential oils has shown to effectively reduce aflatoxin production in crops. In the future, more studies should be conducted to determine the role of over-expression or down-regulation of specific genes in controlling aflatoxin contamination. Moreover, complete detoxification of any aflatoxin residues in agricultural commodities through the combination of different control

measures should be encouraged. Finally, raising awareness regarding the detrimental consequences of aflatoxins should be prioritized, especially in developing nations where aflatoxin-associated disease outbreaks frequently occur due to suitable climates, high levels of illiteracy, and poverty, resulting in consuming contaminated foods.

Acknowledgment: All authors would like to thank the Malaysia Ministry of Education for research grants FRGS/1/2018/STG03/UKM/02/1 and LRGS/1/2019/UKM/01/5/4. The first author would also like to thank Universiti Kebangsaan Malaysia (UKM) for the Zamalah Research Scholarship scheme award.

Conflict of interest: There are no conflict of interest

Authors' contributions: The manuscript was drafted by Y.O.O., while suggestion, comment, improvement of draft, proofreading, editing, and finishing was done by N.A.A.S., I.A.A., O.B.U., N.A.M.Z. and M.N.I.

Ethics approval: Not applicable

References

- Aasa, A.O., Fru, F.F., Adelusi, O.A., Oyeyinka, S.A., Njobeh, P.B. (2023) A review of toxigenic fungi and mycotoxins in feeds and food commodities in West Africa. *World Mycotoxin Journal*, **16**(1), 33-47.
- Abbas, H.K., Wilkinson, J.R., Zablotowicz, R.M., Accinelli, C.E.S.A.R.E., Abel, C.A., Bruns, H.A., Weaver, M.A. (2009) Ecology of *Aspergillus flavus*, regulation of aflatoxin production, and management strategies to reduce aflatoxin contamination of corn. *Toxin Reviews*, **28**(2-3), 142-153.
- Abd El-Ghany, M.N., Hamdi, S.A., Korany, S.M., Elbaz, R.M., Emam, A.N., Farahat, M.G. (2023) Biogenic silver nanoparticles produced by soil rare actinomycetes and their significant effect on *Aspergillus*-derived mycotoxins. *Microorganisms*, **11**(4), 1-17.
- Abubakar, A., Suberu, H.A., Bello, I.M., Abdulkadir, R., Daudu, O.A., Lateef, A.A. (2013) Effect of pH on mycelial growth and sporulation of *Aspergillus parasiticus*. *Journal of Plant Sciences*, **1**(4), 64-67.
- Achaglinkame, M.A., Opoku, N., Amagloh, F.K. (2017) Aflatoxin contamination in cereals and legumes to reconsider usage as complementary food ingredients for Ghanaian infants: A review. *Journal*

- of Nutrition and Intermediary Metabolism*, **10**, 1-7.
- Adejumo, T.O., Adejoro, D.O. (2014) Incidence of aflatoxins, fumonisins, trichothecenes and ochratoxins in Nigerian foods and possible intervention strategies. *Food Science and Quality Management*, **31**, 1-22.
- Ahlberg, S., Grace, D., Kiarie, G., Kirino, Y., Lindahl, J. (2018) A risk assessment of aflatoxin M1 exposure in low and mid-income dairy consumers in Kenya. *Toxins*, **10**(9), 348.
- Agriopoulou, S., Stamatelopoulou, E., Varzakas, T. (2020) Advances in occurrence, importance, and mycotoxin control strategies: Prevention and detoxification in foods. *Foods*, **9**(2), 1-50.
- Ali, N. (2019) Aflatoxins in rice: Worldwide occurrence and public health perspectives. *Toxicology Reports*, **6**, 1188-1197.
- Aliabadi, M., Alikhani, F., Mohammadi, M., Darsanak, R. (2013) Biological control of aflatoxins. *European Journal of Experimental Biology*, **3**(2), 1-6.
- Andrade, P., Caldas, E. (2015) Aflatoxins in cereals: worldwide occurrence and dietary risk assessment. *World Mycotoxin Journal*, **8**(4), 415-431.
- Anonymous (2021) FAO cereal Supply and demand Brief. FAO Website. Food situation. February. Food and Agriculture Organization (FAO) World. http://www.fao.org/world_food_situation/csdb/en [accessed on July 13,2021].
- Atherstone, C., Grace, D., Lindahl, J., Kang'ethe, E. and Nelson, F. (2016) Assessing the impact of aflatoxin consumption on animal health and productivity. *African Journal of Food, Agriculture, Nutrition and Development*, **16**(03), 10949-10966.
- Asurmendi, P., Pascual, L., Dalcero, A., Barberis, L. (2014) Incidence of lactic acid bacteria and *Aspergillus flavus* in brewer's grains and evaluation of potential antifungal activity of these bacteria. *Journal of Stored Products Research*, **56**, 33-37.
- Awuchi, C.G., Ondari, E.N., Eseoghene, I.J., Twinomuhwezi, H., Amagwula, I.O., Morya, S. (2021) Fungal growth and mycotoxins production: Types, toxicities, control strategies, and detoxification. In "Fungal Reproduction and Growth", S. Sultan, G.K.S. Singh (Eds.), pp 1-20, Intech Open.
- Battilani, P., Toscano, P., Van der Fels-Klerx, H., Moretti, A., Camardo Leggieri, M., Brera, C., Rortais, A., Goumperis, T., Robinson, T. (2016) Aflatoxin B1 contamination in maize in Europe increases due to climate change. *Scientific Reports*, **6**(1), 1-8.
- Barajas-Ramirez, J.A., Moncada-Abaunza, D.A., Gómez-Espinoza, M.G. (2021) Mycotoxins in foods, from the field to the plate: A review. *International Food Research Journal*, **28**(2), 230-247.
- Belasli, A., Ben Miri, Y., Aboudaou, M., Aït Ouahioune, L., Montañes, L., Ariño, A., Djenane, D. (2020) Antifungal, antitoxigenic, and antioxidant activities of the essential oil from laurel (*Laurus nobilis* L.): Potential use as wheat preservative. *Food Science and Nutrition*, **8**(9), 4717-4729.
- Benkerroum, N. (2020) Aflatoxins: Producing-molds, structure, health issues and incidence in Southeast Asian and Sub-Saharan African countries. *International Journal of Environmental Research and Public Health*, **17**(4), 1215.
- Bennett, J.W., Klich, M. (2003) Mycotoxins. *Clinical Microbiology Reviews*, **16**, 1-20.
- Blankson, G.K., Mills-Robertson, F.C., Ofori, I.W. (2019) Survey of occurrence levels of aflatoxins in selected locally processed cereal-based foods for human consumption from Ghana. *Food Control*, **95**, 170-175.
- Block, A., Vaughan, M., Schmelz, E., Christensen, S. (2018) Biosynthesis and function of terpenoid defense compounds in maize (*Zea mays*). *Planta*, **249**(1), 21-30.
- Brown, R., Chen, Z., Menkir, A., Cleveland, T. (2004) Using biotechnology to enhance host resistance to aflatoxin contamination of corn. *African Journal of Biotechnology*, **2**, 1-7.
- Daou, R., Joubrane, K., Maroun, R.G., Khabbaz, L.R., Ismail, A., El Khoury, A. (2021) Mycotoxins: Factors influencing production and control strategies. *AIMS Agriculture and Food*, **6**(1), 416-447.
- Dadzie, M., Oppong, A., Ofori, K., Eleblu, J., Ifie, E., Blay, E., et al. (2019) Distribution of *Aspergillus flavus* and aflatoxin accumulation in stored maize grains across three agro-ecologies in Ghana. *Food Control*, **104**, 91-98.
- Das, A., Angayarkanni, J., Bhattacharya, S., Palaniswamy, M. (2012) Evaluation of process parameters influencing aflatoxin B1 synthesis from *Aspergillus flavus* MTCC 2798 using rice straw under submerged fermentation. *International Journal of Pharmacy and Biological Sciences*, **2**(2), 94-105.

- Di Domenico, A.S., Busso, C., Hashimoto, E.H., Frata, M.T., Christ, D., Coelho, S.R. (2016) Occurrence of *Aspergillus sp.*, *Fusarium sp.*, and aflatoxins in corn hybrids with different systems of storage. *Agronomy*, **38**(1), 111-121.
- Elzupir, A.O., Alamer, A.S., AlRajhi, M., Idriss, H. (2018) Assessment of health risks from aflatoxins in rice commercialized in Riyadh, Kingdom of Saudi Arabia. *Quality Assurance and Safety of Crops and Foods*, **10**(3), 255-260
- Echodu, R., Malinga, G.M., Kaducu, J.M., Ovuga, E., Haesaert, G. (2019) Prevalence of aflatoxin, ochratoxin and deoxynivalenol in cereal grains in northern Uganda: Implication for food safety and health. *Toxicology Reports*, **6**, 1012-1017.
- Filazi, A., Tansel Sireli, U. (2013) Occurrence of aflatoxins in food. In "Aflatoxins – Recent Advances and Future Prospects", M. Razzaghi-Abyaneh (Ed.), pp. 143-169. Intech Open.
- Furukawa, T., Sakuda, S. (2019) Inhibition of aflatoxin production by paraquat and external superoxide dismutase in *Aspergillus flavus*. *Toxins*, **11**(2), 107.
- Gacem, M.A., Abd-El Salam, K.A. (2022) Nanomaterials for the reduction of mycotoxins in cereals. In: "Cereal Diseases: Nanobiotechnological Approaches for Diagnosis and Management", pp. 371-406. Singapore: Springer Nature Singapore.
- Garba, M. (2019) Mycotoxins consumption and burden of aflatoxin-Induced hepatocellular carcinoma in people subsisting on sorghum based products in the derived Savannah zone of Nigeria. *Asian Food Science Journal*, **8**(4), 1-10.
- Gbashi, S., Madala, N.E., De Saeger, S., De Boevre, M., Adekoya, I., Adebo, O.A., Njobeh, P.B. (2018) The socio-economic impact of mycotoxin contamination in Africa. Fungi and mycotoxins-their occurrence, impact on health and the economy as well as pre- and postharvest management strategies. In "Mycotoxins– Impact and Management Strategies", P.B. Njobeh, F. Stepman (Eds.), pp. 1-20, Intech Open.
- Giray, B., Girgin, G., Engin, A.B., Aydın, S., Sahin, G. (2007) Aflatoxin levels in wheat samples consumed in some regions of Turkey. *Food Control*, **18**, 23– 29.
- Gong, Y.Y., Egal, S., Hounsa, A., Turner, P.C., Hall, A.J., Cardwell, K.F., Wild, C.P. (2003) Determinants of aflatoxin exposure in young children from Benin and Togo, West Africa: the critical role of weaning. *International Journal of Epidemiology*, **32**, 556–562.
- Hamidou, F., Rathore, A., Waliyar, F., Vadez, V. (2014) Although drought intensity increases aflatoxin contamination, drought tolerance does not lead to less aflatoxin contamination. *Field Crops Research*, **156**, 103-110.
- Hatem, N.L., Hassab, M.H.A., Abd Al-Rahman, E.H., El-Deeb, S.A., El-Sayed Ahmed, R.L. (2005) Prevalence of aflatoxins in blood and urine of Egyptian infants with protein energy malnutrition. *Food and Nutrition Bulletin*, **26**, 49–56.
- Hanvi, D., Lawson-Evi, P., De Boevre, M., Goto, C., De Saeger, S., Eklu-Gadegbeku, K. (2019) Natural occurrence of mycotoxins in maize and sorghum in Togo. *Mycotoxin Research*, **35**(4), 321-327.
- Haque, M. A., Wang, Y., Shen, Z., Li, X., Saleemi, M. K., He, C. (2020) Mycotoxin contamination and control strategy in human, domestic animal and poultry: A review. *Microbial Pathogenesis*, **142**, 1-12.
- Iqbal, S., Asi, M., Hanif, U., Zuber, M., Jinap, S. (2016) The presence of aflatoxins and ochratoxin A in rice and rice products; and evaluation of dietary intake. *Food Chemistry*, **210**, 135-140.
- Ismail, A., Gonçalves, B.L., de Neeff, D.V., Ponzilacqua, B., Coppa, C.F., Hintzsche, H., Oliveira, C.A. (2018) Aflatoxin in foodstuffs: Occurrence and recent advances in decontamination. *Food Research International*, **113**, 74-85.
- Jallow, A., Xie, H., Tang, X., Qi, Z., Li, P. (2021) Worldwide aflatoxin contamination of agricultural products and foods: From occurrence to control. *Comprehensive Reviews in Food Science and Food Safety*, **20**(3), 2332-2381.
- Janik, E., Niemcewicz, M., Ceremuga, M., Stela, M., Saluk-Bijak, J., Siadkowski, A., Bijak, M. (2020) Molecular aspects of mycotoxins-A serious problem for human health. *International Journal of Molecular Science*, **21**, 1-21.
- Joubrane, K., Mnayer, D., EL Khoury, A., EL Khoury, A., Awad, E. (2020) Co-occurrence of aflatoxin B1 and ochratoxin A in Lebanese stored wheat. *Journal of Food Protection*, **83**(9), 1547–1552.
- Kamle, M., Mahato, D., Devi, S., Lee, K., Kang, S., Kumar, P. (2019) Fumonisin: Impact on agriculture, food, and human health and their management strategies. *Toxins*, **11**(6), 1-23.
- Karlovsky, P., Suman, M., Berthiller, F., De Meester, J., Eisenbrand, G., Perrin, I., Dussort, P. (2016) Impact of food processing and detoxification treatments on

- mycotoxin contamination. *Mycotoxin Research*, **32**(4), 179-205.
- Khlangwiset, P., Shephard, G., Wu, F. (2011) Aflatoxins and growth impairment: A review. *Critical Reviews in Toxicology*, **41**(9), 740-755.
- Kobayashi, Y., Higa, M., Higashiyama, K., Nakamura, F. (2020) Drivers of land-use changes in societies with decreasing populations: A comparison of the factors affecting farmland abandonment in a food production area in Japan. *PLOS ONE*, **15**(7), p.e0235846.
- Kumar, P., Mahato, D., Kamle, M., Mohanta, T., Kang, S. (2017) Aflatoxins: A global concern for food safety, human health and their management. *Frontiers in Microbiology*, **07**, 1-10.
- Lahouar, A., Marin, S., Crespo-Sempere, A., Saïd, S., Sanchis, V. (2016) Effects of temperature, water activity and incubation time on fungal growth and aflatoxin B1 production by toxinogenic *Aspergillus flavus* isolates on sorghum seeds. *Revista Argentina de Microbiología*, **48**(1), 78-85.
- Lai, X., Liu, R., Ruan, C., Zhang, H., Liu, C. (2015) Occurrence of aflatoxins and ochratoxin A in rice samples from six provinces in China. *Food Control*, **50**, 401-404.
- Lee, H.S., Nguyen-Viet, H., Lindahl, J., Thanh, H. M., Khanh, T.N., Hien, L.T.T., Grace, D. (2017) A survey of aflatoxin B1 in maize and awareness of aflatoxins in Vietnam. *World Mycotoxin Journal*, **10**(2), 195-202.
- Liu, Z., Gao, J., Yu, J. (2006) Aflatoxins in stored maize and rice grains in Liaoning Province, China. *Journal of Stored Products Research*, **42**(4), 468-479.
- Lukwago, F., Mukisa, I., Atukwase, A., Kaaya, A., Tumwebaze, S. (2019) Mycotoxins contamination in foods consumed in Uganda: A 12-year review (2006–18). *Scientific African*, **3**, 1-11.
- Lv, C., Jin, J., Wang, P., Dai, X., Liu, Y., Zheng, M., Xing, F. (2019) Interaction of wateractivity and temperature on the growth, gene expression and aflatoxin production by *Aspergillus flavus* on paddy and polished rice. *Food Chemistry*, **293**, 472-478.
- Mahato, D., Lee, K., Kamle, M., Devi, S., Dewangan, K., Kumar, P., Kang, S. (2019). Aflatoxins in food and feed: An overview on prevalence, detection and control strategies. *Frontiers in Microbiology*, **10**, 1-10.
- Majeed, S., De Boevre, M., De Saeger, S., Rauf, W., Tawab, A., Rahman, M., Iqbal, M. (2018) Multiple mycotoxins in rice: occurrence and health risk assessment in children and adults of Punjab, Pakistan. *Toxins*, **10**(2), 1-30.
- Marrez, D.A., Ayeshe, A.M. (2022) Mycotoxins: The threat to food safety. *Egyptian Journal of Chemistry*, **65**(1), 1-2.
- Mahfuz, M., Alam, M.A., Fahim, S.M., Gazi, M. A., Raihan, M.J., Hossain, M., Ahmed, T. (2019) Aflatoxin exposure in children living in Mirpur, Dhaka: Data from MAL-ED companion study. *Journal of Exposure Science and Environmental Epidemiology*, **29**(5), 655-662.
- Milani, J. (2013) Ecological conditions affecting mycotoxin production in cereals: A review. *Veterinari Medicina*, **58**(8), 405–411.
- Miri, Y. B., Belasli, A., Djenane, D., Ariño, A. (2019) Prevention by essential oils of the occurrence and growth of *Aspergillus flavus* and aflatoxin B1 production in food systems. *Aflatoxin B1 Occurrence, Detection and Toxicological Effects*, 1-18. doi: 10.5772/intechopen.88247
- Mitchell, N.J., Hsu, H-H., Chandyo, R.K., Shrestha, B., Bodhidatta, L., Tu, Y.K. (2017) Aflatoxin exposure during the first 36 months of life was not associated with impaired growth in Nepalese children: An extension of the MAL-ED study. *PLoS ONE*, **12**(2), e0172124.
- Mngqawaa, P., Shepharda, G.S., Greenb, I.R., Ngobeni, S.H., de Rijk, T.C., Katerere, D.R. (2015) Mycotoxin contamination of home-grown maize in rural northern South Africa (Limpopo and Mpumalanga Provinces). *Food Additives and Contaminants Part B Surveillance*, **9**(1), 1-17.
- Mohadeseh, N., Faezeh, S., Niloofar, R., Haji-Hoseini, R., Farhad, N., Faramarz, K., Hamidreza, J. (2016) Quantitative determination of aflatoxin by high performance liquid chromatography in wheat silos in Golestan Province, North of Iran. *Iranian Journal of Public Health*, **45**, 905-910.
- Mohd Ikmal, A., Nurasyikin, Z., Tuan Nur Aqlili Riana, T.A., Puteri Dinie Ellina, Z., Wickneswari, R., Noraziyah, A.A.S. (2019) Drought yield QTL (*qDTY*) with consistent effects on morphological and agronomical traits of two populations of new rice (*Oryza sativa*) lines. *Plants*, **8**(6), 1-16.
- Mohd Ikmal, A., Noraziyah, A.A.S., Wickneswari, R. (2021). Incorporating drought and submergence tolerance QTL in rice (*Oryza sativa* L.)—The effects under reproductive stage drought and vegetative

- stage submergence stresses. *Plants*, **10**(2), 1-18.
- Mottaghipour, E., Nazari, F., Hosseini, M.J. (2021) Determination of aflatoxin B1 contamination in wheat and rice flour collected from Iranian market using simple and reliable HPLC method. *Journal of Food and Nutrition Research*, **9**(1), 50-54.
- Mousa, W., Ghazali, F., Jinap, S., Ghazali, H., Radu, S. (2013) Modeling growth rate and assessing aflatoxins production by *Aspergillus flavus* as a function of water activity and temperature on polished and brown rice. *Journal of Food Science*, **78**(1), M56-63.
- Mousavi, S.A.A., Pourtalebi, S. (2015) Inhibitory effects of silver nanoparticles on growth and aflatoxin B1 production by *Aspergillus parasiticus*. *Iranian Journal of Medical Sciences*, **40**(6), 1-6.
- Mtega, M., Mgina, C., Kaale, E., Sempombe, J., Kilulya, K. (2020) Occurrence of aflatoxins in maize and maize products from selected locations of Tanzania and the effects of cooking preparation processes on toxin levels. *Tanzania Journal of Science*, **46**(2), 407-418.
- Mupunga, I., Mngqawa, P., Katerere, D.R. (2017) Peanuts, aflatoxins and undernutrition in children in Sub-Saharan Africa. *Nutrients*, **9**(12), 1287.
- Murashiki, T.C., Chidewe, C., Benhura, M.A., Manema, L.R., Mvumi, B.M., Nyanga, L.K. (2018) Effectiveness of hermetic technologies in limiting aflatoxin B1 and fumonisin B1 contamination of stored maize grain under smallholder conditions in Zimbabwe. *World Mycotoxin Journal*, **11**(3), 459-469.
- Nakavuma, J.L., Kirabo, A., Bogere, P., Nabulime, M.M., Kaaya, A.N., Gnonlonfin, B. (2020) Awareness of mycotoxins and occurrence of aflatoxins in poultry feeds and feed ingredients in selected regions of Uganda. *International Journal of Food Contamination*, **7** (1), 1-10.
- Nazir, A., Kalim, I., Sajjad, M., Usman, M., Iqbal, M. (2019) Prevalence of aflatoxin contamination in pulses and spices in different regions of Punjab. *Chemistry International Journal*, **5**(4), 274-280.
- Nazhand, A., Durazzo, A., Lucarini, M., Souto, E., Santini, A. (2020) Characteristics, occurrence, detection and detoxification of aflatoxins in foods and feeds. *Foods*, **9**(5), 1-26.
- Ndagijimana, R., Shahbaz, U., Sun, X. (2020) Aflatoxin B1 in food and feed: An Overview on prevalence, determination and control tactics. *Journal of Academia and Industrial Research*, **8**(8), 144-152.
- Nerilo, S.B., Romoli, J.C.Z., Nakasugi, L.P., Zampieri, N.S., Mossini, S.A.G., Rocha, G.H.O., Machinski Jr, M. (2020) Antifungal activity and inhibition of aflatoxins production by *Zingiber officinale* Roscoe essential oil against *Aspergillus flavus* in stored maize grains. *Ciência Rural*, **50**, 1-10.
- Nguyen, X.T.T., Nguyen, T.T.T., Nguyen-Viet, H., Tran, K.N., Lindahl, J., Randolph, D.G., Ha, T.M., Lee, H.S. (2018) Assessment of aflatoxin B1 in maize and awareness of aflatoxins in Son La, Vietnam. *Infection Ecology & Epidemiology*, **8**(1), 1-9.
- Oyelami, O.A., Maxwell, S.M., Adelusola, K.A., Aladekoma, T.A., Oyelese, A.O. (1997) Aflatoxins in the lungs of children with kwashiorkor and children with miscellaneous diseases in Nigeria. *Journal of Toxicology and Environmental Health*, **51**(6), 623-628.
- Pankaj, S., Shi, H., Keener, K. (2017) A review of novel physical and chemical decontamination technologies for aflatoxin in food. *Trends in Food Science and Technology*, **71**, 1-12.
- Pandey, M.K., Kumar, R., Pandey, A.K., Soni, P., Gangurde, S.S., Sudini, H.K., Varshney, R.K. (2019) Mitigating aflatoxin contamination in groundnut through a combination of genetic resistance and postharvest management practices. *Toxins*, **11**(6), 1-21.
- Passarelli, S., Bromage, S., Darling, A., Wang, J., Aboud, S., Mugusi, F., et al. (2020) Aflatoxin exposure in utero and birth and growth outcomes in Tanzania. *Maternal and Child Nutrition*, **16**(2), 1-10.
- Peles, F., Sipos, P., Kovács, S., Györi, Z., Pócsi, I., Pusztahelyi, T. (2021) Biological control and mitigation of aflatoxin contamination in commodities. *Toxins*, **13**(2), 1-19.
- Petrović, E., Ćosić, J., Vrandečić, K., Godena, S. (2023) Occurrence of mycotoxins in food and beverages. *Journal of Central European Agriculture*, **24**(1), 137-150.
- Pickova, D., Ostry, V., Toman, J., Malir, F. (2021) Aflatoxins: history, significant milestones, recent data on their toxicity and ways to mitigation. *Toxins*, **13**(6), 1-23.
- Reddy, K., Abbas, H., Abel, C., Shier, W., Oliveira, C., Raghavender, C. (2009) Mycotoxin contamination of commercially important agricultural commodities.

- ToxinReviews*, **28**(2-3), 154-168.
- Riba, A., Bouras, N., Mokrane, S., Mathieu, F., Lebrihi, A., Sabaou, N. (2010) *Aspergillus* section Flavi and aflatoxins in Algerian wheat and derived product. *Food and Chemical Toxicology*, **48**, 2772-2777.
- Roy, S., Shaw, D., Sarkar, T., Choudhury, L. (2023) Mycotoxins in fermented foods: A comprehensive review. *Novel Research in Microbiology Journal*, **7**(2), 1897-1917.
- Rudraboyina, S.K., Kammara, M., Yogita, S.S., Potnuri, H.P.L. (2023) Aflatoxins in cereal crops and their integrated management: A review. *The Pharma Innovation Journal*, **12**(3), 4670-467.
- Sabran, M.R., Jamaluddin, R., Ahmad, Z. (2013) A mini review on aflatoxin exposure in Malaysia: past, present and future. *Frontiers in Microbiology*, **4**, 1-8.
- Saladino, F., Luz, C., Manyes, L., Fernández-Franzón, M., Meca, G. (2016) In vitro antifungal activity of lactic acid bacteria against mycotoxigenic fungi and their application in loaf bread shelf life improvement. *Food Control*, **67**, 273-277.
- Sanders, T.H., Cole, R.J., Blankenship, P.D., Dome, J.W. (1993) Aflatoxin contamination of peanuts from plants drought stressed in pod or root zones. *Peanut Science*, **20**, 5-8.
- Singh, S.K., Patel, M.B., Thakker, B.N., Barad, A.J., Gogoi, R., Hooda, K.S. (2019) An overview of challenges and elimination of aflatoxin contamination in maize (*Zea mays*). *Journal of Pharmacognosy and Phytochemistry*, **8**(4), 2895-2906.
- Sirma, A., Lindahl, J., Makita, K., Senerwa, D., Mtimet, N., Kang'ethe, E., Grace, D. (2018) The impacts of aflatoxin standards on health and nutrition in sub-Saharan Africa: The case of Kenya. *Global Food Security*, **18**, 57-61.
- Somda, M. K., Dabire, Y., Mogmenga, I., Ouattara, A., Nikiema, M., Mihini, H.B., Dicko, M.H. (2023) Assessment of maize contamination by aflatoxin in Burkina Faso: A review of methods of control. *Food and Nutrition Sciences*, **14**(2), 135-147.
- Sowley, E.N.K. (2016) Aflatoxins: A silent threat in developing countries. *African Journal of Biotechnology*, **15**(35), 1864-1870.
- Stanley, J., Patras, A., Pendyala, B., Vergne, M.J., Bansode, R.R. (2020) Performance of a UV-A LED system for degradation of aflatoxins B1 and M1 in pure water: kinetics and cytotoxicity study. *Scientific Reports*, **10**(1), 1-12.
- Tai, B., Chang, J., Liu, Y., Xing, F. (2020) Recent progress of the effect of environmental factors on *Aspergillus flavus* growth and aflatoxins production on foods. *Food Quality and Safety*, **4**(1), 21-28.
- Temba, M.C., Njobeh, P.B., Kayitesi, E. (2017) Storage stability of maize-groundnut composite flours and an assessment of aflatoxin B1 and ochratoxin A contamination in flours and porridges. *Food Control*, **71**, 178-186.
- Thakur, S., Singh, R. K., De, P. S., Dey, A. (2022) Aflatoxins in feeds: Issues and concerns with safe food production. *Indian Journal of Animal Health*, **61**(1), 1-13.
- Thakare, D., Zhang, J., Wing, R.A., Cotty, P.J., Schmidt, M.A. (2017) Aflatoxin-free transgenic maize using host-induced gene silencing. *Science Advances*, **3**(3), e1602382.
- Turksoy, S., Kabak, B. (2020) Determination of aflatoxins and ochratoxin A in wheat from different regions of Turkey by HPLC with fluorescence detection. *Acta Alimentaria*, **49**(1), 118-124.
- Udomkun, P., Wiredu, A., Nagle, M., Bandyopadhyay, R., Müller, J., Vanlauwe, B. (2017) Mycotoxins in Sub-Saharan Africa: Present situation, socio-economic impact, awareness, and outlook. *Food Control*, **72**, 110-122.
- Valencia-Quintana, R., Milić, M., Jakšić, D., Šegvić Klarić, M., Tenorio-Arvide, M.G., Pérez- Flores, G.A., Sánchez-Alarcón, J. (2020) Environment changes, aflatoxins, and health issues, a review. *International Journal of Environmental Research and Public Health*, **17**(21), 7850.
- Vijayalakshmi, S., Nadasabhapathi, S., Kumar, R., Kumar, S.S. (2018) Effect of pH and pulsed electric field process parameters on the aflatoxin reduction in model system using response surface methodology. *Journal of Food Science and Technology*, **55**(3), 868-878.
- Wanniarachchi, P.C., Uthpala, T.G.G., Ranaweera, K.K.D.S. (2023) Aflatoxin occurrence, contamination, detection, and decontamination with special emphasis on coconut oil: A review. *Journal of Agricultural Sciences–Sri Lanka*, **18**(1), 1-28.
- Watson, S., Moore, S., Darboe, M., Chen, G., Tu, Y., Huang, Y., et al. (2018) Impaired growth in rural Gambian infants exposed to aflatoxin: a prospective cohort study. *BMC Public Health*, **18**(1), 1-9.

- Wen, H., Shi, H., Jiang, N., Qiu, J., Lin, F., Kou, Y. (2023) Antifungal mechanisms of silver nanoparticles on mycotoxin producing rice false smut fungus. *Isience*, **26**(1), 105763.
- Wilson, D.M., Stansell, J.R. (1983) Effect of irrigation regimes on aflatoxin contamination of peanut pods. *Peanut Science*, **10**(2), 54-56.
- Wu, F. (2015) Global impacts of aflatoxin in maize: trade and human health. *World Mycotoxin Journal*, **8**(2), 137-142.
- Xu, F., Baker, R.C., Whitaker, T.B., Luo, H., Zhao, Y., Stevenson, A., Zhang, G. (2022) Review of good agricultural practices for smallholder maize farmers to minimise aflatoxin contamination. *World Mycotoxin Journal*, **15**(2), 171-186.
- Xue, W., Macleod, J., Blaxland, J. (2023) The use of ozone technology to control microorganism growth, enhance food safety and extend shelf life: A promising food decontamination technology. *Foods*, **12**(4), 1-24.
- Yáñez, L., Ortiz, D., Calderón, J., Batres, L., Carrizales, L., Mejía, J., Martínez, L., García-Nieto, E., Díaz-Barriga, F. (2002) Overview of human health and chemical mixtures: Problems facing developing countries. *Environmental Health Perspectives*, **110**, 1-9.
- Yan, C., Wang, Q., Yang, Q. and Wu, W. (2020) Recent advances in aflatoxins detection based on nanomaterials. *Nanomaterials*, **10**(9), 1-19.
- Yard, E.E., Daniel, J.H., Lewis, L.S., Rybak, M.E., Paliakov, E.M., Kim, A.A., Sharif, S.K. (2013). Human aflatoxin exposure in Kenya, 2007: A cross-sectional study. *Food Additives & Contaminants: Part A*, **30**(7), 1322-1331.
- Zhang, J., Tang, X., Cai, Y., Zhou, W.W. (2023). Mycotoxin contamination status of cereals in China and potential microbial decontamination methods. *Metabolites*, **13**(4), 1-22.
- Zhao, L., Zhang, L., Xu, Z., Liu, X., Chen, L., Dai, J., Sun, L. (2021) Occurrence of Aflatoxin B₁, deoxynivalenol and zearalenone in feeds in China during 2018–2020. *Journal of Animal Science and Biotechnology*, **12**, 1-12.

حدوث الأفلاتوكسين في الحبوب وآثاره الصحية وإدارته: مراجعة حالية

يوسف أوياميحي⁽¹⁾، نورازيا شمس الدين⁽¹⁾، إسماعيل أدبيبايو⁽²⁾، أولانكا عمر⁽³⁾، نور محمد زيني⁽⁴⁾، محمد إسماعيل⁽⁵⁾

⁽¹⁾ قسم العلوم البيولوجية والتكنولوجيا الحيوية- كلية العلوم والتكنولوجيا - جامعة كيبانغسان ماليزيا- 43600 بانغي- سيلانغور- ماليزيا، ⁽²⁾ قسم علم الأحياء الدقيقة والطبليات والمناعة- كلية الطب- جامعة كابل- كابل- أوغندا، ⁽³⁾ قسم بيولوجيا النبات- كلية العلوم- جامعة إيلورين- إيلورين- ولاية كوارا- نيجيريا، ⁽⁴⁾ قسم علوم الأغذية- كلية العلوم والتكنولوجيا- جامعة كيبانغسان ماليزيا، 43600 بانغي- سيلانغور- ماليزيا، ⁽⁵⁾ مركز بحوث الكيمياء الحيوية التحليلية- جامعة ساينز ماليزيا 11800 (USM)- بولاو بينانغ- ماليزيا.

تستهلك الحبوب على مستوى العالم بسبب قيمتها الغذائية وقدرتها على الحد من سوء التغذية. على الرغم من فائدها، فإن الحبوب عرضة للتلوث بالأفلاتوكسين. الأفلاتوكسينات هي منتجات استقلابية ثانوية شديدة السمية ومسببة لتلوث المنتجات الزراعية التي يستهلكها البشر. أظهرت الدراسات أن الأفلاتوكسين موجود في الحبوب على مستوى عالٍ. يؤدي تعرض الإنسان للأفلاتوكسين من خلال الطعام والأعلاف إلى مجموعة واسعة من المشكلات الصحية، بما في ذلك ضعف جهاز المناعة والسرطان. الأسوأ من ذلك، يمكن أن يسبب الوفاة اعتماداً على مستوى ومدى التعرض. يمكن للعديد من العوامل التي يسببها المناخ، مثل الجفاف، أن تؤدي إلى إنتاج الأفلاتوكسين في جميع أنحاء العالم، وخاصة في إفريقيا، حيث تكون البيئة مواتية. تم اتخاذ العديد من الاحتياطات للتخفيف من التعرض البشري للأفلاتوكسين، بما في ذلك اللوائح الصارمة، ومنع التلوث قبل وبعد الحصاد، وإزالة السموم، وإزالة التلوث. وبالإضافة إلى ذلك، يمكن أن تساعد الإدارة والممارسات الجيدة للمزارع، وخلق الوعي والتثقيف في الحد من التلوث بالأفلاتوكسين والتعرض له، إن لم يكن القضاء عليه. وقدم الاستعراض الحالي تفاصيل عن أحدث المعلومات عن حدوث الأفلاتوكسين، والآثار الصحية، وتدابير الرقابة على الأفلاتوكسين في الحبوب لضمان سلامة الأغذية ورفاهية الإنسان.