Knowledge, Attitude, and Practice of Blood Donation among Najran population Kingdom of Saudi Arabia

Amna Mohammed Idris 1, Zeinab Taha Ali Omer²

¹ Assistant Professor in critical care nursing/College of Nursing/Medical Surgical Nursing department, Najran University, Kingdom of Saudi Arabia

²Assistant Professor Medical-Surgical Nursing, Nursing Department, Faculty of Applied Medical Sciences, Al-Baha University, Saudi Arabia: zomer@bu.edu.sa

Abstract

Aim: This study aimed to assess the knowledge, attitude, and practice of the Najran population about blood donation in Saudi Arabia. Methods: A descriptive study was carried out in Najran city to assess people's knowledge, attitude, and practice about donating blood. The study enrolled three hundred and three participants, convenience sampling was used to select participants. Data was collected by questionnaires and analyzed by SPSSResults: There were 303 participants, the majority of whom were female, and 87.5 percent of them had intermediate knowledge regarding blood donation. The majority of participants strongly agreed that they have a good attitude regarding blood donation when asked about it. Additionally, there was no statistical correlation between the research sample's demographic characteristics such as age, social status, education nationality, and knowledge of blood donation, but there was a statistical correlation between these characteristics and their attitude toward blood donation. Conclusion Indicates that the majority of participants had intermediate knowledge levels, and a favorable attitude toward blood donation and 40% of them had experienced an episode of anemia in the past. In addition, 23% of people simply stated that they had donated in the past. Recommendations: The findings showed the critical need for efficient health education that aimed to inform individuals about blood donation and its effects. There should be more research done on a broader population using more samples.

Keywords Najran, The Kingdom of Saudi Arabia, Knowledge, Attitude, Practice, Blood donation

Introduction

Because blood is a key component of the human body that can only be produced naturally through human resources, blood donation has become a procedure that every healthcare establishment consider.(Alharbi et al., 2018) Globally, and particularly in developing nations, the demand for safe and sufficient blood transfusions is rising to meet medical needs and treat clinical conditions. This is primarily because of rising rates of traffic accidents and transfusiondependent hematological disorders like sickle cell anemia, thalassemia, and nephrotic syndrome.(Hamali, 2018) Both developed and developing countries are now finding it difficult to enroll donors to donate at regular times per year. Around 118 million units of blood are gathered from donors each year all around the world. Children under the age of five received more than 50% of blood transfusions in

underdeveloped nations, whereas patients over the age of 60 accounted for up to 76% of all transfusions in wealthy nations. Blood donors can be classified as family replacement, paid, or compensated, or voluntary unpaid donors. Regular, unpaid, voluntary blood donors provide blood that is trustworthy, safe, and sufficient. (World Health Organization, 2023) Additionally, encouraging younger individuals to donate blood voluntarily as the quantity, quality, and safety of blood products have increased.(Hindawi S, 2020) Several variables affected the willingness and recruitment of individuals to give their blood willingly. Numerous studies have demonstrated a correlation between the age, sex, and educational level of the donor population and these demographics.(Alnouri et al., 2019).

1.2 Problem Statement

Blood donation is currently the only method to receive blood and its components. The scarcity of blood in medical institutions is a serious concern for any society. Donor attitudes, knowledge,

motives, and beliefs impact the process of donating blood as well as the number of donors that are now actively involved. (Hamali, 2018)

1.3 Rationale and Justification

The Saudi public's blood donation prevalence was below expectations, most likely as a result of misunderstandings, a lack of awareness, and a negative attitude toward donation. Educational initiatives are required to raise Saudi citizens' awareness of blood donation and to change their attitudes regarding it. (Alharbi et al., 2018).

Aim of the study:

This study aims to assess the knowledge, attitude, and practice of the Najran population about blood donation in the Kingdom of Saudi Arabia.

Research question:

The researcher answered the question: What is people's knowledge about blood donating? What is the people's attitude regarding blood donation? What is people's practice regarding blood donation? &What is the relationship between demographic data and public knowledge about blood donation in Najran City, Kingdom of Saudi Arabian.

Materials and methods:

Study design:

The study was done by using a descriptive community-based study design to assess the knowledge, attitude, and practice of the Najran City population about blood donation.

Study area:

The study was conducted in Najran City. It is in southwestern Saudi Arabia near the border with Yemen. It is the capital of Najran Province.

Study population:

adults aged eighteen years older or above both genders. from the Najran City community.

Inclusion criteria

An adult member of the Najran community whose age is eighteen years older or above both genders and agreed to join in the study.

Exclusion criteria

Any member of the Najran community who age is less than eighteen years old.

Sampling procedure and sample size

A convenience sample method was employed to choose participants from the Najran City Community. In this study, three hundred and three members of the Najran community joined up. For convenience. The sample size was

calculated using the Sample Size Calculator. Confidence Level = 95%., Acceptable error = 5%. Expected frequency =50%.

Data collection techniques:

Data was gathered by distributing to the Najran City population the created questionnaire and outlining the goal of the research.

Data collection tool:

Data was gathered. Data was gathered using an automated questionnaire linked to social media once the study's purpose was communicated to the Najran community. The questionnaire consists of Five parts: the first portion contains demographic information. Personal data, including age, gender, qualification, and occupation, was gathered in this area. Blood donation is covered in part two. Section three concerns the history of chronic disease in order to assess the patient's clinical background; Section four concerns participants' knowledge of blood donation; and Section Five concerns attitudes toward blood donation. The participants' responses were measured using a Likert scale. Scheme grading: 70% or above is considered high, 50% to 70% is considered intermediate, and 50% is considered low.

Data Analysis:

Version 24 of the Statistical Package for Social Sciences (SPSS) was used to handle and analyse the data. descriptive statistics were used to show the data as frequencies, percentages correlation coefficients utilized. A threshold of 0.05 was used to evaluate statistical significance.

Ethical consideration:

Participants' rights were respected throughout the investigation since they were made aware that their participation in the study was voluntary that the results would only be used for research reasons, and that they could withdraw their consent at any time before completing the online survey. The Dean of the Faculty of Nursing Sciences at Najran University granted permission.

Tool reliability was evaluated using the Cronbach-Alpha Coefficient statistical test, which produced a reliability score of (r=0.77).

Pilot research:

To test the tools' applicability, clarity, and feasibility, a pilot study involving 10% of the study sample was performed. The final form had been established when all of the modifications were completed. The study group did not include the person from the pilot research.

Results summary:

303 people in total voluntarily took part in the study. According to Table 1: the participants were mostly between the ages of 18 and 28 years (54.5%), were mostly Saudi (75.6%), had more than half female genders, and had only received a university degree in 49.8% of cases. Less than half of the participants (43.2%) were single, and 26.4% of the participants were employed.

Table 2 illustrates that 23% of participants gave blood.

Table 3: reveals that 23% of the participants have a history of chronic illnesses. 40% of individuals have a history of anemia.

Table 4: demonstrate people's knowledge of donating blood with a total mean and standard deviation (1.9 ± 0.8) .

Figure 1: demonstrates that the majority of participants (87.5%) had intermediate knowledge regarding blood donation, followed by (11.2%) who had high knowledge.

Figure 2: demonstrates that the majority of participants (72.6%) had a positive attitude toward donating blood, whereas 27.40% had a negative attitude about it.

Table 5: showed that the participants provided attitude about blood donation with a mean total and standard deviation (3.6 ± 1.3) .

Table 6: demonstrates that there was no statistically significant connection. between the research sample's demographic characteristics such as age, height, nationality, sex, and social class, and the participants' knowledge of blood donation.

According to Table 7, there was a statistical relationship between the age, social status, educational attainment, nationality of the research sample, and attitude level toward blood donation. However, there was no statistical relationship between the demographic characteristics and the attitude level toward blood donation for the following variables: height, sex, and job.

Tables:

Table 1: Demographic characteristics of the Participants in Najran City

|] | F | % | |
|-------------------|------------------------|-----|-------|
| Deuticination | Yes | 303 | 100% |
| Participation | No | 0 | 0 |
| | 18 to 28 years | 165 | 54.5% |
| Age | 29 to 39 years | 86 | 28.4% |
| | 39> years | 52 | 17.2% |
| | 140-150 | 50 | 16.5% |
| Height | 155-165 | 161 | 53.1% |
| _ | >165 | 92 | 30.4% |
| Nietienelitee | Saudi | 229 | 75.6% |
| Nationality | Non-Saudi | 74 | 24.4% |
| S | Male | 98 | 32% |
| Sex | Female | 205 | 67.7% |
| | read and write | 19 | 6.3% |
| Educational level | Intermediate education | 41 | 13.5% |
| Educational level | high school education | 92 | 30.4% |
| | University education | 151 | 49.8% |
| | Married | 124 | 40.9% |
| Social status | Single | 131 | 43.2% |
| Social status | Widower | 37 | 12.2% |
| | Divorced | 11 | 3.6% |
| | Work | 80 | 26.4% |
| Job | do not work | 108 | 35.6% |
| | Student | 115 | 38.0% |

Table 2: The percentage of participants Donating of blood in Najran City

| Items | F | % | |
|----------------|-----|-----|-------|
| Blood donation | Yes | 70 | 23.1 |
| | No | 233 | 76.9% |

Table 3: The Medical history of participants in Najran City

| Items | Yes | No | Total |
|---------------------------------|-----|-----|-------|
| Previous history of any chronic | 23% | 77% | 100% |
| diseases | | | |
| History of anemia | 40% | 60% | 100% |

Table 4: The participants knowledge about blood donating in Nairan City

| able 4: The participants knowledge about blood donating in Najran City | | | | | | | |
|--|------|-------------------|--|--|--|--|--|
| L Items | Mean | Std. Deviation | | | | | |
| 1 1-Do you think that donating blood is healthy? | 2.40 | 0.78 | | | | | |
| 2 2-Does donating blood cause infection? | 1.91 | 0.80 | | | | | |
| 3 3- Can a pregnant woman donate blood? | 1.61 | 0.78 | | | | | |
| 4 4-Can a person with chronic diseases donate blood? | 1.58 | 0.76 | | | | | |
| 5– Is it possible for a menstruating woman to donate blood? | 1.66 | 0.74 | | | | | |
| 6 6- Is there a certain age for donating blood? | 2.07 | 0.86 | | | | | |
| 7 7 - Can a breastfeeding woman donate blood? | 1.79 | 0.77 | | | | | |
| 8 - When people need a blood transfusion, do they have to pay? | 1.50 | 0.75 | | | | | |
| 9 10- Is blood from one donor sufficient for one needy person? | 1.81 | 0.77 | | | | | |
| 10 11-Does donating blood make you lose or gain weight? | 1.62 | 0.73 | | | | | |
| 11 12-Can people under 16 years of age donate blood? | 1.66 | 0.77 | | | | | |
| 12 13-Does donating blood help patients who need it? | 2.36 | 0.84 | | | | | |
| 13 14-Can people of any blood type donate blood? | 2.32 | 0.83 | | | | | |
| 15- If the donor develops a fever on the day of donation, can he donate blood? | 1.59 | 0.77 | | | | | |
| 16-Do you think community awareness helps with blood donation opportunities? | 2.32 | 0.84 | | | | | |
| 16 17-Can a smoker donate blood? | 1.83 | 0.82 | | | | | |
| Total knowledge | 1.9 | 0.8 | | | | | |

Figure 1: The level of participants knowledge about donating blood. (n=303)




Figure 2: The participants attitude towards blood donation. (n=303)




Table 5: The participants attitudes about blood donation.

| NO | Attitude | Mean | Std. Deviation |
|----|--|------|-------------------|
| 1 | My donation will encourage others to donate. | 4.03 | 0.98 |
| 2 | I will donate blood if it is confirmed that the donated blood will be delivered to one of my family members in the future. | 3.35 | 1.44 |
| 3 | I will donate blood if there are incentives or rewards. | 3.06 | 1.41 |
| 4 | Donating blood saves lives. | 3.96 | 1.27 |
| 5 | Donating blood is an ethical activity. | 3.69 | 1.33 |
| 6 | Young people should donate frequently. | 3.57 | 1.29 |
| 7 | People who have greater knowledge of donating blood donate more often. | 3.80 | 1.23 |
| 8 | The best way to donate blood is to volunteer without payment. | 3.85 | 1.23 |
| 9 | Do you agree to pay money to the donor? | 2.99 | 1.43 |
| 10 | Donating blood is important. | 3.96 | 1.21 |
| | Total | 3.6 | 1.3 |

Table 6: The relationship between the demographic characteristics of the research sample and the participants knowledge about donating blood:

| | | | Level of knowledge | | | | | | | |
|-------------------------------|------------------------|-------|--------------------|------|--------------------|----|---------------------|------|--|--|
| Demographical characteristics | | I don | I don't know | | I know somewhat | | I know perfectly | | | |
| | F | % | F | % | F | % | | | | |
| Age | 18 to 20 years | 21 | 61.8 | 141 | 53.2 | 3 | 75 | | | |
| | 29 to 39 years | 8 | 23.5 | 77 | 29.1 | 1 | 25 | 0.75 | | |
| | 39> years | 5 | 14.7 | 47 | 17.7 | 0 | 0 | | | |
| height | 140-150 | 9 | 26.5 | 39 | 14.7 | 2 | 50 | | | |
| | 155-165 | 18 | 52.9 | 142 | 53.6 | 1 | 25 | 0.13 | | |
| | >165 | 7 | 20.6 | 84 | 31.7 | 1 | 25 | | | |
| Nationality | Saudi | 25 | 73.5 | 201 | 75.8 | 3 | 75 | 0.06 | | |
| | Non-Saudi | 9 | 26.5 | 64 | 24.2 | 1 | 25 | 0.96 | | |
| Sex | Male | 11 | 32.4 | 17.7 | 92.9 | 28 | 75.7 | 0.05 | | |
| | Female | 23 | 67.6 | 86 | 32.5 | 1 | 24.3 | 0.95 | | |
| Educational | read and write | 2 | 5.9 | 17 | 6.4 | 0 | 0 | | | |
| level | Intermediate education | 4 | 11.8 | 37 | 14 | 0 | 0 | | | |
| | high school education | 12 | 35.3 | 79 | 29.8 | 1 | 25 | 0.94 | | |
| | University education | 16 | 47.1 | 132 | 49.8 | 3 | 75 | | | |
| Social status | Married | 13 | 38.2 | 109 | 41.1 | 2 | 50 | | | |
| | Single | 16 | 47.1 | 113 | 42.6 | 2 | 50 | 0.02 | | |
| | Widower | 3 | 8.8 | 34 | 12.8 | 0 | 0 | 0.93 | | |
| | Divorced | 2 | 5.9 | 9 | 3.4 | 0 | 0 | 1 | | |
| Job | Work | 6 | 17.6 | 74 | 27.9 | 0 | 0 | | | |
| | Don't work | 12 | 35.3 | 95 | 38.5 | 1 | 25 | 0.31 | | |
| | Student | 16 | 47.1 | 96 | 36.2 | 3 | 75 | | | |

Table 7: The relationship between the demographic characteristics of the research sample

and the level of participants attitude about blood donation:

| Level of attitude | | | | | | | | | | | | |
|-------------------------------|-----------------------|------------------|-----|-----------------|------|--------------------|------|--------|------|-------------------------|------|-----------------|
| Demographical characteristics | | I don't o kno | - | I don't know | | I know somewhat | | I know | | I know perfectl y | | P- Valu e |
| | | | % | F | % | F | % | F | % | F | % | |
| | 18 to 20 years | 0 | 0 | 11 | 26.2 | 84 | 57.5 | 55 | 57.3 | 15 | 83.3 | |
| Age | 29 to 39 years | 0 | 0 | 20 | 47.6 | 36 | 24.7 | 27 | 28.1 | 3 | 16.7 | 0.00 |
| | 39> years | 1 | 100 | 11 | 26.2 | 26 | 17.8 | 14 | 14.6 | 0 | 0 | |
| 1 1. | 140-150 | 0 | 0 | 7 | 16.7 | 25 | 17.1 | 15 | 15.6 | 3 | 16.7 | |
| heigh t | 155-165 | 1 | 100 | 18 | 42.9 | 83 | 56.8 | 50 | 52.1 | 9 | 50 | 0.79 |
| ' | >165 | 0 | 0 | 17 | 40.5 | 38 | 26 | 31 | 32.3 | 6 | 33.3 | |
| Nati | Saudi | 0 | 0 | 23 | 54.8 | 114 | 78.1 | 75 | 78.1 | 17 | 94.4 | |
| onali ty | Non-Saudi | 1 | 100 | 19 | 45.2 | 32 | 21.9 | 21 | 21.9 | 1 | 5.6 | 0.00 |
| Sex | Male | 0 | 0 | 16 | 38.1 | 43 | 29.5 | 33 | 34.4 | 6 | 33.3 | 0.76 |
| Sex | Female | 1 | 100 | 26 | 61.9 | 103 | 70.5 | 63 | 65.6 | 12 | 66.7 | |
| | read and write | 0 | 0 | 4 | 9.5 | 13 | 8.9 | 2 | 2.1 | 0 | 0 | 0.00 |
| Educa | | 0 | 0 | 13 | 31 | 20 | 13.7 | 8 | 8.3 | 0 | 0 | |
| tional level | high school education | 0 | 0 | 11 | 26.2 | 50 | 34.2 | 29 | 30.2 | 2 | 11.1 | |
| | University education | 1 | 100 | 14 | 33.3 | 63 | 43.2 | 57 | 59.4 | 16 | 88.9 | |
| | Married | 1 | 100 | 18 | 42.9 | 61 | 41.8 | 40 | 41.7 | 4 | 22.2 | 0.01 |
| Social | Single | 0 | 0 | 10 | 23.8 | 59 | 40.4 | 48 | 50 | 14 | 77.8 | |
| status | Widower | 0 | 0 | 11 | 26.2 | 19 | 13 | 7 | 7.3 | 0 | 0 | |
| | Divorced | 0 | 0 | 3 | 7.1 | 7 | 4.8 | 1 | 1 | 0 | 0 | |
| | Work | 1 | 100 | 13 | 31 | 33 | 22.6 | 29 | 30.2 | 4 | 22.2 | 0.16 |
| Job | Don't work | 0 | 0 | 19 | 45.2 | 50 | 34.2 | 35 | 36.5 | 4 | 22.2 | |
| | Student | 0 | 0 | 10 | 23.8 | 63 | 43.2 | 32 | 33.3 | 10 | 55.6 | |

Discussion:

the Najran population about blood donation. The results of the current study were consistent with those of a study conducted by Kanani in India, which revealed that the majority of participants (80.2%) had never donated blood. (Kanani AN, 2018) In contrast, research conducted in Ethiopia by Aschale revealed that the majority of respondents (83.7%) had a history of blood

[.] A person's life can be saved by a blood transfusion in a variety of acute and chronic diseases. if they have blood loss as a result of an accident, surgery, or other medical issues (Thurn et al., 2019) This study's goal is to assess the knowledge, attitude, and practice of

donation. (Aschale et al., 2021)

In terms of the participants' medical histories, 23% of the participants have a history of chronic diseases. Among the participants, anemia has a (40%) history. Most participants do not donate blood because of this. The survey found that most participants had moderate levels of understanding about blood donation. This outcome is consistent with research conducted by Melku in Ethiopia, which found that most participants (79.2%) had a sufficient understanding of blood donation(Melku et al., no date) Additionally, a different study conducted in Saudi Arabia by Alsalmi revealed that the majority of participants had enough awareness about blood donation.(Alsalmi et al., 2019) On the other hand, research by Baseer found that the majority of participants knew little to nothing about blood donation.(Baseer et al., 2017) Also study done by Tan, found that most participants had a lack of awareness about blood donation. (T, Hafizuddin and Rosnah, 2019)

The present study found that most participants had favorable beliefs toward blood donation. This outcome was consistent with a study performed by Alharbi, which found that nearly all the participants had a favorable attitude toward blood donation. (Alharbi et al., 2018) A survey executed by Melku also revealed that almost all the participants (79.2%) had positive beliefs toward blood donation. (Melku et al., no date) on the contrary, Baseer's study found that the majority of participants had negative attitudes concerning blood donation. (Baseer et Alsalmi's study additionally $2017)^{1}$ discovered that a larger proportion of individuals had a negative attitude toward blood donation.(Alsalmi et al., 2019)

Our study discovered that there was no statistically significant relationship between the demographic characteristics of the research sample such as age, height, nationality, sex, the people's knowledge of blood donation. This outcome is consistent with research by Enawgaw and Malako in Ethiopia in terms of age and marital status.(Enawgaw, Yalew and Shiferaw, 2019)(Malako, Yoseph and Bekele, 2019) .In contrast, a study carried out in India by Razdan identified a statistical relationship age, educational attainment, and socioeconomic

position, and their understanding of blood donation.(Razdan et al., 2018) The current study revealed a statistical relationship between the age, status, educational attainment, nationality of the research sample and the attitude level toward blood donation; however, there was no statistical relationship between the attitude level toward blood donation and the relationship between height, sex, or employment. Similarly, a study conducted in Ethiopia by Enawgaw revealed a statistical relationship between the demographic characteristics of the research sample and the attitude level toward blood donation. (Enawgaw, Yalew and Shiferaw, 2019) On the other hand, a study by Gill found a statistical relationship between sex and attitude toward blood donation.(Gill, 2021)

Conclusion:

The study concluded that the majority of individuals in the study had never given blood, according to the findings. While the majority of respondents had a good attitude and just a moderate amount of understanding regarding blood donation. The demographic traits of persons, such as age, height, nationality, sex, social position, degree of education, and employment, did not statistically correlate with their knowledge of donation. Furthermore, blood a statistical association was observed between demographic attributes of the research sample, including age, social status, educational attainment, and the attitude level toward blood donation.

Recommendations:

The findings demonstrated the urgent need for effective health education that focused educating people about blood donation and its consequences. More research should be done with a larger sample size, across the nation, and with different methods of study.

References:

social status, educational attainment, job, and ALHARBI, S. ALATEEQ ,F., AHMED ,I ., **ALSOGAIR AL-RASHDI** Y, A., T.& ALDUGIEMAN. Ahmed.H.G. 2018. Assessment of Levels of Awareness Towards Blood Donation in Saudi Arabia J. AIMS Public Health .;5(May), pp. 324–337. doi: 10.3934/publichealth.2018.3.324.

between participant demographics, including ALNOURI ,A.K., MAGHRABI, L.A., HAMDI, S.S., EL-GHANY, S.M.A., &ALNOURI,

- Analysis of the K.A. 2019. Common Causes of Blood Donor Deferral in Northern Jeddah: a single-center study. Journal of Blood Medicine. pp. 47–51.
- ALSALMI, M., ALMALKI, H., ALGHAMDI, Attitude and Practice of Blood Donation among Health Professions Students in Saudi Arabia; A cross-sectional study. Journal. Family Medicine and Primary MELKU, 2322–2327.doi: Care. 7. pp. 10.4103/jfmpc.jfmpc.
- BASEER, S., EJAZ, S., NOORI, M. & FAISAL A. 2017 .'Knowledge , Attitude Perceptions of University Students towards Blood Donation: An Assessment of Motivation and Barriers. Isra Med J. 9(6): 406-10.
- Enawgaw, B., Yalew, A. & Shiferaw E. 2019. Blood Donors knowledge and Attitude towards Blood Donation at North Gondar districtBloodBank, Northwest Ethiopia: a cross - sectional study BMC Research Notes. pp. 1–6. doi: 10.1186/s13104-019-4776-0.
- Gill ,N.P. 2021. The Statistical Analysis of Factors Explaining the Intention of Public for Blood Donation in Jamshoro, Sindh . Journal of Business and Social Review in Emerging Economies.;7(2), pp. 231–238.
- , Attitude Hamali, Н. 2018. knowledge Awareness, and Motivation of Blood Donation Among Academic Staff in Health Colleges in The Southern Region of Saudi Arabia . Journal of Health Sciences;3(2):1–7.
- ASCHALE, A., FUFA, D., KEKEBA, T& BIRHANU ,Z. 2021. Intention to voluntary blood donation among private higher education students, Jimma town, Oromia, Ethiopia: Application of the theory of planned behaviour. pp. 1–14. doi: 10.1371/journal.pone.0247040.
- Kanani, A. A. 2018 Study on Knowledge and Awareness about Blood Donation amongst Government Medical, Para-Medical and Nursing Undergraduate Students in Jamnagar, Gujarat . Global 46–51. doi: 3(1),pp.10.4103/GJTM.GJTM.
- Yoseph ,F. Malako, D., &Bekele, M.L.

- 2019. Assessment of knowledge, attitude and practice and associated factors of blood donation among health care workers in Ethiopia: a cross-sectional study. BMC Hematology. 19:10 pp. 1–8.
- A. &ALJASIR, B. 2019. Knowledge, HINDAWI, S. 2020. Evolution of Blood Transfusion Medicine in Saudi Arabia. TRANSFUSION;60(S1):S2–S3.doi: 10.1111/trf.15681.
 - E., M., ASRIE, F., SHIFERAW, WOLDU, B., ASMELASH, & ENAWGAW, B. 2018. Knowledge, Attitude and Practice Regarding Blood Donation among Graduating Undergraduate Health Science Students at the University of Gondar , Northwest Ethiopia. Ethiop J Health Sci.28(5):571
 - RAZDAN, N., KUMAR, R., BHARDWAJ, A. & SINGH, P. 2018. The knowledge, attitude, and practice toward blood donation among voluntary and replacement blood donors in Ambala , India. International Journal of Medical Science and ALHARBI, ALATEEQ ,F., AHMED ,I ., ALSOGAIR A., AL-RASHDI Y, ALDUGIEMAN, T.& Ahmed,H.G. 2018. Assessment of Levels of Awareness Towards Blood Donation in J. Saudi Arabia AIMS PublicHealth .;5(May), 324–337. doi: pp. 10.3934/publichealth.2018.3.324.
 - ALNOURI ,A.K., MAGHRABI, L.A., HAMDI, S.S., EL-GHANY, S.M.A., &ALNOURI, K.A. 2019. Analysis of the Most Common Causes of Blood Donor Deferral in Northern Jeddah: a single-center study. Journal of Blood Medicine. pp. 47–51.
 - ALSALMI, M., ALMALKI, H., ALGHAMDI, A. &ALJASIR, B. 2019. Knowledge, Attitude and Practice of Blood Donation among Health Professions Students in Saudi Arabia; A cross-sectional study. Journal. Family Medicine and Primary Care. 7. pp. 2322-2327.doi: 10.4103/jfmpc.jfmpc.
 - BASEER, S., EJAZ, S., NOORI, M. & FAISAL A. 2017 . 'Knowledge', Attitude, Perceptions of University Students towards Blood Donation: An Assessment of Motivation and Barriers. Isra Med J. 9(6): 406-10.
- Journal of Transfusion Medicine AATM. Enawgaw, B., Yalew, A. & Shiferaw E. 2019. Blood Donors knowledge and Attitude towards Blood Donation North at Gondar districtBloodBank, Northwest Ethiopia: a

- cross sectional study BMC Research 4776-0.
- Gill ,N.P. 2021. The Statistical Analysis of Factors Explaining the Intention of Public for Blood Donation in Jamshoro, Sindh . Emerging Economies.;7(2), pp. 231–238.
- Hamali, H. 2018. knowledge, Attitude, Awareness, and Motivation of Blood Donation Among Academic Staff in Health Colleges in The Southern Region WORLD of Saudi Arabia . Journal of Health Sciences;3(2):1–7.
- ASCHALE, A., FUFA, D., KEKEBA, T& BIRHANU ,Z. 2021. Intention to voluntary blood donation among private higher education students, Jimma town, Oromia, Ethiopia: Application of the theory of planned behaviour. pp. 1–14. doi: 10.1371/journal.pone.0247040.
- Kanani, A. A. 2018 Study on Knowledge and about Blood Donation Awareness amongst Government Medical, Para-Medical and Nursing Undergraduate Students in Jamnagar, Gujarat . Global Journal of Transfusion Medicine AATM. 46–51. 3(1),pp.doi: 10.4103/GJTM.GJTM.
- Malako, D., Yoseph ,F. &Bekele, M.L. 2019. Assessment of knowledge, attitude and practice and associated factors of blood donation among health care workers in Ethiopia: a cross-sectional study. BMC Hematology. 19:10 pp. 1–8.
- HINDAWI ,S. 2020. Evolution of Blood Transfusion Medicine in Saudi Arabia. TRANSFUSION;60(S1):S2–S3.doi: 10.1111/trf.15681.
- MELKU, M., ASRIE, F., SHIFERAW, E., WOLDU, B., ASMELASH, D. & ENAWGAW,B. 2018. Knowledge Attitude and Practice Regarding Blood Donation among Graduating Undergraduate Health Science Students at the University of Gondar, Northwest Ethiopia. Ethiop J Health Sci.28(5):571
- RAZDAN, N., KUMAR, R., BHARDWAJ, A. & SINGH, P. 2018. The knowledge attitude, and practice toward blood donation among voluntary replacement blood donors in Ambala, India. International Journal of Medical

- Science and Public Health. 7(6) pp. 444–452. Notes. pp. 1–6. doi: 10.1186/s13104-019- (January). doi: 10.5455/jjmsph.2018.0307109032018.
 - T, P. P., HAFIZUDDIN, M. F. & ROSNAH, B. (2019). 'Knowledge and Perceptions of Blood Safety among Blood Donors in Kelantan, Malaysia.26(4), pp. 127–136.
- Journal of Business and Social Review in THURN, L., WIKMAN, A., WESTGREN, M. & LINDOVIST, P.G. 2019. 'Incidence and risk factors of transfusion reactions in postpartum blood transfusions.3(15). 10.1182/bloodadvances.2019000074.
 - HEALTH ORGANIZATION. 2023. Blood safetv and availability. In https://www.who.int/news-room/factsheets/detail/blood-safety-and-availability.