#### SINAI Journal of Applied Sciences 12 (6) 2023 915-930



Available online at <u>www.sinjas.journals.ekb.eg</u> SCREENED BY SINAI Journal of Applied Sciences Vint ISSN 2314-6079 Online ISSN 2682-3527



# EFFECT OF DIFFERENT TYPES OF ORGANIC FERTILIZERS AND PLANTING DATES ON THE PRODUCTIVITY OF SEEDS, ROOTS AND THEIR MUCILAGE CONTENT OF Althea rosea PLANTS

Samah, A. Abo-Baker<sup>\*;</sup> Dina A. Soliman and M.A. Mahmoud

Dept. Plant Prod., Fac. Environ. Agric. Sci., Arish Univ., Egypt.

### ARTICLE INFO

#### ABSTRACT

Article history: Received: 25/11/2023 Revised: 02/12/2023 Accepted: 30/12/2023

Keywords: *Althea rosea*, organic fertilizers, vermicompost, chicken manure, compost, mucilage.



2022-2023 at the Experimental Farm of Ornamental and Medicinal Plants, Faculty of Environmental Agriculture Science at El-Arish, Arish University, North Sinai, to study the effect of three types of organic fertilizers, *i.e.*, chicken manure (ChM), vermicompost (VC), compost (C) and there mixture as well as three sowing dates were as October  $15^{\text{th}}$ , November  $1^{\text{st}}$  and November  $15^{\text{th}}$  for both seasons on growth, seeds, roots and mucilage yields of *Althea rosea* plant. The result showed that the highest values of all growth parameters: plant height, number of leaves and fresh and dry weight of leaves recorded at the first sowing date October  $15^{\text{th}}$  with chicken manure fertilization at (5 t/fad.) plus of vermicompost (3 t/fad.). In addition, the highest increase in root length, fresh and dry weight of root, yield of root and seeds, percentage and yield of mucilage in root and seeds achieved in early sowing date mid-October and mix of vermicompost and chicken manure.

A field experiment was carried out during winter seasons of 2021-2022 and

# **INTRODUCTION**

The Malvaceae family includes a plant known as garden hollyhock or marshmallow (Althea rosea). This plant is native to China, Southern Europe, the Middle East, the Mediterranean, and Central Asia. The stem, which is either erect or sparsely branched, has stellate hairs, and can reach a height of 0.5-2.0 meters. The seeds receive coverage from stellate hairs as well (Aslam et al., 2012). Althea rosea has a history of traditional use in the treatment of various health conditions as managing fever. respiratory issues, gastrointestinal problems, sore throat, and skin burns. Additionally, it demonstrated effectiveness against has bacterial infections and inflammation (Nawaz et al., 2022). The economic part of this plant is the root because it contains vitamin C (26.80 mg/100g), crud protein (7.49%), crud fibre (3.28%), ash (5.28%)

and a high percentage of polysaccharides (83.94%) represented by starch (27.16%), pectin (10.22%) and mucilage (16%) which is considered the most important active substance in the plant.

Mucilage is synthesized in certain plants during their growth and its primary function inside the plant is to provide water, nutrients, and promote seed germination. The main sources for producing mucilage in the Althea rosea plant are the roots (20%), seeds (9%), leaves (4.3%), and flowers (1.6 MDa) (Tomoda et al., 1985; Classen and Blasheck, 1998; Niknam, 1999; Detersa et al., 2010). Mucilage is playing a crucial role in various medical applications in addition used in the food industry, as natural preservatives in juices and jams are contributed to improving the stability of some nutritional components while preserving their quality over time makes them a healthier option

\* **Corresponding author: E-mail address:** bakr@agri.aru.edu.eg https://doi.org/ 10.21608/sinjas.2023.260784.1246

2023 SINAI Journal of Applied Sciences. Published by Fac. Environ. Agric. Sci., Arish Univ. All rights reserved..

### than synthetic preservatives (Amini and Razavi, 2012; Husain *et al.*, 2019; Al Masum *et al.*, 2020; Maher, 2022).

Sowing date is the most effected on the vegetative and reproductive phases, as well as on crop productivity and quality. The quality and components of chemicals within the plants are greatly affected by the day length, temperature, and relative humidity (Ghasemi et al., 2020). The highest amount of mucilage and yield was achieved when cultivating okra early (Rad et al., 2022). Early cultivation of the soybean crop also increased the number of pods in each plant and weighed 1,000 seeds and higher protein content in the seeds than late cultivation of soybean (Jarecki and Bobrecka-Jamro, 2021). Additionally, Overdue Agriculture was the cause of low productivity in the jute crop (Saha et al., 2015).

Non-target organisms, biological ecosystems, and soil microorganisms have been affected by the intensive use of chemical fertilizers (Maddahi et al., 2021). So, the current chemical application systems are replaced by organic fertilization, which enhances the soil's chemical, physical, and biological properties while preserving its ability to retain moisture by increasing the amount organic matter of in it. Consequently, crop growth is enhanced, and its quality and quantity are maintained. (Ofosu-Anim et al., 2006; Heidarzadeh et al., 2022; Prisa and Testa, 2022). Organic fertilizers such as vermicompost, compost, and chicken manure are considered environmentally friendly ways to enhance soil fertility and production, which reduces the need for chemical fertilizers (Khosravi Shakib et al., 2019; Maddahi et al., 2021). Animal manure can be used as organic sources of nutrients to achieve sustainable plant production. Improving soil fertility, structure, and water and air permeability of soils is achieved by increasing organic matter, which in turn enhances host plant growth and yield (Rahimi et al., 2019).

Compost is a process that uses aerobic or anaerobic fermentation to turn organic waste into a stable form (**Talyan** *et al.*, **2008**), its purpose is to enhance the soil's chemical and physical characteristics. The soil is made easier to grow by promoting the growth of roots and providing nutrients to the plant, increasing its potential for the roots to absorb them. Additionally, the compost aids in keeping moisture in the soil (Lea Master *et al.*, **1998**).

is of Vermicompost made up microorganisms and earthworms, which are interacting to produce nutrient-rich organic matter. Sustainable agriculture could benefit from using it to enhance soil porosity, which would increase nutrient availability. Vermicompost contains microscopic organisms that produce a variety of organic acids, including humic acid 17-36% and fulvic acid 30-31% of the total amount of organic matter, thus enhance the solubility of minerals, particularly potassium and phosphorus (Senesi et al., 1992; Ievinsh et al., 2020; Celikcan et al., 2021). Compared to compost, Vermicompos has a higher quality because, vermicompost's higher organic matter decomposition rate and nutrient content in the final product are. Furthermore, vermicompost generated a greater quantity of hormones and enzymes that could enhance plant growth and discourage plant pathogens (Wu et al., **2014**). The microorganisms in vermicomposts and those in traditional composts were compared conventional compost had a lower population of bacteria  $(5.7 \times 107)$ , fungi  $(22.7 \times 104)$  and actinomycetes (17.7) $\times$  106) than vermicomposts (Nair et al., 1997).

This study aims to assess the influence of sowing dates and organic fertilizer treatments on the vegetative growth characteristics and mucilage percentage of the *Althea rosea* plant.

### **MATERIALS AND METHODS**

### **Experimental Site**

This experiment was conducted at Experimental Farm of Ornamental and Medicinal Plants, Faculty of Environmental Agriculture Science, Arish University, North Sinai (13°07'33.3"N, 13°49'22.3"E, 226 M above mean sea-level) during two consecutive winter seasons in 2021-2022 and 2022-2023 and the laboratories of the Faculty of Environmental Agriculture Science, Arish University. The chemical and physical analyses of the soil used. As well as organic fertilizer were tabulated in Tables 1 and 2 and irrigation water used in Table 3 which was analyzed at the National Research Centre. Seeds of Althea rosea were procured from the Agricultural Research Center. Chicken manure (ChM) was sourced from a poultry farm Faculty of Environmental Agriculture Science, Arish University, and vermicompost (VC) and compost (C) were procured from а nearby company. Meteorological data, temperature degrees (C°) and total rain (mm./month) for El-Arish region during (2021/2022 and 2022/ 2023) seasons are shown in Table 4 by the Center Laboratory for Agriculture Climate.

### **Experiment Design**

The experiment was planned in a split plot design with three replicates. The main plot consisted of three different sowing dates as  $15^{\text{th}}$  October,  $1^{\text{st}}$  November, and  $15^{\text{th}}$  November for both seasons. The subplots included applying three types of organic fertilizers: control = recommendation for using chicken manure at 10 tons per feddan, chicken manure at (5 t/fad.) mixture with vermicompost at (3 t/fed) and chicken manure at (5 t/fad.) mixture with compost at (3 t/fad.). Organic fertilizers were added during the soil preparation for cultivation.

A drip irrigation system was used, and *Althea rosea* seeds were directly planted in hill with a spacing of 1.0 m between rows and 50 cm between plants on the same row,

where each plant contained an area of 50  $\text{cm}^2$  to again plot area of 12  $\text{m}^2$ , containing 24 plants (8400 plants/fed). After one month, the plants were thinned to one plant/ hill and all natural agriculture practices were carried out as needed by the plants.

# Vegetative growth parameters of *Althea rosea*

After planting, the plant's vegetative growth was recorded after 5 months for two seasons. And were taken the height of a plant from the soil surface to its highest point was taken, as well as the number of leaves/plants, the fresh and dry weights of leaves/plants (g), the length of the root (cm), the fresh and dry weights of roots/ plant (g). Furthermore, the plant's leaves and roots were dried at 70°C until they reached a constant weight, and their dry weights were measured in (g).

### Yield parameters of Althea rosea

The yield of seeds and roots were recorded on the *Althea rosea* plant at the end of both seasons (after 7 months of cultivation) in two seasons. Measurements were taken for the following parameters:

- 1- Seeds yield (kg/fed): = the weight of seeds per plant × number of plants per feddan/1000
- 2- Root yield (kg/fed<sup>-</sup>): = the weight of roots per plant × number of plants per feddan/1000.
- 3- Mucilage in root and seed (kg/ fed): = percentage of mucilage in root or seed × yield of root or seed/ 100

The process of extracting and estimating the percentage of mucilage:

After the end of both seasons (7 months after planting), Extraction of mucilage from *Althea rosea* roots and seeds according to **Le** *et al.* (2020). The roots and seeds were washed to remove dirt and the roots were chopped to 5 mm, taken to dry at a temperature of  $50^{\circ}$ c until constant weight. The extraction process was executed in two steps.

	Particles size distribution (%)								
	First season 2021-2022	Second season 2022-2023							
Coarse sand%	58.3	58.4							
Fine sand%	27.4	27.5							
Silt%	12.3	12.0							
Clay%	2.0	2.1							
Soil texture	Sandy loam	Sandy loam							
Bulk density (Mgm <sup>-3</sup> )	1662	1661							
	<b>Chemical properties</b>								
Ca+ (meq. <sup>L-1</sup> )	10.37	10.37							
Mg+ (meq. <sup>L-1</sup> )	5.19	5.25							
$Na+(meq.^{L-1})$	15.06	15.09							
K+ (meq. <sup>L-1</sup> )	0.07	0.07							
CO3 <sup>-</sup> (meq. <sup>L-1</sup> )	-	-							
HCO3 <sup>-</sup> (meq. <sup>L-1</sup> )	4.52	4.70							
Cl <sup>-</sup> (meq. <sup>L-1</sup> )	20.63	20.57							
<b>SO4</b> (meq. <sup>L-1</sup> )	5.54	5.51							
EC (dSm-1)	3.11	3.15							
рН	7.52	7.50							
Organic matter %	0.153	0.160							
<u>CaCo3 %</u>	22.43	22.48							

Table 1. Some initial chemical and physical characteristics of soil	

Table 2. Organic fertilizer content of nitrogen, phosphorus and potassium

Organic fertilizers	N%	P%	K%
Chicken manure	1.20	0.45	0.80
Vermicompost	2.35	1.60	1.50
Compost	0.71	0.39	0.42

Table 3. Analyzing of the chemical composition of irrigation water

рН	FC	Soluble ions (meq. <sup>L-1</sup> )									
	$(dSm^{-1})$		Cati	ons		Anions					
	(usin )	Ca <sup>++</sup>	$Mg^{++}$	Na <sup>+</sup>	$\mathbf{K}^{+}$	Cl.	HCO3 <sup>-</sup>	CO3	SO4 <sup></sup>		
	First season (2021/2022)										
7.55	5.56	19.50	17.36	18.50	0.24	45.92	2.90	-	6.78		
Second season (2022/2023)											
7.60	5.71	21.00	17.05	18.80	0.25	46.77	2.99	-	7.34		

918

	2021-2	022	2022-2023			
Months	Average Temperature(°C)	Precipitation (mm)	Average Temperature (°C)	Precipitation (mm)		
October	23.76	5.80	24.06	16.30		
November	21.25	5.40	20.03	10.10		
December	15.64	25.10	17.10	13.90		
January	11.86	107.50	14.50	38.40		
February	13.06	21.90	12.93	33.30		
March	13.06	30.80	17.21	23.90		
April	19.57	2.20	19.06	29.20		
May	22.05	2.40	22.33	2.70		
June	25.86	1.70	25.21	5.60		
Average and Sum	18.46	202.80	19.16	173.40		

Table 3. Meteorological data: temperature degrees (C°) and total rain (mm. /month) forEl-Arish region during (2021/2022 and 2022/2023) seasons

Table 4. Effect of chicken manure, vermicompost, compost and sowing date as well as interaction treatments on plant height, number of leaves, fresh and dry weights of leaves of *Althea rosa* plant

Treatments		Plant heig	ht (cm)	Number of leaves/plant		Fresh we leaves/p	eight of lant (g)	Dry weight of leaves/plant(g)	
		First season	Second season	First season	Second season	First season	Second season	First season	Second season
th	ChM	160.73 cd	169.50 de	43.33 c	53.33 d	98.08 d	176.24 b	40.07 bc	50.37 ab
15 <sup>th</sup> October	ChM +VC	213.30 a	223.40 a	64.33 a	77.33 a	149.56 a	228.54 a	45.24 a	55.41 a
occoser	ChM+C	208.80 a	216.50 b	62.33 a	69.66 b	148.98 a	226.50 a	44.57 ab	54.86 a
. et	СМ	143.20 e	156.53 f	27.66 e	34.66 fg	74.70 e	125.09 e	26.13 f	35.26 d
1 <sup>ន</sup> November	ChM+VC	169.30 b	176.23 c	52.33 b	61.00 c	124.55 b	161.86 c	42.61 ab	51.58 a
	ChM+C	165.06 bc	167.36 e	43.00 c	58.66 cd	124.29 b	160.12 c	33.49 de	42.67 c
th	ChM	124.16 f	145.40 g	27.00 e	31.66 g	67.14 f	114.61 f	23.76 f	28.82 e
15 <sup>th</sup> November	ChM+VC	156.53 d	173.20 cd	33.33 d	41.33 e	106.30 c	131.80 d	30.78 e	37.52 d
	ChM+C	139.06 e	154.90 f	28.66 de	38.00 ef	110.07 c	135.42 d	37.44 cd	46.27 bc

ChM= chicken manure, VC= vermicompost, C= compost

Step 1: mucilage separation, Dried roots or dried seeds were mixed with distilled water (ration 1:8 W/V) and heated at 60°C in water bath for 1.0 hour (the mixture was stirred regularly). After that, the mixture was taken out and cooled at room temperature within 1 hour to increase the amount of mucilage dissolved in distilled water. Thereafter, the mixture wase filtered through 8 layers of muslin cloth to collect the filtrate containing the mucilage.

Step 2: mucilage precipitation and collection. The mucilage was precipitated from the filtration by acetone with the volume ration of 3:1 and it was dried at a temperature of 40°C until a constant weight according to Lia and Liang (2011) and **Prabakran** *et al.* (2011). The mucilage yield was calculated as:

 $Mucilage (\%) = \frac{weight of dried mucilage (g)}{weight of dry root or seeds (g)} \times 100$ 

# Experiment Design and Statistical Analysis

The complete randomized block design (RCBD) in a split plot design system was used experiment with three replicates. All collected data were analyzed with analysis (ANOVA) procedure variance using MSTAT.C statistical software package (Michigan State University, 1983). Differences between means were compared by using Duncan multiple range test at 0.05 (Duncan, 1955)

### **RESULTS AND DISCUTION**

## Effect of Chicken Manure, Vermicompost, Compost and Sowing Date As Well as Their Interaction Treatments on Plant Height, Number of Leaves, Fresh and Dry Weights of Leaves of *Althea rosa* Plant

Data presented in Table 5 show that chicken manure at (5 t/fad.) mixture with vermicompost at (3 t/fad.) treatments generally increased plant height (213.30 cm and 223.40 cm), number of leaves (64.33 and 77.33), fresh and dry weights of leaves (149.56 g, 228.54 g, 45.24g and 55.41 g), respectively during the first sowing date at 15<sup>th</sup> October compared with those of chicken manure ones alone or chicken manure mixed with compost during the second and third sowing date.

Using previous results, the highest values for plant height, number of leaves, and both fresh and dry weights of leaves were attained in early sowing dates (mid-October). In Table 2, the average air temperature for the mid-October planting season under El-Arish conditions was 23.76 and 24.06°C in the first and second seasons, respectively. For seed germination and vegetative growth of marshmallow plants,

this temperature is more favorable. Abuo El-Kasem et al. (2017); Wafaa et al. (2017) and Hasanvand et al. (2018) found that the most significant growth parameters were recorded on the early sowing date of October 15<sup>th</sup> in Pisum sativum, Foeniculum vulgare and Borago officinalis, respectively. The early sowing date in October was reported by El-Mekawy (2012) in Nigella sativa; Megawer et al. (2017) in Faba bean and Islam et al. (2010) in Capsicum annum) have caused an increase in plant height, number of leaves, and fresh and dry weight of plants compared to the later sowing date. The plant's increased growth may have been influenced by the appropriate weather conditions during developmental stages, which may have led to an increase in cell division and elongation. On the other hand, plants that are sowing late may be subjected to unfavorable environmental conditions, which could result in a short duration of vegetable growth, as high temperatures accelerate growth stages.

The increase in plant height, number of leaves, fresh weight of leaves and dry weight of leaves resulting from using vermicompost and chicken manure together. These results agree with those obtained by Abha et al. (2019) reported that okra vegetable growth significantly increased after using vermicompost 2.5 t/ha combination with 2.5 t/ha of poultry manure. Also, Jamkatel et al. (2020) found that vermicompost and chicken manure were most effective in increasing the height of okra plants. Furthermore, organic manure, particularly vermicompost, had a positive impact on plant height compared of the other organic fertilizer (Gutiérrez et al., 2007; Adhikari and Piya, 2012; Muqtadir et al., 2019; Masud et al., 2022), whilst Naseer et al. (2017) observed an increase in plant biomass and number of Abelmoschus esculentus plant leaves when vermicompost was applied at a rate of 5 tons/acre. Whereas, Bhandari et al. (2019), Qulsum et al. (2020) and Samanhudi et al. (2021) in different plants

Treatments		Root length (cm)		Fresh weight of root/plant (g)		Dry weight of root/plant (g)		Yield of roots/ feddan (kg)		Yield of seeds/feddan (kg)	
		First season	Second season	First season	Second season	First season	Second season	First season	Second season	First season	Second season
	ChM	38.63 d	43.33 d	99.88 b	116.15 c	45.52 b	53.06 b	382.37 b	445.73 b	848.09 c	965.65 c
15 <sup>th</sup> October	ChM+VC	59.53 a	68.13 a	112.69 a	145.89 b	53.09 a	65.81 a	445.98 a	552.83 a	1172.86 a	1350.19 a
	ChM+C	50.50 b	57.63 b	115.42 a	157.36 a	49.57 a	55.04 b	416.41 a	462.33 b	859.12 c	962.40 c
1 St	ChM	33.83 e	35.16 e	55.33 d	65.61 f	23.06 e	30.12 de	193.73 e	253.03 de	676.19 e	737.98 e
I" November	ChM+VC	42.56 c	50.66 c	55.85 d	83.40 e	30.70 c	41.94 c	257.93 с	352.35 c	906.35 b	1137.16 b
November	ChM+C	51.76 b	56.70 b	76.90 c	94.66 d	27.48 cd	33.80 d	230.83 cd	283.92 d	772.91 d	857.95 d
15 <sup>th</sup> November	ChM	25.10 f	31.53 f	39.69 f	53.47 g	17.53 f	23.92 e	147.25 f	200.93 e	342.19 g	459.63 g
	ChM+VC	34.83 e	36.93 e	47.60 e	64.07 f	24.39 de	26.82 de	204.93 de	225.34 de	664.38 e	741.7 e
	ChM+C	39.33 d	42.73 d	54.17 d	64.37 f	21.49 e	25.77 e	180.51 e	216.46 e	546.85 f	642.10 f

Table 5. Effect of chicken manure, vermicompost, compost and sowing date as well as interaction treatments on root length, fresh and dry weights of root, yield of roots and seeds of Althea rosa plant

ChM= chicken manure, VC= vermicompost, C= compost

at Malvaceae family they reported increased vegetable growth by improving plant height, number of leaves, and fresh and dry weight of leaves after applied the poultry fertilizer treatment. Also, **Agustanto** *et al.* (2022) in *Abelmoschus esculentus* found that vegetative growth was enhanced when chicken manure was applied at a rate of 20 t/ha.

This increase of vegetative growth parameters reflects that richness vermicompost and poultry manure with N, P, K, micronutrients, and amino acids which a play many important roles as formation of DNA, RNA, cell building, cell division, cell enlargement, formation the coenzyme, synthesis proteins and bud differentiation as well as branching (El-Agrodi *et al.*, 2011; Mistry *et al.*, 2015; Dume *et al.*, 2023).

Effect of Chicken Manure, Vermicompost, Compost and Sowing Date As Well As Interaction Treatments on Root Length, Fresh And Dry Weights of Root As Well As Yield of Roots and Seeds of *Althea rosa* Plant

Results in Table 6 data show that root length, fresh and dry weight of root and yields of roots as well as seeds per plant gradually increased in mid-October by using the treatments of chicken manure (5 t/fad.) plus vermicompost (3 t/fad.). Moreover, addition these treatments gave the highest values of root length (59.53 cm and 68.13 cm), (540.15 kg and 682.64 kg) and (1172.86 kg and 1350.19 kg) in the first and second seasons, respectively and showed significant increase compared to other treatments under study. The advantage of planting early could be linked to the favorable temperature and day length, which could result in better vegetative growth of Althea rosea plants. These findings agree with those reported by (El-Hosary et al., 2010) that the root length and root yield of sugar beet significantly increased with early sowing on October 15th as compared to other sowing dates. According to Abuo El-Kasem et al. (2017), Wafaa et al. (2017) and Hasanvand et al. (2018), the early sowing date on October 15<sup>th</sup> resulted in the greatest vields. In Pisum sativum, Foeniculum vulgare and Borago officinalis respectively. The early sowing date in October was reported by (Megawer et al., 2017 and Islam et al., 2010) have caused an increase yield in Faba bean and Capsicum annum respectively.

Table 6. Effect of chicken manure, vermicompost, compost and sowing date as well as interaction treatments on mucilage % and yield in seeds and roots of *Althea rosa* plant

Treatments		Mucilag (%	e in seeds %)	Mucilag (	ge in roots %)	Yield of m seeds/fed	ucilage in dan (kg)	Yield of mucilage in roots/feddan (kg)	
		First season	Second season	First season	Second season	First season	Second season	First season	Second season
	ChM	3.60 cde	4.38 bcd	11.83 ab	16.53 b	30.51 cd	42.31 cd	45.36 b	73.24 c
15 <sup>th</sup> October	ChM+VC	5.80 a	6.93 a	13.99 a	20.93 a	68.14 a	93.66 a	62.33 a	115.66 a
	ChM+C	4.32 bc	5.24 abc	13.72 ab	19.86 a	37.30 c	50.21 c	57.18 a	92.01 b
1 <sup>st</sup> November	СМ	3.01 de	3.53 cd	7.97 de	10.38 d	20.41 e	26.21 de	15.59 ef	26.24 e
	ChM+VC	5.07 ab	6.24 ab	11.82 ab	15.92 b	45.94 b	71.04 b	30.45 c	56.33 d
	ChM+C	3.12 de	4.90 bc	11.28 bc	11.72 cd	24.02 de	42.08 cd	26.03 cd	33.02 e
1 <b>c</b> th	ChM	2.49 e	2.99 d	7.26 e	9.59 d	8.60 f	13.93 e	10.81 f	19.45 e
November	ChM+VC	4.30 bc	5.19 abc	9.96 cd	13.82 bc	28.60 cde	38.46 cd	20.19 de	31.37 e
	ChM+C	3.98 bcd	4.49 bcd	8.46 de	10.50 d	21.72 de	28.81 de	15.22 ef	22.55 e

ChM= chicken manure, VC= vermicompost, C= compost.

922

These findings agree with those reported by Singh et al. (2016) in Raphanus sativus and Abha et al. (2019) in okra, they reported that all aspects of vegetative and root growth and yield parameters were greatly improved by combining 50% vermicompost and 50% poultry manure. Abul-Soud et al. (2014) reported that the utilization of vermicompost at the rate of 15 to 25  $m^3/fad$ . can result in an improvement in the total yield, number of pods/plants, seed yield, weight of 100 dry seeds, and number of dry seeds per plant in peas plant. The highest rates of chicken manure applied (10 and 20 t/ha) resulted in a significantly substantial high yield of pods and a significantly higher vield fresh and dry weight in Corchorus olitorius plant (Naim et al., 2015).

The use of vermicompost in conjunction with chicken manure is responsible for the increase in root length and fresh weight of root, dry weight root, yield of root and seeds, because the mixture enhances the soil's physical structure, these include things like porosity, ventilation, drainage, corrosion resistance, and infiltration (Lim et al., 2014; Sha et al., 2023) Thus providing a better medium for root growth. According to Edwards and Bohlen (1996) vermicompost is rich in polysaccharides. Polysaccharide's role as a cementing substance in the soil resulted in aggregate stability. Humic acid and fulvic acids have been found in vermicompost, as reported by (Arancon et al., 2006; Singh et al., 2008; and Hussain et al., 2017a,b). Promotes plant growth and causes an increase in cell division, which leads to the absorption of nutrients, the retention of water, the formation of microbial aggregations with soil, the potential to elongate side roots and improve root activity. Moreover, poultry fertilizer has a substantial amount of phosphorus. The activity of phosphorus involves growing roots, increasing root mass, and increasing the absorption of nutritional staples. and is also part of many enzymes and co-enzymes and ATP-rich energy, which increases the growth of roots, increase crop productivity (Mangal, 1985).

## Effect of Chicken Manure, Vermicompost, Compost and Sowing Date As Well As Interaction Treatments on Mucilage % and Yield in Seeds and Roots of *Althea rosa* Plant

The data of mucilage percentage and their yields in dried seeds and roots were shown in Table 7. The results clearly demonstrated that chicken manure at (5 t/fad.) plus vermicompost (3 t/fad.) caused a slight increase in the percentage of mucilage constituents in seeds and roots during both seasons. However, the differences between chicken manure plus vermicompost treatments and other treatments were a high significant in the first and second seasons during the first sowing date. Data also reveal that there were significant differences among treatments in the two seasons. Also, the superior treatment which increased yield of mucilage in seeds and roots per plant and per feddan, yield of mucilage in seeds (68.14 kg and 94.80 kg) and yield of mucilage in roots (93.70 kg and 118.34 kg) during the first sowing date and two seasons respectively.

When plants are grown early (in winter), differences in daytime length and relative daytime temperature system provide better conditions for growing and creating more robust plants. As such, their seed productivity is enhanced. Its spring growth also starts early and shortens the period from sowing to flowering, avoiding the synchronization of the seed filling period with higher environmental temperatures, lower plant height and loss of seed crop This is reflected in the productivity of mucilage in seeds and roots As such, it was found that delayed sowing reduced seed yield and mucilage yield since the growth period was shortened and the flowering and seed filling periods coincided with hot summer temperatures (Ghasemi et al., 2020). According to Hekmat et al. (2002) and Ebrahimi et al. (2010), the earlier sowing date caused a rise in mucilage percentage and yield in

*Plantago ovata* and *Borago officinalis* respectively.

The use of 20 ton/ha of manure had a significant impact on the mucilage content of *Plantago ovata*, according to **Mirshekari** and Forouzandeh (2015). Whereas **Shahbazi** *et al.* (2019a) in *Borago officinalis*, **Shahbazi** *et al.* (2019b) in *Borago officinalis* and **Sadeghi** *et al.* (2014) in *Althea officinalis*) suggest that utilizing 10 t/ha of vermicompost is the best way to obtain the greatest amount of mucilage.

An increase in the proportion of nutrients in the soil can be attributed to both the increase of mucilage percentage on seeds and roots as well as the increase of mucilage yield in seeds and roots. According to Shahbazi et al. (2019a), the plant may have used mucilage as an energy source, which is why mucilage decreased that is comprised of polysaccharides that are specific to plants or long chains of sugar molecules. Based on these findings, Mycorrhiza in has been suggested to increase the availability of water and nutrients for plants and likely plays a significant role in promoting the plant's photosynthesis. According to Pouryousef et al. (2012), organic fertilizers cause an increase in the mucilage percentage of Plantago psyllium. Nitrogen absorption, phytohormone synthesis, and mineral solubility can be organic fertilizer, which it is possible for plant production to experience a significant influence (Goswami et al., 2017; Shakib et al., 2019). By increasing mineral uptake, manure application can improve growth, biomass production, and seed yield, it enhances the efficiency of the leaf water supply, photosynthetic efficiency, transpiration rate, and water uptake of the root tissues of plants (Askary et al., 2018). Thus, it will be reflected in productivity of mucilage yield.

### Conclusion

The result showed that the highest values of all growth parameters: plant height, number of leaves, fresh and dry weight of leaves recorded at the first sowing date October 15<sup>th</sup> with using chicken manure (5 t/fad.) plus vermicompost (3 t/fad.), in addition to the increase in root length, fresh and dry weight of root, root and seeds yield, percentage and yield of mucilage in root and seeds achieved in early sowing date mid-October and mix of vermicompost and chicken manure. Therefore, we recommend planting *Althea rosea* plants on October 15<sup>th</sup> using organic fertilizer consisting of chicken manure (5 t/fad.) and vermicompost (3 t/fad.).

## **REFERENCES**

- Abha, R.; Meena, M.; Singh, R. and Mandal, R.K. (2019). Effect of various organic manures on growth and yield of okra [*Abelmoschus esculentus* (L.) Moench]. J. Pharm. and Phytochem., 8 (6): 1203-1205.
- Abul-Soud, M.; Refaie, K.M. and Abdelraouf, R.E. (2014). Influence of vermicompost and plant density on sustainable production of peas. J. Agri. Sci., Ain Shams Univ. Cairo, 22 (1): 55-65.
- Abuo El-Kasem, S. and Elkassas, M. (2017). Effect of sowing Dates, irrigation intervals and organic fertilizers on growth and productivity of pea plants (*Pisum sativum* L.) under sandy soil conditions. J. Plant Prod., 8 (11): 1159-1175.
- Adhikari, A. and Piya, A. (2012). Effect of different sources of nutrient on growth and yield of okra (*Abelmoschus esculentus* L. Monech). Int. J. Environ. Agric. Res., 6 (1): 45-50.
- Agustanto, R.; Herawati, N. and Warnita,
  W. (2022). Application effect of chicken manure on growth and yield of okra (*Abelmoschus esculentus* L. Moench). Asian Plant Res. J., 9(3):1-7.
- Al Masum, A.; Pal, K.; Saha, I.; Ghosh, D.; Roy, S.; Chowdhury, S.G.; Islam, M.D. and Karmakar, P. (2020). Facile synthesis of antibiotic encapsulated

biopolymeric okra mucilage nanoparticles: molecular docking, *in vitro* stability and functional evaluation; Vietnam Acad. Sci. and Technol., 11:1-10.

- Amini, M. and Razavi, S. (2012). Dilute solution properties of Balangu (*Lallemantia royleana*) seed gum: effect of temperature, salt, and sugar. Int. J. Biol. Macromol., 51(3): 235-43.
- Arancon, N.Q.; Edwards, C.A. and Bierman, P. (2006). Influences of vermicomposts on field strawberries. Part 2. Effects on soil microbiological and chemical properties. Bioresour Technol., 97:831-840.
- Askary, M.; Behdani, M.A.; Parsa, S.; Mahmoodi, S. and Jamialahmadi, M. (2018). Water stress and manure application affect the quantity and quality of essential oil of *Thymus daenensis* and *Thymus vulgaris*. Indust. Crops and Prod., 111: 336-344.
- Aslam, M.; Sddiqui, A.; Shiffa, M.; Husain, A.; Ahmed, S. and Javed, K. (2012). Pharmacognostical, psychological and phytochemical investigations of *Althaea rosea* linn. Int. J. Pharm. Res. and Dev. (IJRD), 4 (03):129-140.
- Bhandari, S.; Pandey, S.R.; Giri, K.;
  Wagle, P.; Bhattarai, S. and Neupane,
  R.B. (2019). Effects of different fertilizers on the growth and yield of okra (*Abelmoschus esculentus* L.) in summer season in Chitwan, Nepal. Archives of Agric. and Environ. Sci., 4 (4): 396-403.
- Celikcan, F.; Kocak, M.Z. and Kulak, M. (2021). Vermicompost applications on growth, nutrition uptake and secondary metabolites of *Ocimum basilicum* L. under water stress: A comprehensive analysis. Indust. Crops and Prod., 171: 113973.
- Classen, B. and Blasheck, W. (1998). High molecular weight acidic polysaccharides

from *Malva sylvestris* and *Alcea rosea*. Planta Med., 64(7): 640-644

- Detersa, A.; Zippela, J.; Hellenbranda, N.; Pappaib, D.; Possemeyera, C. and Hensela, A. (2010). Aqueous extracts and polysaccharides from marshmallow roots (*Althea officinalis* L.): Cellular internalization and stimulation of cell physiology of human epithelial cells in vitro. J. Ethnopharmacol., 127: 62–69.
- Dume, B.; Hanc, A.; Svehla, P.; Michal, P.; Chane, A.D. and Nigussie, A. (2023). Composting and vermicomposting of sewage sludge at various C/N ration: Technological feasibility and end-product quality. Ecotoxicol. and Environ. Safety, 263 (9): 1-11.
- **Duncan, B.D. (1955).** Multiple ranges and multiple F test. Biometrics, 11:1-24.
- Ebrahimi, A.; Moaveni, P. and Farahani, H.A. (2010). Effect of planting dates and compost on mucilage variations in borage (*Borago officinalis* L.) under different chemical fertilization systems. Int. J. Biotechnol. and Molec. Biol., 1 (5): 58-61.
- **Edwards, C.A. and Bohlen, P.J. (1996).** Biology and Ecology of Earthworms (3<sup>rd</sup> Ed.). Chapman and Hall, London.
- El-Agrodi, M.; El-Zehery, T. and Issa, H. (2011). Effect of chicken manure and gypsum on sugar beet (*Beta vulgaris*, Var. *Saccharifora*, L.) under saline condition. J. Soil Sci. and Agric. Eng., 2(6):701-716. doi: 10.21608/jssae. 2011. 55652.
- El–Hosary, A.A.; Salwau, M.I.; Saad, A.M.M.; El-Geddawy, I.H. and Ibrahim, B.S. (2010). Sugar beet yield and its components as affected by sowing date, mineral nitrogen and bio- fertilizers. Egypt. J. Appl. Sci., 25 (8A): 349 – 366.
- **El-Mekawy, M.A.M. (2012).** Growth and yield of *Nigella sativa* L. plant influenced by sowing date and irrigation treatments.

Ame. Euras. J. Agric. and Environ. Sci., 12: 499 - 505.

- Ghasemi, V.M.; Moghaddam, S.S.; Rahimi, A.; Pourakbar, L. and Popović-Djordjević, J. (2020). Winter cultivation and nano fertilizers improve yield components and antioxidant traits of dragon's head (*Lallemantia iberica* (M.B.) Fischer and Meyer). Plants, 252 (9): 1-15.
- Goswami, L.; Nath, A.; Sutradhar, S.; Bhattacharya, S.S.; Kalamdhad, A.; Vellingiri, K. and Kim, K.H. (2017). Application of drum compost and vermicompost to improve soil health, growth, and yield parameters for tomato and cabbage plants. J. Environ. Manag., 200: 243-252.
- Gutiérrez, M.; Federico, A.; Santiago, B.; Jorge, M.M.; Joaquin, A.; Carlos, N.; Camerino, A.; Miguel, O.L.; Maria, A.; Rincón, R. and Dendooven, L. (2007). Vermicompost as a soil supplement to improve growth, yield and fruit quality of tomato (*Lycopersicum esculentum*). Bioresource Technol., 98 (15): 2781-2786.
- Hasanvand, H.; Siadat, S.A.; Bakhshandeh,
  A.; Moradi Telavat, M.R. and Poshtdar,
  A. (2018). Assessment of required growing degree days phenologic stages and growth of borage in different sowing and densities. Iranian J. Field Crops Res., 16 (2): 511-523.
- Heidarzadeh, A.; Modarres-Sanavy, S.A. M. and Mokhtassi-Bidgoli, A. (2022). Changes in yield and essential oil compositions of *Dracocephalum kotschyi* boiss in response to azocompost, vermicompost, nitroxin, and urea under water deficit stress. J. Soil Sci. and Plant Nutr., 22 (1):896–913.
- Hekmat, M.Y.A.; Ahmed, H.A. and El-Gamal, S.M.A. (2002). Effect of sowing dates and NPK levels on active substances

in seeds of isabgol plants (*Plantago ovata* Forsk, L.). J. Agric. Sci. Mansoura Univ., 27 (4): 2451-2457.

- Husain, M.; Wadud, A.; Sofi, H.G.; Perveen, S. and Hafeez, K.A. (2019). Physicochemical standardization of mucilage obtained from *Althaea officinalis* Linn-root. Phoog Mag., 15 (S1): 55-61.
- Hussain, N.; Abbasi, T. and Abbasi, S.A. (2017a). Toxic and allelopathic ipomoea yields plant-friendly organic fertilizer. J. Clean Prod., 148:826-835.
- Hussain, N.; Abbasi, T. and Abbasi, S.A. (2017b). Detoxification of parthe- nium (*Parthenium hysterophorus*) and its metamorphosis into an organic fertilizer and biopesticide. Bioresour Bioprocess, 26 (4): 1-9.
- Ievinsh, G.; Andersone-Ozola, U. and Zeipiņa, S. (2020). Comparison of the effects of compost and vermicompost soil amendments in organic production of four herb species. Biol. Agric. and Hort., 36 (4): 267–282.
- Islam, M.; Saha, S.; Akand, H. and Rahim, A. (2010). Effect of sowing date on the growth and yield of sweet pepper (*Capsicum annum* L.). Agronomski Glasnik., 72:3-14.
- Jamkatel, D.P.; Khatri, S.; Bista, A. and Ghimire, A. (2020). Responses of organic and inorganic fertilizers on growth and yield of okra (*Abelmoschus Esculentus* L. Moench Cv. Arka Anamika). Tropical. Agroecosystems, 1(2): 64-66.
- Jarecki, W. and Bobrecka-Jamro, D. (2021). Effect of sowing date on the yield and seed quality of soybean *Glycine* max L. Merr. J. Elementol., 26 (1): 7-18.
- Khosravi Shakib, A.; Rezaei Nejad, A.;
  Khandan Mirkohi, A. and Kalate Jari,
  S. (2019). Vermicompost and manure compost reduce water-deficit stress in pot marigold (*Calendula officinalis* L. cv. *Candyman orange*). Compost Scie. and Utilization, 27 (1): 61–68.

- Lai, L.S. and Liang, H.Y. (2011). Chemical compositions and some physical properties of the water and alkaliextracted mucilage from the young fronds of *Asplenium ustralasicum* (J. Sm.) Hook. Food Hydrocoll, 26: 344-349.
- Le, O.T.H.; Tran, L.T.; Doan, V.T.; Pham, Q.V.; Ngo, A.V. and Nguyen, H.H. (2020). Mucilage extracted from dragon fruit peel (*Hylocereus undatus*) as flocculant for treatment of dye wastewater by coagulation and flocculation Process. Inter. J. Polymer. Sci., 9.
- Lea Master B.; Holly, J.R. and Sullivan, J.L. (1998). Compost animal manures: Precaution and processing. Anim. Waste Manag., 6: 100-105.
- Lim, S.L.; Wu, T.Y.; Lim, P.N. and Shak, K.P. (2014). The use of vermicompost in organic farming: overview, effects on soil and economics. J. Sci. Food Agric., 95 (6) :114356.
- Maddahi, S.; Rahimi, A.; Siavash Moghaddam, S.; Pourakbar, L. and Popović-Djordjević, J. (2021). Effects of sowing time and chemical, organic and biological fertilizer sources on yield components and antioxidant properties of dragon's Head (*Lallemantia iberica* (M. Bieb.) Fisch. and CA Mey). J. Plant Growth Regulation, 40:1–15.
- Maher, N. (2022). Utilization of hollyhock (*Althaea officinalis* L.) roots in manufacture of juices and jam. Food Sci. and Nutr. Health. Int. J. Family Studies, 3 (2): 193-211.
- Mangal, J.L. (1985). Effect of nitrogen and phosphorus application on growth and yield of onion. Indian J. Hort., 42 (2): 152 154.
- Masud, A.A.C.; Akhter, S. and Amin, A.K.M.R. (2022). Comparative performances of organic amendments on germination, growth and yield indices of three selected okra [*Abelmoschus*]

*esculentus* (L.) Moench] cultivars. J. Agric. Food Environ., 3(1): 15-20.

- Megawer, E.A.; El-Sherif, A.M. and Mohamed, M.S. (2017). Performance of five Faba bean varieties under different irrigation intervals and sowing dates in newly reclaimed soil. Int. J. Agron. Agric., 10 (4): 57-66.
- Michigan State University (1983). MSTAT-C Micro Computer Statistical Program, Version2 Michigan State Univ., East Lan sing.
- Mirshekari, S. and Forouzandeh, M. (2015). Analysis of effect of organic fertilizer on yield and active substance of psyllium herbal plant (*Plantago ovata* L.). Inter. J. Biol.Forum., 7(1): 436-440.
- Mistry, J.; Mukhopadhyay, A.P. and Baur, G.N. (2015). Status of NPK in vermicompost prepared from two common weed and two medicinal plants. Int. J. Appl. Sci. and Biotech., 3 (2): 193 - 196.
- Muqtadir, M.A.; Islam, M.A.; Haque, T. and Nahar, A. (2019). Growth and yield of okra influenced by different types of fertilizers and netting. Progress Agric., 30 (1): 1-9.
- Naim, A.H.; Ahmed, K.M. and Ahmed, F.E. (2015). Effects of chicken manure on growth and yield of jute mallow (*Corchorus olitorius* L.) under rain-fed conditions of Sudan. Open Access Library J., 2: e 2042.
- Nair, S.K.; Naseema, A.; Meenakumari, S.K.; Prabhakumari, P. and Peethambaran, C.K. (1997). Microflora associated with earthworms and vermincomposting. J. Trop. Agric., 35:93 - 98.
- Naseer, H.; Tasneem, A. and Abbasi, S.A. (2017). Enhancement in the productivity of ladies finger (*Abelmoschus esculentus*) with concomitant pest control by the vermicompost of the weed salvinia (*Salvinia molesta*, Mitchell). Int. J. Recycl. Org. Wast Agric., 1: 1-9.

- Nawaz, H.; Akram, H.; Ishaq, Q.H.M.;
  Khalid, A.; Zainab, B. and Mazhar, A. (2022). Polarity-dependent response of phytochemical extraction and antioxidant potential of different parts of *Alcea rosea*. Free Radicals and Antioxidants, 12 (2): 49 54.
- Niknam, V. (1999). Identification of secondary metabolits (N- aliphatic composition, mucilage polyssacharides, saponines, sterols and phenolic compositions). Ph.D. Thesis, School Wang, Sci., Tehran Univ., 2(1): 3-9.
- Ofosu-Anim, J.; Blay, E.T. and Frempong, M.E. (2006). Effects of organic manure on okra (*Abelmoschus esculentus* L. Moench) production. J. Appl. Hort., 8 (2): 155-158.
- Oraee, A.; Shoor, M.; Oraee, T.; Tehranifar, A. and Nemati, H. (2022). Organic amendments role in reducing drought stress in *Alcea rosea* L. adv. Hort. Sci., 36(3): 201-214.
- Pouryousef, M.; Mazaheri, D.; Chaiechi, M.R.; Rahimi, A. and Avakoli, A.T. (2012). Effect of different soil fertilizing treatments on some of agro morphological traits and mucilage of Isabgol (*Plantago ovate* Forsk). Electronic J. Crop Prod., 3 (2): 193-213.
- Prabakran, L.; Murthy, V.S.N. and Karpakavali, M. (2011). Extraction and characterization of *Hibiscus rosa-sinensis* leaves mucilage for pharmaceutical applications. J. Pharm Sci., 1: 232-238.
- Prisa, D. and Testa, A. (2022). Use and evaluation of new organic fertilizers in the cultivation of *Abelmoschus esculenthus* L. (okra). World. J. Biol. Pharm. and Health Sci., 11(03): 77-84.
- Qulsum, U.; Meem, F.F.; Promi, R.S.; Zaman, J.R.; Ara, M.F. and Rahman, M.K. (2020). Growth performance of jute (*Corchorus capsularies* L.) as

influenced by different organic manures. J. Biodiversity Conserv. and Biores. Manag., 6(1):17-24.

- Rad, S.K.; Madani, H.; harifabadi, H.H. S.; Mahmoudi, M. and Nourmohamadi, G. (2022). Effects of different irrigation intervals and sowing time on yield attributing traits of okra (*Abelmoschus esculentus* L.). Arabian J. Geosci., 740: 1-9.
- Rahimi, A.; Siavash Moghaddam, S.;
  Ghiyasi, M.; Heydarzadeh, S.;
  Ghazizadeh, K. and PopovićDjordjević,
  J. (2019). The influence of chemical, organic and biological fertilizers on agrobiological and antioxidant properties of Syrian Cephalaria (*Cephalaria syriaca* L.). Agric., 9(6):1-13.
- Sadeghi, A.A.; Kelarestaghi, K.B. and Ghalibaf, K.H. (2014). The effects of vermicompost and chemical fertilizers on yield and yield components of marshmallow (*Altheae officinalis* L.). Agroecol., 6(1): 42- 50.
- Saha, B.C.; Iqbal Hossen, A.S.M.; Rahman, T.; Hossain, M.J. and Zakaria, K.M. (2015). Influence of sowing time and variety on growth and yield of jute (*Corchorus capsularis* L.) at Southern Region of Bangladesh. Europ. Acad. Res., 3(4): 4392-4402.
- Samanhudi, P.H.; Handayanta, E.;
  Hartanto, R.; Yunus, A. and Prabawati,
  K. (2021). Effects of manure types on the growth and yield of sweet sorghum (Sorghum bicolor L.) in dryland. IOP Conf. Series: Earth and Environ. Sci., 807: 1-11.
- Senesi, N.; Saiz-Jimenez, C. and Miano, T.M. (1992). Spectroscopic characterization of metal-humic acid-like complexes of earthworm-composted organic wastes. The Sci. Total Environ., 117/118: 111-120.
- Sha, C.; Wu, J.; Wu, J.; Ye, C.; Shen, C.; Su, J. and Wang, M. (2023). Effects of different fertilizers on soil microbial

diversity during long-term fertilization of a corn field in Shanghai, China. Diversity, 15 (1): 1-16.

- Shahbazi, Z.; Salehi, A.; Dehnav, M.M. and Farajee. H. (2019a). The effect of organic fertilizer and mycorrhizal fungus on morphological characteristics, shoot biomass and mucilage of borage (*Borago officinalis*). Iranian. J. Hort. Sci., 50 (3): 561-570.
- Shahbazi, Z.; Salehi, A.; Hazrati, S. and Dehnavi, M.M. (2019b). Enhancing the quality and yield of European Borage (*Borago officinalis*) by simultaneously Application of Granulated compost, vermicompost and Mycorrhiza. Int. J. Hort. Sci., 6(2):283-298.
- Shakib, K.A.; Nejad, A.R.; Mirkohi, A.K. and Jari, S.K. (2019). Vermicompost and manure compost reduce waterdeficit stress in pot marigold (*Calendula* officinalis L. cv. Candyman orange). Compost Sci. and Utilization, 27 (1): 61–68.
- Singh, R.; Sharma, R.; Kumar, S.; Gupta R. and Patil, R. (2008). Vermicompost substitution influences growth, physiological disorders, fruit yield and quality of strawberry (*Fragaria x ananassa* Duch.). Bioresour. Technol., 99: 8507-8511.
- Singh, V.; Naseeruddin, K.H. and Rana, D.K. (2016). Effect of organic manures on growth, yield and quality of radish (*Raphanus sativus* L.) cv. Pusa Desi. Hort. Flora., 5 (2): 129-133.
- Talyan, V.; Dahiya, R.P. and Sreekrishnan, T.R. (2008). State of municipal solid waste management in Delhi, the capital of India. Waste Manag., 28: 1276-1287.
- Tomoda, M.; Gonda, R.; Shimizu, N.; Akiyama, S. and Arai, H. (1985). Plant Mucilages. A Representative Mucilage, Althaea-Mucilage R from the Leaves of *Althaea rosea*. Chm. Pharm. Bull. Tokyo, 33 (10): 4320-4325.

- Wafaa, A.E.A.; Hendawy, S.F.; Hamed, E.S. and Toaima, W.I.M. (2017). Effect of planting dates, organic fertilization and foliar spray of algae extract on productivity of Dutch fennel plants under Sinai conditions. J. Med. Plants Studies, 5(3): 327-334.
- Wu, T.Y.; Lim, S.L.; Lim, P.N. and Shak, K.P.Y. (2014). Biotransformation of biodegradable solid wastes into organic fertilizers using composting or/and vermicomposting. Chem. Eng. Transactions, 39: 1579-1584.

الملخص العربى

تأثير أنواع مختلفة من الأسمدة العضوية ومواعيد الزراعة على إنتاجية البذور والجذور ومحتوياتها من الميوسيلاج لنبات الخطمية

> سماح أبوبكر أحمد، دينا عبد العاطي سليمان، محمد أحمد محمود قسم الانتاج النباتي، كلية العلوم الزراعية البيئية، جامعة العريش، مصر

تم إجراء تجربة حقلية خلال موسم الشتاء لعامي 2021-2022 و 2022-2023 في المزرعة التجريبية، كلية العلوم الزراعة البيئية بالعريش، جامعة العريش، شمال سيناء، لدر اسة تأثير ثلاثة أنواع من الأسمدة العضوية: معاملة الكنترول وتم إستخدام الكمية الموصي بها من سماد الدواجن بمعدل 10 ط/فدان، ومعاملة سماد الدواجن بمعدل 5 ط/فدا +سماد الفير ميكمبوست بمعدل 3 ط/فدان، و معاملة سماد الدواجن بمعدل 5 ط/فدان بسماد الكمبوست بمعدل 5 ط/فدان، بالإضافة إلى ثلاثة مواعيد زراعة (15 أكتوبر، 1 نوفمبر و 15 نوفمبر) على نمو نبات الخطمية ومحصول البذور والجذور ومحتوي الميوسيلاج. وأظهرت النتائج أن أعلى قيم لجميع مقاييس نمو النبات مثل: ارتفاع النبات، وعدد الأوراق، والوزن الطازج والجاف للأوراق والتي تم تسجيلها عند الزراعة في 15 أكتوبر واستخدام السماد العضوي المكون من سبلة الدواجن بمعدل (5 ط/فدان) بالفير ميكومبوست بمعدل (3 ط/فدان)، كما أدى الي واستخدام السماد العضوي الموراق، والوزن ومحتوي الميوسيلاج. وأظهرت النتائج أن أعلى قيم لجميع مقاييس نمو النبات مثل: ارتفاع النبات، وعدد الأوراق، والوزن الطازج والجاف للأوراق والتي تم تسجيلها عند الزراعة في 15 أكتوبر واستخدام السماد العضوي المكون من سبلة الوازج والجاف للأوراق والتي تم تسجيلها عند الزراعة في 15 أكتوبر واستخدام السماد العضوي المكون من سبلة والعازج والجاف للأور ال والتي تم تسجيلها عند الزراعة وي 15 أكتوبر واستخدام السماد العضوي المكون من سبلة والحاز والحاف للأور والي والوزن

الكلمات الإسترشادية: الخطمية، الأسمدة العضوية، زرق الدواجن، الكمبوست، الفرميكمبوست، والميوسيلاج.

REVIEWERS:| dr\_2hms@yahoo.comDr. Hany M.S. Hassan| dr\_2hms@yahoo.comDept. Plant Prod., Fac. Environ. Agric. Sci., Arish Univ., Egypt.| mohammed Ahmed 1980@yahoo.comDr. Mohammed Ahmed Abdelkader| mohammedahmed1980@yahoo.comHort. Dept., Fac. Agric., Zagazig Univ., Zagazig, Egypt.