

CONSTRUCTION AND CHARACTERIZATION OF BACTERIAL ARTIFICIAL CHROMOSOME LIBRARY OF EGYPTIAN COTTON (*GOSSYPIUM BARBADENSE* L.)

OSAMA, A.MOMTAZ¹, ABEER, A.IBRAHIM¹,
ISMAIL.A. KAMEL², AND M. A. MADKOUR

1. Agricultural Genetic Engineering Research Institute, Agricultural Research Center.
2. Department of Botany, Faculty of Science, Cairo University.

(Manuscript received 3 October, 1999)

Abstract

BAC library for the Egyptian cotton *Gossypium barbadense* Giza 70, Giza 86, and Giza 75 varieties have been constructed and characterized. The isolation and purification of high molecular weight DNA from nuclei embedded in agarose microbeads was an essential part of this work. Several experimental parameters were investigated including optimization of megabase-size DNA restriction enzyme digest and CHEF gel conditions to achieve the highest resolution and separation of such DNA. Fragments ranging in size from 200Kb to 500Kb were size selected and recovered from agarose gel to be used in BAC library construction. The BAC vector, pBeloBAC II derived from the endogenous *E.coli* F-factor plasmid was used in library construction. Different insert: vector ligation ratios examined indicated 15:1 insert:vector to be the optimum ligation ratio. The maximum construct-host transformation efficiency calculated at 1.8 KV electroporation voltage. Purification and isolation of BAC clones were followed by restriction enzyme digest using NotI to characterize insert sizes. Characterization of insert sizes and integrity was achieved by using CHEF gel electrophoresis conditions at 6V/cm, 5 and 15 sec initial and final switch times respectively, and 12°C temperature for 12 hrs. BAC libraries for Giza 70, Giza 86, and Giza 75 varieties contained 45,237; 45,742; and 46,531 clones respectively with an average insert size of 100 Kb. Considering the haploid genome size for cotton to be 2,118 Mb, BAC libraries contains 2.12 haploid genome equivalent. Egyptian cotton BAC library constructed for *Gossypium barbadense* Giza 70, Giza 86, and Giza 75 varieties provide 88% probability of isolating a specific genomic region and are optimized for future gene identification, isolation, target gene mapping, and chromosome walking.

INTRODUCTION

Intensive efforts to construct a high resolution physical map of many plant crops are currently underway (Ahn *et al.*, 1993; Coulson *et al.*, 1986; Kurata *et al.*, 1994; Zhang and Wing, 1997; Zwick *et al.*, 1998; Bush and Wise 1998). Eventually, these maps are composed of overlapping fragments of DNA and allow direct acquisition of DNA fragments that correspond to specific genes. Completion of the physical

map requires the availability of comprehensive libraries of DNA clones in appropriate vectors. Furthermore, accuracy and efficiency of physical mapping increase progressively with the size of the cloned fragments in these libraries. While chromosome walking is a straight forward process in organisms with small genomes, it is quite difficult to apply it in most plant species which typically have large complex genomes in the absence of large-insert DNA libraries (Tanksley *et al.*, 1995; Hong 1997; and Zhu *et al.*, 1997).

Recently, some laboratories have explored the use of BAC cloning system for plants and animals (Woo *et al.*, 1994; Kim *et al.*, 1996; Danesh *et al.*, 1997; Yang *et al.*, 1997; Zhang and Wing 1997; Marek and Shoemaker 1997; Bent *et al.*, 1998; and Miller *et al.*, 1998). It is emerging as one of the systems of choice for plant genomic libraries with average insert sizes of 150 Kb. Cloning of exogenous DNA into bacterial artificial chromosomes (BACs) provides a new approach to the analysis of genomes of higher organisms (Shizuya *et al.*, 1992; Zhang and Wing 1997; Marek and Shoemaker 1997; and Bent *et al.*, 1998). BAC libraries containing genomic DNA inserts are important tools for physical mapping, genomic sequencing and positional cloning. Since 1994, Several human and plant BAC libraries have been constructed (e.g., *sorghum*: Woo *et al.*, 1994; *arabidopsis*: Choi S *et al.*, 1995; rice:Zhang HB *et al.*, 1996; and human: Kim *et al.* 1996, Danesh *et al.*, 1998). The average size of the plant libraries was 100 to 160 Kb. This system has advantage over both cosmid and YAC (yeast artificial chromosome) systems (Zhang, *et al.*, 1996; Bent *et al.*, 1998). potential advantage of BAC system over YAC system include low level of chimerism, increased stability over high copy vectors (Kim, *et al.*, 1992), and ease of library generation and increased manipulation (O'Conner, *et al.*, 1989; Woo, *et al.*, 1994; and Miller *et al.*, 1998). It has been demonstrated that BAC system has permitted cloning of DNA fragments up to 7-fold larger than cosmid system but it is much smaller than those of YACs (Shizuya *et al.*, 1992; Woo *et al.*, 1994; Zhang *et al.*, 1996; and Zhu *et al.*, 1997).

The basis of high degree of stability of DNA fragments containing repetitive sequences is the single-copy number of vector (Kim *et al.*, 1992; Shizuya *et al.*, 1992; and laonnou *et al.*, 1994). Results shown by Shizuya *et al.*, (1992) indicated that stable structure for cloned human DNA in the BAC vector represents a stable source of specific fragments. Very little or no rearrangement of the inserts or chimerism have been observed (Shizuya *et al.*, 1992; Woo *et al.*, 1994; Jiang *et al.*, 1995; Cai *et al.*, 1995; Zhu *et al.*, 1997). All current methods for isolating high molecular weight DNA resulted in a mechanical shearing of DNA with the largest fragments being about 50Kb (Couch 1990; Paterson *et al.*, 1993). Recently, techniques for the isolation of megabase-size DNA have been developed for a number of plant

species (Ganal and Tanksley, 1989; Daelen *et al.*, 1989; Cheung and Gale, 1990; Sorbal *et al.*, 1990; Honeycutt *et al.*, 1992; Hatano *et al.*, 1992; Guidet and Langridge, 1992; Wing *et al.*, 1993; Zhang *et al.*, 1995, Frijters *et al.*, 1997). The DNA isolated by these techniques have been successfully applied to large-scale physical mapping for specific chromosome regions (Ganal *et al.*, 1989; Cheung *et al.*, 1991; Wing *et al.*, 1994; Zhang and Wing, 1997; Zwick *et al.*, 1998; and Bush and Wise 1998) and map-based cloning of economical important genes in plant genomes (Arondel *et al.*, 1992; Martin *et al.*, 1992; Leyser *et al.*, 1993; Marek and Shoemaker 1997; and Miller *et al.*, 1998).

In this work, we report the construction and characterization of BAC library for Egyptian cotton *Gossypium barbadense* Giza 70 (extra-long staple), Giza 86, and Giza 75 (long staple) varieties. Isolation of high molecular weight DNA from nuclei embedded in agarose microbeads, followed by optimization of megabase-size DNA restriction enzyme digest using HindIII enzyme are examined. Fragments ranging in size from 200 to 500Kb are targeted and recovered for BAC library construction using pulsed field gel electrophoresis. Different insert: vector ligation ratios are tested for optimization of ligation conditions. The maximum construct-host transformation efficiency is optimized using electroporation technique. Characterization of constructed BAC clones took place using pulsed field gel electrophoresis technique.

MATERIALS AND METHODS

BAC vector, pBeloBAC II, is derived from endogenous *E. coli* F-factor plasmid which contains genes for strict copy number control with unidirectional origin of DNA replication (Figure 1). Additionally, pBeloBAC II has three unique restriction enzyme sites (HindIII, BamHI & SphI) located within the LacZ gene which can be used as cloning sites for megabase-size DNA.

BAC vector Preparation

pBeloBACII (7.4 Kb) in *E. coli* strain DH10B was provided by Dr.M.Simon, Cal Tech university (Figure 1). pBeloBACII vector was isolated as described by Woo *et al.* (1994) and Wang *et al.* (1995). Cells were harvested from an overnight culture in LB medium containing 12.5 µg/ml chloramphenicol (O.D. = 1.4). Plasmid DNA was isolated using the Plasmid Maxi Kit (Qiagen).

pBeloBAC II (20 µg) was digested to completion with 3 units of HindIII (Biolabs) per microgram DNA at 37°C for 5 hrs. One µg aliquots of dephosphorylated pBeloBAC II plasmid were stored at -80°C until needed.




Figure 1. Diagram of pBeloBAC II. Size fractionated fragments of 250 kbp of Hind III digested of HMW DNA of Egyptian Cotton varieties Giza 70, Giza 86 and Giza 75 were introduced to the Hind III site of pBeloBAC II vector. Insertional inactivation of the lacZ gene by the cloned DNA fragment results in white colonies after transformation.

Plant material and megabase DNA isolation

Cotton seeds of *Gossypium barbadense* Giza 70, Giza 86, and Giza 75 varieties were obtained from Cotton Research Institute, Agricultural Research Center (ARC), Giza, Egypt. Seeds were grown in the greenhouse of the Agricultural Genetic Engineering Research Institute. Young leaves were obtained from four to six weeks old seedlings, and placed immediately in liquid nitrogen. Cotton nuclei were isolated as described by Zhang *et al.* (1995). DNA homogenization buffer (1X) as described by Zhang *et al.* (1995) was substituted by cotton DNA extraction buffer as described by Paterson *et al.*, (1993). 0.5% Triton X-100 (w/w) was added immediately before use. Nuclei were pelleted at relatively low speed centrifugation (200 g instead of 1800g). The nuclei were then embedded in agarose plugs (75 μ l of 1% agarose). Plugs were incubated in nuclei lysis buffer which consists of 0.5 M EDTA (pH 9.0), 1% (w/v) sodium lauryl sarcosine and 0.2 mg/ml proteinase K (Boehringer Mannheim) at 50°C for two days. Nuclei lysis buffer was renewed after 24 hrs. Nuclei suspended in TE buffer were stored at 4°C. Prior to enzyme digestion, plugs were dialysed overnight against TE buffer plus 40 μ g/ml PMSF (phenyl methyl sulphonyl fluoride) at 4°C. The concentration of PMSF was increased to 80 μ g/ml followed by an additional incubation for 1h at room temperature to insure complete inactivation of proteinase K. Plugs were washed four times in TE for 30 min at room temperature. For partial digestions using Hind III, the plugs were equilibrated twice with Hind III restriction enzyme buffer for 1hr at room temperature. Plugs were melted at 65°C for 15 min. and held at 37°C for 5 min. For each plug (75 μ l), containing approximately 15 μ g DNA, 7 U HindIII was added and incubated for 45 minutes at 37°C. The reaction was stopped by the addition of 1/10 volume of 0.5 EDTA, pH 8.0. Partially digested cotton DNA was size fractionated on a low melting point agarose CHEF gel at 6 V/cm, with a 90s pulse, for 20h, at 11°C, in TAE buffer (50 mM Tris-acetate, pH 7.7; 0.5 mM EDTA pH 8.0) (Chu, *et al.*, 1986; Orbach, *et al.*, 1988). DNA ranging from 300 to 500 Kb was cut from the gel, and used for ligation or subjected to a second size fractionation. A second size selection was conducted for 16 h at 11°C running in TAE buffer, to generate a compression zone in excess of 1 Mb using a contour clamped homogeneous, electric field (CHEF DR III apparatus, Biorad). The second size selection was carried out at 4.0 V/cm, with a 5s pulse in TAE buffer. The compressed DNA band was excised from the CHEF gel and stored at 4°C. A gel piece containing size selected DNA was melted at 67°C for 5 min. and digested with β -agarase (New England Biolabs). One unit of β -agarase per 100 μ l of melted agarose gel was incubated at 40°C for 1hr.

Construction of BAC library

After digestion, about 50 ng of size selected cotton DNA was ligated to 750 ng dephosphorylated pBeloBAC II vector in a molar ratio of 1 (Cotton DNA) to 15 (vector DNA) in a total volume of 100 μ l. 400 unit T4 DNA ligase was added to the ligation mixture and incubated overnight at 16°C. Reaction was stopped by incubation for 10 min at 65°C, then stored at 4°C until used. 15 μ l of ligation product was transformed into 100 μ l of electrocompetent ElectroMAX DH10B cells (Gibco BRL) using a cell porator system (Biorad) at the following settings: voltage : 1.8, capacitance: 25 μ F; impedance Low ohms; charge rate: fast; Voltage booster resistance: 200 ohms. Transformed cells were resuspended immediately in 1 ml of 30°C medium (2% Bactotryptone, 0.5% Bactoyeast extract, 10mM NaCl, 2.5 mM KCl, 10 mM MgCl₂, 10 mM Mg SO₄, 20 mM glucose, pH 7.0), incubated at 37°C with shaking speed 225 rpm for 1h and then plated on LB plates containing chloramphenicol (12.5 μ g/ml), X-GAL, and IPTG for 36-48 hrs. White recombinant BAC clones were picked with a toothpick to a second LB plus CM (12.5 μ g/ml), X-GAL, and IPTG plate to verify the color selection. White colonies from the second plate were then transferred to 96 well microtiter dish containing 200 μ l of LB Freezing buffer (36 mM K₂HPO₄, 13.2mM KH₂ PO₄, 1.7mM sodium citrate, 0.4 mM Mg SO₄, 6.8 mM (NH₄)₂SO₄, 4.4% (v/v) glycerol, LB). Microtiter dishes were stored at -80°C for long term storage. The library was replicated in triplicate and stored in - 80°C.

Analysis of BAC clones

Individual BAC clones were inoculated in 5 ml of LB broth containing 12.5 μ g/ml chloramphenicol and grown at 37°C with shaking speed 250 rpm overnight. The circular BAC DNA was isolated by the alkaline lysis method used for the preparation of plasmid DNA. One quarter of the BAC DNA from the overnight culture was digested with NotI to release cotton DNA genomic insert from the cloning vector for at least 3 hrs. The digested BAC DNA was analyzed by PFGE with the CHEF III (BioRad, USA) using Lambda concatemer DNA ladder as the molecular weight mark. Conditions of the PFGE for the CHEF III were: 1% agarose gel, 0.5 x TBE at 11°C, and 6.0V/cm for 12 h with linear ramping time from 5 to 15 s. The gel was stained with ethidium bromide and photographed. The insert size for each isolated BAC clone was determined.

RESULTS AND DISCUSSION

Recently, techniques for megabase-size DNA isolation have been developed for number of plant species (Zhao *et al.*, 1994; Zhang *et al.*, 1995; Frijters *et al.*, 1997). These techniques involve protoplasts or nuclei embedded in agarose plugs or microbeads. The agarose acts as a solid yet porous matrix which allows for the diffusion reagents for DNA purification and subsequent manipulations while preventing the DNA from being sheared. Most protocols for the isolation of megabase-size DNA from plants utilize the protoplast method (Raymond *et al.*, 1989; Wing and Gale, 1990; and Woo *et al.*, 1995). Although the protoplast method yields megabase-size DNA of high quality, yet the process is costly and labor intensive. Zhang *et al.* (1995) have recently developed a nuclei method that works well for several divergent plant taxa. Nuclei are isolated and embedded in agarose plugs or microbeads. The quality of migrated DNA from agarose plugs is as good as DNA prepared from protoplasts, is often more concentrated, and has been shown to contain lower amounts of chloroplast DNA. The primary advantage of this method is that it is economical and not as labor intensive as protoplast method. High molecular weight (HMW) DNA was prepared from Giza 70, Giza 75, and Giza 86 using the previous mentioned method with some modifications (Zhao *et al.*, 1994; Zhang *et al.*, 1995; and Frijters *et al.*, 1997).

To test the quantity and quality of HMW DNA, megabase-sized DNA in agarose plugs was analyzed against DNA prepared by the conventional method (Figure 2A) by PFGE using *Saccharomyces cerevisiae* as a HMW molecular marker ranging from 1.9 Mb to 245 Kb in size. It was then evaluated for size, restriction enzyme digestibility, and yield. Genomic DNA was analyzed in a 0.6% chromosomal grade agarose gel (GibcoBRL) in 0.5 x TBE for 72 hrs at 14°C with 120°C angle, 30 min pulse time and 1.5V/cm voltage gradient using a contour clamped homogenous electric field (CHEF) DRIII apparatus (Biorad) as described by Zhao *et al.*, (1994). The results indicated that the majority of cotton HMW DNA prepared from agarose embedded nuclei is retained in the well of the gel and in the compression zone which is located in a size range greater than 1Mb (Figure 2A). For comparison, the size of cotton DNA prepared by a conventional method shows the majority of DNA between 50-200 Kb in size. There was no significant difference between megabase-size DNA embedded in plugs or in microbeads in yield or quality (Figure 2A). Microbeads were preferred over plugs in restriction enzyme digest because the use of microbeads increases the surface surrounding the tissue sample by approximately 1000 fold thereby allowing for more efficient and rapid diffusion of chemicals and enzyme into and out of the

agarose beads (Zhang and Wing 1997).

Digestion of HMW for genome analysis by PFGE

To determine the quality of HMW DNA in terms of its digestability, cotton DNA embedded in microbeads was partially digested using EcoRI and HindIII enzymes, and was subjected to PFGE. CHEF gel showed several restricted bands on a smeared background for each restriction pattern (Figure 2B). It was noted that all restriction fragments of HMW DNA generated with either enzymes were larger than 50 Kb. EcoRI generated fragments were in the size range between 200-450 Kb, while the size range of fragments resulting from partial digest using HindIII enzyme were in the size range between 200-600 Kb.

G. barbadense BAC library

pBeloBACII has three unique cloning sites within the β -galactosidase gene-BamHI, SphI, and Hind III. Previous experiments showed that HindIII restricts *Gossypium barbadense* DNA frequently with complete digest giving a size range below 50 Kb. To test the BAC system for library construction, *G. barbadense* DNA was partially digested with HindIII and the DNA fragments were size selected by pulsed field gel electrophoresis (Wing *et al.*, 1994; Zhao *et al.*, 1994; Zhang *et al.*, 1995). Figure 2B shows the optimal partial digestion conditions giving DNA in the size range of 200-500 Kb using HindIII. DNA between 300-500 Kb was isolated from this gel. In an attempt to increase the average insert size, a second size selection was made using a low melting point agarose gel, size selected at 4.0 V/cm, with a 5s pulse for 10h, at 11°C. After electrophoresis, the compressed DNA band was excised from the CHEF gel, ligated to pBeloBACII vector and transformed into *E. coli*. Approximately 450 transformants per μ l of ligation reaction was obtained which yielded 45, 742 BAC clones per library.

Figure 4 shows the analysis of 12 BAC clones derived from insert DNA from two size selections. The DNA was digested with NotI to release the insert from the vector and resolved on a CHEF gel. The common band in all the lanes (7.4 Kb) is the BAC vector. The remaining bands are *G. barbadense* DNA fragments. Six of 12 inserts have a size in the range of 97 to 145 Kb. One of the 12 inserts has an internal NotI site while 11/12 do not have an internal NotI site. A hundred and eight BAC clones were similarly analyzed. The average insert size per library ranged between 50-250 Kb.

The clones were grouped by insert size and the insert size of each clone was




Figure 2. **Panel A.** Size fractionation of Megabase DNA isolated from *Gossypium barbadense* L. using CHEF gel. High molecular weight (HMW) DNA was fractionated by PFGE using CHEF III (Biorad). A 0.6% chromosomal grade agarose gel was run in 0.5 x TBE buffer for 72 hrs at 14°C with a 120°C angle, 30 min pulse time and 1.5V/cm voltage gradient. (M) lambda ladder PFG marker (Biolabs), lane (1) Yeast chromosome marker (Biolabs), lanes (2 and 3) HMW DNA of Giza 86 in plugs and microbeads respectively, and lane (4) DNA isolated from Giza 86 using conventional method (as illustrated in materials and methods).

Panel B. Restriction endonuclease enzymes digestion and size fractionation of megabase DNA from Egyptian cotton. Cotton DNA from three varieties were embedded in 50 ul of agarose microbeads, and digested with EcoRI and HindIII enzymes. The DNA was subjected to CHEF on 1% agarose gel in 0.5 TBE buffer using a switch time of 90s at 6V/cm and 12°C with a 120°C angle, for 16 hrs. (M) Yeast chromosome marker (Biolabs), lanes (1,2 and 3) Giza 70, Giza 86, and Giza 75 partially digested DNA using HindIII respectively., lanes (4,5 and 6) Giza 70, Giza 86, and Giza 75 partially digested DNA using EcoRI respectively.

plotted versus the frequency of each group of clones presented in the library (Figure 3B). BAC libraries for Giza 70, Giza 86, and Giza 75 varieties contain 45, 237; 45,742; and 46,742; and 46,531 clones respectively. The average insert size of the library is 100 Kb, which is equivalent to 2.12 haploid genome equivalent. The insert sizes of clones range from 50 to 250 Kb, and approximately 65% of inserts were in the size range between 81 to 145 Kb. Nearly 1.5% of the clones were in the size range between 146 to 250 Kb. Theoretically, the probability of obtaining a particular clone from this cotton BAC library is approximately 88%. We have generated a 2.12 haploid genome equivalent BAC libraries for three Egyptian cotton varieties Giza 70, Giza 86, and Giza 75 using HindIII digest. There is a strong negative correlation between the numbers of colonies obtained and the average insert size. The cotton BAC libraries Giza 70, Giza 86, and Giza 75 contain 45,237; 45,742; and 46,531 clones with an average insert size 100 Kb comprising 4, 523 Mb, 4, 574 Mb and 4,653 Mb DNA respectively from total cotton genome size (2118 Mb DNA).

Figure 3A shows a range of field strengths that was tested to determine the optimum electroporation conditions. Electroporation of *E.coli* strain DH10B electrocompetent cells (Gibco BRL) at 10 KV/cm resulted in a higher number of colonies per 15 μ l ligation mixture which is less than the recommended settings of the Cellporator system (12.5 KV/cm.).

Critical factors to be considered in the construction of BAC libraries

The quantity and quality of genomic DNA targeted for cloning are two of the most critical factors. While preparation of the genomic HMW DNA using the conventional method, it resulted in significant shearing of DNA. Optimization protocol isolation of the mega-base sized DNA embedded in agarose microbeads gave a satisfactory intact DNA profile. Sufficient DNA is required to compensate for the lower transformation efficiencies of large plasmids (Leonardo and Sedivy 1990; Sheng *et al.*, 1995; Wang *et al.*, 1995); nevertheless, if the pulsed field gels are overloaded, there will be excessive trapping of smaller DNA fragments on the PFG. These smaller molecules transform with much higher efficiency, and the average insert size drops dramatically. A significant amount of these smaller fragments can be excluded by a second size fractionation; this results in an increase in the average insert size of BACs (Frijters *et al.*, 1997).

Large genomic fragments have to be circularized with much smaller size plas-

mid. The importance of diluting the large fragments sufficiently to reduce the likelihood of intermolecular ligation has been demonstrated for jumping libraries. The larger the fragments to be cloned, the more they should be diluted. In addition, for the ligation of a small vector molecule to one end of the long DNA fragment followed by ring closure, the concentration of the vector should be roughly equal to the local concentration of the other end of the same molecule. However, diluting the large fragments results in a low concentration of circularized DNA fragments, and yields of transformants by electroporation are directly proportional to input DNA concentrations (Yuling *et al.*, 1995). Three ratios were tested 15:1, 1:10 and 1:15 (insert:vector). The greatest number of BACs per microliter ligation was obtained when approximately 50 ng of large cotton fragments were ligated to 750 ng of dephosphorylated pBelollBAC vector in a total volume of 100 μ l. Based on considerations of stability, ease of isolation, yield, and size, the Egyptian cotton BAC library constructed in this study is suited for use in cotton genome sequencing, gene identification and isolation by chromosome walking. In addition, the library is appropriate for genome mapping and subsequent map-based cloning.




Figure 3. Curve showing the relationship between field strength during electroporation and the number of transformants obtained. 15µl aliquot of ligation product was added to 100 µl of electro MAX DH10B cells (Gibco BRL) and placed between 0.2-cm gap electrodes. Different field strengths were tested with a constant pulse time of 6 mins. The settings for the Cell-porator system (Biorad) were 1-2.5 KV; 25µF capacitor; and 200 ohms resistor.

Panel B. A histogram showing the insert size distribution of BAC clone BAC library. DNA was isolated from 120 random BAC clones and were size fractionated on CHEF gel. The clones were grouped by insert size and were plotted versus its frequency present in the library. Isolated clones have an insert size range from 50 Kb to 250 Kb. The average insert size of the Giza 70 BAC library is 100 Kbp.




Figure 4. Analysis of Giza 70 BAC clone. BAC clones were analyzed by pulsed-field gel electrophoresis (PFGE). The BAC DNA was isolated, digested with NotI to release the insert DNA from the cloning vector, and fractionated on a 1% PFG. The 7.4 kbp band is the cloning vector pBeloBAC II. The insert size of each BAC clone is sum of sizes of all the bands of the lane (M) Lambda concatemer DNA ladder, lanes (1-12) BAC clones. The digested BAC clones was subjected to CHEF on a 1% agarose gel in 0.5 TBE buffer using a switch time of 5 to 15 sec. at 6 V/cm and 11°C with a 120° angle, for 12 hrs.

REFERENCES

- 1 . Ahn, S., J. A. Anderson, M.E. Sorrells, and S.D. Tanksley. 1993. Homologous relationships of rice, wheat and maize chromosomes. *Mol Gen Genet* 241: 483-490.
- 2 . Arondel, V., B. Lemieux, I. Hwang, S.Gibson, H.N. Goodman, and C.R. Somerville. 1992. Map-based cloning of gene controlling omega-3 fatty acid desaturation in *Arabidopsis* *Science*, 258, 1353-1355.
- 3 . Bent, E., S. Johnson, and I. Bancroft. 1998. BAC representation of two low-copy regions of the genome of *Arabidopsis thaliana*. *Plant J.* 13 (6): 849-855.
- 4 . Bush, A.L., and R.P. Wise. 1998. High-Resolution mapping adjacent to the PC71 Crown-Rust Resistance locus in hexaploid Oat. *Mol. Breeding* 4 (1): 13-21.
- 5 . Cai, L, J.F. Taylor, R. A. Wing, D.S. Gallagher, S.S. Woo, and S.K. Davis. 1995. Construction and characterization of a bovine bacterial artificial chromosome library. *Genomics* 29: 413-425.
- 6 . Cheung, W.Y., and M.D. Gale. 1990. The isolation of HMW DNA from wheat, barley and rye for analysis by pulsed-field gel electrophoresis. *Plant Mol.* 14: 881-888.
- 7 . Cheung, W.Y., S. Chao and M.D. Gale. 1991. Long range physical mapping of the α -amylase-1 (α -Amy-1) loci on homoeolodous group 6 chromosomes of wheat. *Mol. Gen. Genet.* 229: 373-379.
8. Choi, S., R.A. Creelman, J. E. Mullet, and R.A. Wing. 1995. Construction and Characterization of a bacterial artificial chromosome library of *Arabidopsis thaliana*. *Plant Mol Biol Rep* 13: 124-129.
9. Chu, G., D. Vollrath and R.W. Davis. 1986. Separation of large DNA molecules by controur clamped homogeneous electric fields. *Science* 234:1582-1585.
10. Couch, J.A., and P.fritz. 1990. Extraction of DNA from plants high in polyphenolics. *Plant Mol. Bio. Repr.* 8 (1): 8-12.
11. Coulson, A., J. Sulston, S.Brenner, and J.Karn. 1986. Toward a physical map of the genome of the nematode *Caenorhabditis elegans*. *Proc Natl Acad Sci USA* 83:7821-7825.

12. Daelen, R.A.J.V., J.J. Jonkers and A.Zabel. 1989. Preparation of megabase-sized tomato DNA and separation of large restriction fragments by field inversion gel electrophoresis (FIGE). *Plant Mol. Biol.* 12:341-342.
13. Danesh, D., S. Penuela, J. Mudge, R. Denny, H. Nordstrm, J.P. Martinez, and N.D. Young. 1997. Construction and characterization of a soybean bacterial artificial chromosome (BAC) library. *Soybean genet. Newsletter.* 24: 196-198.
14. Danesh, D., S.Penuela, J. Mudge, R.Denny, J.P. Martinez, and N.D. Young. 1998. A bacterial artificial chromosome library for Soybean and identification of clones near a major cyst nematode resistance gene. *Theoret. & appl. Genet.* 96 (2): 196-202.
15. Frijers, A.C.J., Z. Zhang, M. van Damme, G.W. Wang, P.C. Ronald, and R. W. Micheltmore. 1997. Construction of a bacterial artificial chromosome library containing large EcoRI and HindIII genomic fragments of lettuce. *Theological Applied Genetics.* 110-115.
16. Ganal, M.W. and S.D. Tanksley. 1989. Analysis of tomato DNA by pulse-field gel electrophoresis. *Plant Mol Biol Rep* 7: 41.
17. Ganal, M.W., N.D. Young and S.D. Tanksley. 1989. Pulsed field gel electrophoresis and physical mapping of large DNA fragments in the Tm-2 α region of chromosome 9 in tomato. *Mol. Gen Genet.* 226:484-490.
18. Guidet, F. and P. Langridge. 1992. Megabase DNA preparation from plant tissue. *Meth. Enzymol.* 216:3-12.
19. Guidet, F. and P.Langridge. 1992. Megabase DNA preparation from plant tissue. *Meth. Enzymol.* 216:3-12.
20. Hatano, S., J. Yamaguchi and A. Hirai. 1992. The preparation of high-molecular weight DNA from rice and its analysis by pulsed-field gel electrophoresis. *Plant Sci.* 83:55-64.
21. Hong, G. 1997. A rapid and accurate strategy for rice counting map construction by combination of fingerprinting and hybridization. *Plant. Mol. Biol.* 35 (1-2): 129-133.
22. Honeycutt, R.J., B. W.S. Sorbal, M. McClelland and A.G. Atherly. 1992. Analysis of large DNA from soybean (*Glycine max* L.Merr.) by pulsed-field gel electrophoresis. *Plant J.* 2: 133-135.

23. Ioannou, P.A., C.T. Amemiya, J. Garnes, P.M. Kroisel, H. Shizuya, C.Chen, M.A. Batzer, and P.J. de Jong. 1994. A new bacteriophage λ -derived vector for the propagation of large human DNA fragments. *Nature Genetics* 6:84-90.
24. Jiang, J., B.S. Gill, G.L. Wang, P.C. Ronald, D.C. Ward. 1995. Metaphase and interphase fluorescence in situ hybridization mapping of the rice genome with bacterial artificial chromosomes. *Proc Natl Acad Sci UA* 92: 4487-4491.
25. Kim, U.J., H. Shizuya, P.J. de Jong, B.W. Birren, and M.I. Simon. 1992. *Nucleic Acids Res.* 20, 1083-1085.
26. Kim, U.J., B.W. Birren, T.Slepek, V. Mancino, C. Boysen, H.L. Kang, M.I. Simon, and H. Shizuya. 1996. Construction and characterization of a human bacterial artificial chromosome library. *Genomics* 34: 213-218.
27. Kurata, N, Y. Nagamura, K. Yamamoto, Y. Harushima, N. Sue, J. Wu, B.A. Antonio, A. Shomura, T. Shimizu, S.Y. Lin, T. Inoue, A. Fukuta, T. Shimano, Y.Kuboki, A. Toyama, Y. Miyamoto, T. Kirihara, K.Hayasaka, A. Miyao, L. Monna, H.S. Zhong, Y. Tamura, Z.X. Wang, T.Momma, Y. Umehara, M. Yano, T. Sasaki, and Y. Minobe. 1994. A 300 Kilobase interval genetic map of rice including 883 expressed sequences. *Nature Genet* 8: 365-372.
28. Leonardo, E.D, and J.M. Sedivy. 1990. A new vector for cloning large eukaryotic DNA segments in *Escherichia coli*. *Biotchnology* 8:841-844.
29. Leyser, H.M.O., C.A. Lincoln, C. Timppte, D.Lammer, J. Turner, and M. Estelle. 1993. *BAC artificial chromosomes*. *Nature*. 364: 161-164.
30. Marek, LF., and R.C. Shoemaker. 1997. BAC contig development by fingerprinting analysis in soybean. *Genome* 40:420-427.
31. Martin, G.B. M.W. Ganal, and S.D. Tanksley. 1992. *Map-based cloning* *Mol. Gen. Genet.* 233:25-32.
32. Miller, J.T., S.A. Jackson, S.Nasuda, B.S. Gill, R.A. Wing, and J. Jiang. 1998. Cloning and characterization of a centromre-specific repetitive DNA element from *Sorghum bicolor*. *Theoret. & appl. Genet.* 96 (6-7) : 832-839.
33. Orbach, M.J., D. Vollrath, R.W. Davis, and C. Yanofsky. 1988. *Construction of BAC library*. *Mol. Cell. Biol.* 8: 1469-1473.

34. O'Connor, M., M. Peifer, and W. Bender. 1989. Construction of large DNA segments in *Escherichia coli*. *Science* 244:1307-1312.
35. Paterson, A. H., L. Brubaker, and J. Wendel. 1993. A rapid method for extraction of Cotton Genomic DNA suitable for RFLP or PCR analysis. *Plant. Mol. Biol. Reprtr* 11 (2): 122-127.
36. Raymond, A.J., J. Jonkers, and P.Zabel. 1989. Preparation of megabase-sized tomato DNA and separation of large restriction fragments by field inversion gel electrophoresis (FIGE). *Plant. Mol. Biol.* 12:341-352.
37. Sheng, Y., V. Mancino, and B. Birren. 1995. Transformation of *Escherichia coli* with large DNA molecules by electroporation. *Nucl Acids Res.* 23: 1990-1996.
38. Shizuya, H., B. Birren, U. J. Kim, V. Mancino, T. Slepak, Y. Tachiri, and M. Simon. 1992. Cloning and stable maintenance of 300-kilobase-pair fragments of human DNA in *Escherichia coli* using F-factor-based vector. *Proc Natl Acad Sci USA* 89:8794-8797.
39. Sorbal, B.W.S., R.J. Honeycutt, A.G. Atherly and M.McCliland. 1990. Analysis of rice genome using pulsed-field gel electrophoresis and rare cutting restriction endonucleases. *Plant. Mol. Biol. Reprtr* 8: 253-275.
40. Tanksley, S.D., M.W. Ganai, and G.B. Martin. 1995. Chromosome landing: a paradigm for map-based gene cloning in plants with large genomes. *Trends Genet* 11:63-68.
41. Wang, G.L., T. E. Holsten, W. Song, H.P. Wang, and P.C. Ronald. 1995. Construction of a rice bacterial artificial library and identification of clones linked to the Xa21 disease resistance locus. *Plant J* 7:525-533.
42. Wing, R.A., V.K. Rastogi, H.B. Zhang, A. H. Paterson, and S.D. Tanksley. 1993. An improved method of plant megabase DNA isolation in agarose microbeads suitable for physical mapping and YAC cloning. *Plant J* 4: 893-898.
43. Wing. R.A., H.B. Zhang, and S.D. Tanksley. 1994. Map-based cloning in crop plants: Tomato as a model system: Genetic and physical mapping of jointless. *Mol. Gen Genet.* 242: 681-688.
44. Wing, Y. Cheung and M.D. Gale. 1990. The isolation of high molecular weight DNA from wheat, barley and rye for analysis by pulsed-field gel electrophoresis. *Plant Mol. Bio.* 14: 881-888.

45. Woo, S.S., J. Jiang, B.S. Gill, A.H. Paterson, and R.A. Wing. 1994. Construction and characterization of a bacterial artificial chromosome library of *Sorghum bicolor*. *Nucl Acids Res.* 22: 4922-4931.
46. Woo, S.S., V.K. Rastogi, H.B. Zhang, A.H. Paterson, K.F. Schertz, and R.A. Wing. 1995. Isolation of Megabase-Size DNA from Sorghum and Applications for Physical mapping and Bacterial and Yeast Artificial chromosome library construction. *Plant. Mol. Biol. Reprtr* 13 (1): 82-94.
47. Yang, D., A. Parco, S. Nandi, P. Subudhi, H. Zhu, G. Wang, and N. Huang. 1997. Construction of a bacterial artificial chromosome (BAC) library and identification of overlapping BAC clones with chromosome 4-specific RFLP markers in rice. *Theoret. & appl. Genet.* 95 (7): 1147-1154.
48. Yuling, S., M. Valeria, and B. Birren. 1995. Transformation of *Escherichia coli* with large DNA molecules by electroporation. *Nucliec Acids Res.* 23 (1): 1990-1996.
49. Zhang, H.B., S. Choi, S.S. Woo, Z. Li, and R.A. Wing. 1996. Construction and characterization of two rice bacterial artificial chromosome libraries from the parents of a permanent recombinant inbred mapping population. *Molecular Breeding* 2: 11-24 (Choi constructed Lemont, and Zhang constructed Teqing).
50. Zhang, H.B., X.P. Zhao, X.D. Ding, A.H. Paterson, and R.A. Wing. 1995. Preparation of megabase-sized DNA from plant nuclei. *Plant J:* 175-184.
51. Zhang, H.B. and R.A. Wing. 1997. Physical mapping of the rice genome with BACs. *Plant Mol. Biol.* 35 (1-2): 115-127.
52. Zhao, X.P., H.B. Zhang, R.A. Wing, and A.H. Paterson. 1994. A simple method for isolation of Megabase DNA from Cotton. *Plant. Mol. Biol. Raptr.* 12 (2): 110-115.
53. Zhu, H., S.D. Choi, A.K. Johnston, R.A. Wing, and R.A. Dean. 1997. A large-insert (130Kb) bacterial artificial chromosome library of the rice blast fungus *Magnaporthe grisea* genome analysis, contig assembly, and gene cloning. *Fungal Genet. & Biol.* 21 (3): 337-347.
54. Zwick, M.S., M.N. Islamfaridi, D.G. Czeschin, R.A. Wing, G.E. Hart, D.M. Stelly, and H.J. Price. 1998. Physical mapping of the liguleless linkage group in *Sorghum bicolor* using rice RFLP-Selected BACs. *Genetics.* 148 (4) 1983-1992.

بناء وتمييز مكتبات المادة الجينية من أصناف القطن المصري بواسطة العائل البكتيري *E.coli*

أسامة أحمد ممتاز^١ - عبير إبراهيم^١ - إسماعيل كامل^٢ - مجدي مذكور^١

^١ معهد بحوث الهندسة الوراثية الزراعية - مركز البحوث الزراعية - جيزة.
^٢ قسم النبات - كلية العلوم - جامعة القاهرة.

تم بناء وتمييز مكتبات جينية للقطن المصري أصناف جيزة ٧٠ وجيزة ٨٦ وجيزة ٧٥. كذلك تم عزل وتنقية المادة النووية DNA المتميزة بوزن جزيئي عالي وذلك من أنوية الخلية النباتية الموجودة في داخل قطع من الاجاروز كروية الشكل وصغيرة الحجم وهذه تعتبر الخطوة الاساسية في هذا العمل.

وقد تمت دراسة عملية لعدد من المعايير المتحكمة في هذه العملية ومنها الوصول الي الطريقة المثلي لعزل وقطع المادة النووية بأحجام كبيرة متماسكة والقيام باختبارها باستخدام تكنولوجيا ال Chef Gel. وقد قدر حجم ال DNA المعزول ما بين ٢٠٠ الي ٥٠٠ الف قاعدة نووية. وقد تم نقل هذه المادة النووية الي ناقل جيني متميز PBeloBACII الخاص بالعائل البكتيري ال *E.coli* وذلك لاستخدامه في بناء هذه المكتبات الجينية المتماسكة.

وبعد العديد من الدراسات العملية ثبت أن أفضل نسبة من ال Vector : Insert كانت ١٥ : ١ وأن أعلى كفاءة لعملية التحور الوراثي عن طريق اكتساب مادة نووية جديدة تم باستخدام ١,٨ كيلوفولت في عملية التحور بالتيار الكهربائي. أحتوت مكتبات ال BAC بالنسبة للقطن المصري أصناف جيزة ٧٠ وجيزة ٨٦ وجيزة ٧٥ علي ٤٥٢٢٧ - ٤٥٧٤٢ - ٤٦٥٢١ مجموعة كلون وكان متوسط حجم ال DNA ١٠٠ الف قاعدة نيروجينية. وتعتبر هذه المكتبات الوسيلة الاساسية المستقبلية في عزل الجاميع الجينية المسئولة عن أي صفات اقتصادية هامة في القطن المصري.