



## Reproduction Potential of Medaka Fish (*Oryzias profundicola*) as an Endemic Fish in Lake Towuti

Andi Tamsil<sup>1\*</sup>, Nursyahran<sup>2</sup>, Jayadi<sup>1</sup>, Harlina<sup>1</sup>

<sup>1</sup>Faculty of Fisheries and Marine Science Universitas Muslim, Indonesia, Makassar 90231

<sup>2</sup>Institute of Maritime Technology and Business Balikpapan, Makassar 90245

\*Corresponding Author: [andi.tamsil@umi.ac.id](mailto:andi.tamsil@umi.ac.id)

### ARTICLE INFO

#### Article History:

Received: Oct. 1, 2023

Accepted: Jan. 17, 2024

Online: Feb. 24, 2024

#### Keywords:

Medaka fish,  
*Oryzias profundicola*,  
Reproduction potential,  
Fecundity,  
Gonado-somatic index,  
Egg diameter,  
Lake Towuti

### ABSTRACT

This research aimed to determine the reproductive potential of medaka fish (*Oryzias profundicola*), including fecundity, egg diameter, and gonado somatic index (GSI). This research was carried out from January to June 2022. The sampling location was Lake Towuti in South Sulawesi. Medaka fish sample collection was carried out at each research station using a modified bag seser net with a mesh size of  $\frac{3}{4}$  inch, a total length of one fishing gear unit of 15 meters, a height of 1.50 meters, and a bag length of 3 meters.

The fecundity of medaka fish at station 1 (Tanjung Timbala) ranged from 27- 87 eggs ( $56 \pm 15$  eggs) for TKG III and 65- 175 eggs ( $104 \pm 21$  eggs) for TKG IV, while the total fecundity recorded 15,157 eggs. Station 2 (Tanjung Lengkobutangan) fecundity for TKG III ranged from 31- 84 eggs ( $47 \pm 11$  eggs) and TKG IV ranged from 97- 182 eggs ( $153 \pm 20$  eggs), and total fecundity was 11,369 eggs. The diameter of medaka fish eggs in Lake Towuti at all stations was found to range between 0.03 & 1.19mm, with the highest frequency for TKG III in the range of 0.03- 1.08mm, while for TKG IV, it was in the range of 0.15- 1.19mm. The IKG tendency for female fish is relatively greater than that of male fish, except in April and May at station 3 (Tanjung Bakara) and in April at station 5 (Tanjung Tominanga). Reproductive potential is the product of the number of fish and the average fecundity. The reproductive potential of medaka fish at station 1 was 15,157 eggs, station 2 was 11,369 eggs, station 3 was 10,723 eggs, station 4 was 14,842 eggs, and station 5 was 18,385 eggs.

### INTRODUCTION

Sulawesi is one of Indonesia's large islands with high biota richness. This island is included in the Wallacea region, together with the Philippines and Nusa Tenggara, which is a transitional area between Oriental and Australian zoogeography (Whitten *et al.*, 1987). Therefore, there are many endemic types of flora and fauna that attract the attention of biological researchers.

Lake Towuti stores a variety of biodiversity and represents a multifunctional lake, including having the potential for freshwater fisheries resources with food fish and ornamental fish commodities, which have economic value and contribute to the main ecological processes in the aquatic environmental system, controlling the food web as consumers and prey for other organisms, contributing on nutrient cycles, and shaping biophysical habitats through ecological engineering (Villéger *et al.*, 2017).

Based on information in the study of **Hadiaty (2018)**, 17 types of endemic fish were found in Lake Towuti. Meanwhile, **Husnah *et al.*, (2006)** reported that the endemic fish species found included those from the genera *Oryzias*, *Glossogobius*, *Telmatherina*, *Paratherina*, and *Dermogenys*.

Medaka fish (*Oryzias profundicola*) is one of the species of the *Oryzias* genus in Lake Towuti. In the international language, it is called yellow-finned medaka or ricefish, while in the local language, it is called pangkilang fish (**Said *et al.*, 2015**). Medaka fish have important economic and ecological value; therefore, their existence needs to be preserved. To preserve the medaka fish, it is necessary to study its reproductive potential. The data on the reproductive potential of trees can serve as a reference for maintaining its sustainability through domestication. This has been domesticated with different natural feeds on medaka fish (*O. profundicola*) (**Nursyahran *et al.*, 2023**).

Basic information regarding the reproductive potential of fish can be obtained by reviewing the phenomena of gonad development. This phenomenon is used to predict the fish reproductive process, starting from gonad development until the fish spawn and produce seeds. The reproductive potential of each type of fish is different and can be influenced by human intervention, feed quality, and environmental factors. The fish reproductive potential includes spawning patterns, gonado somatic index, fecundity, egg diameter, and maturation time. Some freshwater fish such as *O. kalabu*, a relative of *O. vittatus* (nilem fish), have mature gonads every month, however peak spawning occurs in certain months at the end of the year (**Nasution *et al.*, 2006**).

The distribution of mature egg diameters in the ovary can be used to estimate the frequency of spawning, namely by looking at the mode that is formed. The length of time for spawning can be predicted from the diameter of the eggs. If the fish specimen has a short spawning time, then all the eggs that mature in the ovaries will be the same size. However, if the spawning time of the fish is long or continues over a long period of time, the eggs in the ovaries will have different sizes (**Omar, 2010**).

The aim of the present research was to determine the reproductive potential of freshwater medaka fish (*Oryzias profundicola*), including fecundity, egg diameter and gonado-somatic index (GSI).

## MATERIALS AND METHODS

### Study area

The research was conducted for 6 months from January to June 2022 in Lake Towuti, South Sulawesi, Indonesia. The fish sampling locations were at Tanjung Timbala, Lengkobutanga Cape, Tanjung Bakara, Tanjung Saone, and Tanjung Tominanga (Fig. 1).

### Sample collection

Medaka fish sample collection was carried out at each research station using a modified bag seser net with a mesh size of  $\frac{3}{4}$  inch, a total length of one fishing gear unit of 15 meters, a height of 1.50 meters, and a bag length of 3 meters. The net is stretched at the bottom of the water, and each end is held by a fisherman, then another fisherman guides the fish into the net, and the net is simultaneously lifted to the surface and the entire catch is used as a sample. The location for sampling the medaka fish was determined using the global positioning system (GPS).




Fig. 1. Map of research locations (Nursyahran *et al.*, 2022)

### Fish measurements

The catches obtained from each station were separated by gender and counted. The fish samples were preserved in 4% formalin for  $\pm 1$  hour, and then the fish samples were rinsed using clean water and fixed with 70% alcohol. Subsequently, the caught fish were coded on a label stating the location where it was found.

### Fecundity

The total fecundity was calculated using the direct calculation method, namely all eggs (of gonad maturity levels III & VI: TKG III and TKG IV) in the female's gonads used as samples (Andy Omar *et al.*, 2012).

### Egg diameter

The results of each egg diameter measurement were displayed with a histogram on TKG III and TKG IV. The egg diameter was calculated using the method of Andy Omar *et al.* (2012), as follows:

$$D_s = \sqrt{D_h \times d_v}$$

Where,  $D_s$ = Actual egg diameter (mm),  $D_h$ = Horizontal egg diameter (mm),  $d_v$ = Vertical egg diameter (mm)

### Gonado somatic index

The gonado-somatic index (GSI) was calculated using the formula of Johnson (1971), as follows:

$$GSI = \frac{W_g}{W} \times 100\%$$

Where, GSI= Gonado somatic index (%),  $W_g$ = Gonad weight (g),  $W$ = Total weight (g).

## Reproductive potential

The reproductive potential was calculated based on the percentage of the number of individuals and average total fecundity (FT) in each age group or mode. The reproductive potential was calculated based on **Conand (1987)** method, as follows:

$$PR = N \times FT$$


Where, PR= Reproductive potential; N= Number of individuals, and FT= Average total fecundity.

## RESULTS

### Fecundity

The fecundity of medaka fish at station 1 (Tanjung Timbala) ranged from 27 to 87 eggs ( $56 \pm 15$  eggs) for gonad maturity level III and from 65 to 175 eggs ( $104 \pm 21$  eggs) for gonad maturity level IV, as well as total fecundity was 15,157 eggs. At station 2 (Tanjung Lengkobutangan), fecundity for gonad maturity level TKG III ranged from 31 to 84 eggs ( $47 \pm 11$  eggs) and TKG IV ranged from 97 to 182 eggs ( $153 \pm 20$  eggs), and total fecundity was 11,369 eggs. At station 3 (Tanjung Bakara), fecundity ranged from 32 to 82 eggs ( $48 \pm 15$  eggs), gonad maturity level IV ranged from 58 to 182 eggs ( $105 \pm 37$  eggs), and total fecundity was 10,723 eggs. At station 4 (Tanjung Saone), fecundity ranged from 53 to 87 eggs ( $67 \pm 10$  eggs) for gonad maturity level III, gonad maturity level IV ranged from 81 to 146 ( $99 \pm 13$  eggs), and total fecundity was 14,842 eggs. Additionally, at station 5 (Tanjung Tominanga), fecundity ranged from 43 to 105 ( $73 \pm 17$  eggs) for gonad maturity level III, gonad maturity level IV ranged from 84 to 156 eggs ( $103 \pm 84$  eggs), as well as total fecundity was 18,385 eggs. Overall, the fecundity of medaka fish in Lake Towuti was 70,467 eggs, consisting of 24,292 TKG III eggs, and 46,184 TKG IV eggs.


The relationship between fecundity and total length of medaka fish is at a value of  $R^2 = 0.3872$ , meaning that 38.72% of fecundity is influenced by total length, while the relationship between fecundity and total weight is at a value of  $R^2 = 0.6064$ , meaning that 60.64% of fecundity is influenced by total weight (Fig. 2). The graph show that the heavier the fish, the higher the fecundity. Fish weight correlates with fecundity.



**Fig. 2.** Relationship between fecundity and total length: (a) Weight (b) At all stations

**Egg diameter**


The egg diameter of medaka fish in Lake Towuti at all stations was found to be between 0.03- 1.19mm with the highest frequency for gonad maturity level III in the range 0.03-1.08mm, while for gonad maturity level IV, it was in the range of 0.15- 1.19mm. Based on the analysis of the distribution of egg diameters at all stations, both gonad maturity levels (III and IV) found one peak mode. For all stations (1– 5), the peak mode egg diameter was found in the size range of 0.03- 0.17mm for gonad maturity level III, while for gonad maturity level IV, it was found in the size range of 0.93- 1.07mm (Fig. 3).




**Fig. 3.** The egg diameter of medaka fish for TKG III and IV at all studied stations

### Gonado-somatic index (GSI)

The results of the analysis of the average GSI values for male and female fish at each station based on sampling time are displayed in Fig. (4). Fig. (5) shows that the average GSI of male fish at station 1 during the research ranged from 0.77 to 1.56% and female fish ranged from 1.52 to 1.66%. The female GSI recorded every month was greater than the male GSI. The average GSI of male fish at station 2 was 1.13- 1.50% and female fish was 0.59- 1.67%. The female GSI was greater than the male GSI in each month. The results showed that the average GSI of male fish at station 3 during the research ranged from 0.87 to 1.60%, and female fish ranged from 1.37 to 1.65% (Fig. 4).






**Fig. 4.** The average value of gonado somatic index (%) of male and female medaka fish based on sampling time at station 1 (a) Tanjung Timbala, station 2 (b) Tanjung Lengkobutanga, station 3 (c) Tanjung Bakara, station 4 (d) Tanjung Saone, and station 5 (e) Tanjung Tominanga

The female GSI was larger than the male GSI in January, February, March and June. The data showed that the average GSI of male fish at station 4 during the study ranged from 0.78 to 1.48%, and female fish was at 1.44- 1.61% (Fig. 4). The female GSI was larger than the male GSI at the time of sampling. Furthermore, the results showed that the average GSI of male fish at station 5 during the study ranged from 0.85 to 1.56% and female fish was at 1.40- 1.59%. The female GSI was larger than the male GSI in every month except April (Fig. 4).

Based on the present data, different GSI fluctuations were shown in each month and at each station. The tendency for gonado-somatic index (GSI) of female fish was relatively greater than that of male fish, except in April and May at station 3 and April at station 5 (Fig. 4).

### Reproductive potential

The reproductive potential is the product of the number of fish and the average fecundity. The reproductive potential of medaka fish at station 1 was 15,157 eggs, station 2 was 11,369 eggs, station 3 was 10,723 eggs, station 4 was 14,842 eggs, and station 5 was 18,385 eggs (Fig. 5).



**Fig. 5.** The Reproductive potential of medaka fish in Lake Towuti

## DISCUSSION

Fecundity is a variable of reproductive strategy in fish, in addition to sex ratio, size of first mature gonads, period and type of spawning, and oocyte development (**Gomiero et al., 2008**). Knowledge about the fecundity of a fish is very important since it can be used to evaluate the stock potential, life cycle, cultivation, and management of the species itself (**Hussain et al., 2007**). Fecundity is the number of eggs previously released by female fish during spawning (**Nursyahrhan et al., 2021**).

The fecundity of medaka fish in Lake Towuti based on location and time of research ranged from 27 to 182 eggs. The fecundity of this fish species is smaller than several endemic fish of the same species, including lunjar fish (*O. marmoratus*) ranging from 20- 760 eggs (**Sulistiono, 2012a**), and *O. nigrimas* fish ranging from 143- 243 eggs (**Serdiati, 2019**). In Matano medaka fish (*O. matanensis*) a greater fecundity was recorded compared to the medaka fish (*O. profundicola*), with 22- 180 eggs (**Eragradhini, 2020**). The occurrence of low fecundity in a fish species is thought to be related to its small body shape, as is the case with the beseng-beseng fish in the Maros River (**Jayadi et al., 2016**).

The amount of fecundity in the same species can be influenced by body size, age, environment, and egg diameter. The fish fecundity tends to increase with increasing body size, which is influenced by the amount of food and other environmental factors, such as temperature and season (**Kara & Bayhan, 2008**).

The length of spawning time can be predicted from the frequency of egg diameter measurements. Egg diameter analysis was carried out to determine fish spawning patterns. There are two types of spawning patterns, namely total spawning and partial spawning. Fish ovaries that contain all the same or uniform-sized cooked eggs indicate a short spawning time. On the other hand, a long and continuous spawning time is indicated by the number of fish eggs of different sizes in the ovary (**Katiandagho & Marasabessy, 2017**). Variations in average egg diameter and increase in egg diameter along with increasing total length, total weight, and ovary weight are caused by environmental factors and the level of fish gonad maturity (**Mostafa et al., 2008**).

The peak mode diameter of medaka fish eggs for TKG III was found in the size range of 0.03- 0.15mm, while for TKG IV it was found in the size range of 0.93- 0.1.07mm at all stations. Thus, the results of this study show that the distribution of egg diameter at all stations has only one highest peak mode at TKG III and TKG IV. The distribution of egg diameters varies at each station. The frequency of egg diameter in TKG III decreased with increasing egg diameter, however in TKG IV the frequency of egg diameter increased with increasing egg diameter. The distribution of egg diameter at the level of gonad maturity shows the fish spawning pattern. The frequency distribution of the diameter of medaka fish eggs varies in size at different TKG. Fish with asynchronous ovulators are known as multiple spawners or partial spawning (**Muchlisin, 2014**). Therefore, medaka fish are included in gradual spawning or partial spawning or long spawning. Fish that partially lay eggs have various sizes of eggs in their ovaries, indicating that all the eggs are not ready to be spawned (**Hunter, 1980**). The length of egg-laying time is indicated by the number of differences in egg size in the ovary; therefore it can be said that, the egg diameter at each level of gonad maturity will reflect the spawning pattern (**Nursyahrhan et al., 2021**). The polymodal nature of the egg size frequency distribution over almost the entire length range predicts that the species may spawn one or more times over a long seasonal period (**Usman et al., 2013**).



The frequency distribution of the diameter of medaka fish in the ovaries did not reveal large differences between the levels of gonad maturity. From these results, it is known that the medaka fish is a partial spawning type that can spawn several times in one spawning season. This is found in other types of medaka fish, such as *O. matanensis* (Eragradhini, 2020), *O. marmoratus* (Said & Mayasari, 2020), and *O. woworae* (Firmansyah *et al.*, 2021). Gradual or partial spawning is found in endemic fish, viz. *T. ladigesi* (Andriani, 2000; Nasution *et al.*, 2006; Kariyanti *et al.*, 2014; Jayadi *et al.*, 2016), *T. celebensis* (Nasution *et al.*, 2007; Jayadi *et al.*, 2010), *Glossolepis incisus* (Siby *et al.*, 2009), and *T. bonti* (Nursyahrhan *et al.*, 2021).

The diameter of lunjar fish (*O. marmoratus*) eggs in Lake Towuti is 0.01- 1.10mm (Sulistiono, 2012b), *O. nigrimas* fish individual is 266- 1166 microns (Serdiati 2019), and medaka fish (*O. matanensis*) in Lake Matano is 0.05- 1.15mm (Eragradhini, 2020). In research on *O. marmoratus* fish, *O. nigrimas* and *O. matanensis*, the egg diameter range is slightly smaller than that of *O. profundicola* fish. Several endemic fish in Lake Towuti that are reported to have a larger egg diameter range than medaka fish include the bonti-bonti fish specimen (*Telmatherina bonti*), which has an egg diameter range of 0.33- 1.85mm (Nursyahrhan *et al.*, 2021).

Quantitative changes in the gonads can be identified as an index of gonadal maturity. By monitoring changes in the gonado somatic index from time to time, you can know the size at which fish individual starts to spawn. The IKG value of female medaka fish was higher than that of male medaka fish at all stations (Tanjung Timbala, Tanjung Lengkobutanga, Tanjung Bakara, Tanjung Saone and Tanjung Tominanga). Moreover, the IKG value from the results of this study is in accordance with the statement of Effendie (2002) that the IKG in male fish is generally smaller than in females. Furthermore, the IKG value of the endemic bonti-bonti fish in Lake Towuti is relatively small for male fish compared to female fish (Nasution *et al.*, 2010; Andy Omar *et al.*, 2012). In addition, the IKG value of female beseng-beseng fish is greater than the IKG value of males (Jayadi *et al.*, 2016). The IKG value of female *Glossogobius giuris* fish is higher than that of males in Lake Limboto (Juliana *et al.*, 2018). The IKG value from the results of this research is in accordance with research by Eragradini (2020) who elucidated that female Matano medaka fish individual has a higher IKG value than male fish.

The highest IKG value of male and female medaka fish at station 1 Tanjung Timbala was found in March 2022. Moreover, the highest IKG value of male medaka fish at station 2 Tanjung Lengkobutanga was found in March 2022 and female fish in May 2022. Furthermore, the highest IKG value of male medaka fish was in station 3 Tanjung Bakara was found in March 2022 and females in February and May 2022. In addition, the highest IKG value of male medaka fish at station 4 Tanjung Saone was found in April 2022 and females in February 2022, while the highest IKG value of male medaka fish was recorded at station 5 Tanjung Tominanga during February 2022 and females during March 2022. Thus, the spawning peak of male and female medaka fish in Lake Towuti ranged from February to March 2022.

The gonado somatic index value of medaka fish is quite high in both male and female fish. This shows that this fish species can spawn in almost all stations. However, station 3 Tanjung Bakara and station 5 Tanjung Tominanga are thought to be the spawning places for medaka fish in Lake Towuti since fish were found with the highest gonado somatic index values for both male and female fish, and it is possible that the microhabitat at each station is different.

Several endemic fish species are reported to have different spawning peaks. The peak of bonti-bonti fish spawning in Lake Towuti is in May and November (Nasution *et al.*, 2010). The peak of the spawning of *O. marmoratus* fish is in July (Sulistiono, 2012b). The peak of peras fish spawning in the Menduk River occurs from May to August (Suhendra *et al.*, 2017). Moreover, the peak spawning of *O. nigrimas* fish occurs in June and August (Serdiati, 2019). Furthermore, the peak spawning of *O. matanensis* fish occurs from August to September (Eragradhini, 2020). Additionally, the peak spawning of Pleco fish, *Pterygoplichthys pardalis* in the Ciliwung River is in February and May (Elfidasari *et al.*, 2022).

## CONCLUSION

Based on the research results, it can be concluded that the reproductive potential of the medaka fish (*O. profundicola*) in Lake Towuti, an endemic species, is robust. This conclusion is based on the fecundity values, egg diameter, gonado somatic index (GSI), and the results of calculations of reproductive potential. These findings indicate that the population is well-protected from extinction.

## REFERENCES

- Andriani. (2000). Bioekologi, morfologi, kariotip dan reproduksi ikan hias rainbow Sulawesi (Telmatherina ladigesii) di Sungai Maros, Sulawesi Selatan. IPB.
- Andy Omar, S. Bin. (2010). Aspek reproduksi ikan nilem, *Osteochilus vittatus* (Valenciennes 1842) di Danau Sidenreng, Sulawesi Selatan. Jurnal Iktiologi Indonesia 10(2): 111-122.
- Andy Omar, S.; Salam, R.; Umar, M.T. and Kune, S. (2012). Biologi reproduksi ikan endemik bonti-bonti, *Paratherina striata* Aurich, 1935 di Danau Towuti. Seminar Nasional Tahunan IX Hasil Penelitian Perikanan Dan Kelautan, July 2012, BP-03.
- Conand, F. (1987). Biologie et Ecologie des Poissons Pelagiques du Lagon de Nouvelle – Caledonie Utilisable Comme Appat Thonier. These de Doctorat d’Etat, Univ. Bretagne Occidentale, Brest. 235 pp.
- Effendie.M.I. (2002). Biologi perikanan. Yayasan pustaka nusatama.
- Elfidasari, D., Puspaningtias, F.C. and Fahmi, M.R. (2022). Reproductive Biology Pleco (*Pterygoplichthys pardalis* Castelnau 1855) in Ciliwung River. Jurnal Pembelajaran Dan Biologi Nukleus, 8(2): 247–262.
- Eragradhini, A. R. (2020). Ekobiologi dan Reproduksi Ikan Matano Medaka, *Oryzias matanensis* (Aurich, 1935) di Danau Towuti Sulawesi Selatan. Universitas Hasanuddin Makassar.
- Firmansyah, M. A., Mustahal, M., Syamsunarno, M. B. and Herjayanto, M. (2021). Optimization of reproduction of ricefish endemic to Southeast Sulawesi *Oryzias woworae* Parenti & Hadiaty, 2010 through different sex ratios in spawning. Jurnal Iktiologi Indonesia, 21(3), 235–251.
- Gomiero, L.M.; Garuana, L. and Braga, F.M.S. (2008). Reproduction of *Oligosarcus hepsetus* (Cuvier, 1829) Characiformes in the Serra do Mar State Park, São Paulo, Brazil. Brazilian Journal of Biology, 68(1): 187–192.

- Hadiaty, R. K. (2018).** Status taksonomi iktiofauna endemik perairan tawar Sulawesi. *Jurnal Iktiologi Indonesia*, **18**(2): 175-190.
- Hunter, J.R. (1980).** The Feeding Behavior and Ecology of Marine Fish Larvae. *Fish Behavior and Its Use in the Olpture and Culture of Fishes*, 287–327.
- Husnah, S. and Makmur, S. (2006).** Riset keanekaragaman hayati dan bahan perumusan pengelolaan jenis ik an endemik perairan pedalaman di Sulawesi. Laporan Teknis. PusatRiset Perikanan Tangkap.
- Hussain, M.A.; Khatun, M.R. and Hossain, M.A. (2007).** On the fecundity and sex-ratio of *Botia Dario* (Hamilton) (Cypriniformes : Cobitidae). *University Journal of Zoology, Rajshahi University*, **26**: 27–29.
- Jayadi, J.; Hadijah, S.; Tang, B. and Husma, A. (2016).** Biologi reproduksi ikan beseng-beseng (*Marosatherina ladigesi* Ahl, 1936) di beebraja sungai di Sulawesi Selatan. *Jurnal Iktiologi Indonesia*, **16**(2): 185–198.
- Jayadi, J.; Hamal, R. and Arifuddin, (2010).** Reproduksi ikan endemik rainbow Sulawesi *Telmatherina celebensis* di Danau Matano Sulawesi Selatan. *Torani (Jurnal Ilmu Kelautan Dan Perikanan)*, **20**(1): 44–48.
- Johnson, J.E. (1971).** Maturity and fecundity of threadfin shad, *Dorosoma petenense* (*Gunther*), in Central Arizona reservoirs. *Transaction of the American Fisheries Society*, **100**(1): 74-85
- Juliana, Koro, Y., and Lamadi, A. (2018).** Domestikasi dan Apalikasinya Terhadap Ikan Manggabei (1st ed.)
- Kara, A. and Bayhan, B. (2008).** Length-weight and length-length relationship of the bogue, *Boops boops* (Linneaus, 1758) in Izmir Bay (Aegean Sea of Turkey). *Belgian Journal of Zoology*, **138**(2): 154–157.
- Kariyanti, Omar, S.B.A. and Tresnati, J. (2014).** Analisis Fekunditas dan Diameter Telur Ikan Beseng-Beseng (*Marosatherina ladigesi* Ahl, 1936) Sungai Pattunuang Asue dan Sungai Bantimurung, Kabupaten Maros, Sulawesi Selatan. Simposium Nasional I Kelautan Dan Perikanan, November, 1–10.
- Katiandagho, B. and Marasabessy, F. (2017).** Potensi Reproduksi, Pola Pemijahan Serta Alternatif Pengelolaan Ikan Kembung Laki-Laki (*Rastrelliger kanagurta*) Di Sekitar PesisirTimur Perairan Biak. *Agrikan: Jurnal Agribisnis Perikanan*, **10**(2): 51.
- Mostafa, Z.; Haque, M.R.; Afsana, D. and Tareque, A.M.H. (2008).** Fecundity of *Liza parsia* (Hamilton) and relationship between length-weight and fecundity. *Fisheries (Bethesda)*, **16**(Salojarvi 1987): 65–70.
- Muchlisin, Z.A. (2014).** A General overview on some aspects of fish reproduction. *Aceh International Journal of Science and Technology*, **3**(1): 43–52.
- Nasution, S.H.; Muschsin, I. and Sulistiono, S. (2010).** Potensi rekrut Ikan Bonti-Bonti (*Paratherina striata* Aurich) Di Danau Towuti, Sulawesi Selatan. *BAWAL Widya Riset Perikanan Tangkap*, **3**(1): 45. <https://doi.org/10.15578/bawal.3.1.2010.45-55>
- Nasution, S.H.; Lukman, S.S D. and Fauzi, T.H. (2006).** Aspek Reproduksi Ikan Beseng Beseng (*Telmatherina ladigesi* Ahl) darl beberapa sungai dl sulawesi selatan. *Prosiding Seminar Nasional Ikan I V, Jatiluhur*, 29-30 Agustus (2006), pp.: 83-93
- Nasution, S.H.; Sulistiono, S.; Sjafei, D.S. and Haryani, G.S. (2007).** Distribusi Spasial dan Temporal Ikan Endemik (*Telmatherina celebensis* Boulenger) di Danau Towuti

Sulawesi Selatan. Jurnal Penelitian Perikanan Indonesia, **13**(2): 95.

- Nursyahran, N.; Jayadi, J.; Tamsil, A. and Harlina. (2023).** Domestication of yellow-finned medaka fish (*Oryzias profundicola*). ACL Bioflux, **16**(1): 524–533.
- Nursyahran, N.; Kariyanti, K.; Ilmiah, Jayadi, J. and Yusuf, A. (2021).** Fecundity and egg diameter of endemic fish ( *Telmatherina bonti* Weber and De Beaufort , 1922 ) from Towuti Lake , South Sulawesi , Indonesia. International Journal of Fisheries and Aquatic Studies, **9**(3), 378–382.
- Said, D. S. and Hidayat. (2015).** 101 Ikan Hias Air Tawar. LIPI Press
- Said, D.S. and Mayasari, N. (2020).** *Oryzias mamoratus*, Si Marmer dari Danau Towuti, Sulawesi Selatan. WARTA IKTILOGI, **4**(3): 19–23.
- Serdiati, N. (2019).** Karakterisasi ikan padi oryzias nigrimas (kottelat, 1990) sebagai upaya pengelolaan ikan endemik di danau poso sulawesi tengah. Universitas Brawijaya.
- Siby, L. S., Rahardjo, M. F. and Sjafei, D. S. (2009).** Biologi Reproduksi Ikan Pelangi Merah (*Glossolepis incisus*, Weber 1907) di Danau Sentani. Jurnal Iktiologi Indonesia, **9**(1), 49–61
- Suhendra, C.; Utami, E. and Umroh, U. (2017).** Reproductive Biology Of fish Keperas (*Cyclocheilichthys apogon*) in Menduk River, Bangka Regency. Akuatik Jurnal Sumberdaya Perairan, **11**(1): 1–11.
- Sulistiono, S. (2012a).** Reproduction of tank goby (*Glossogobius giuris*) in Ujung Pangkah waters, East Java. Jurnal Akuakultur Indonesia, **11**(1): 64-75.
- Sulistiono, S. (2012b).** Reproduksi Ikan Lunjar (*Oryzias marmoratus*) di Danau Towuti Sulawesi Selatan. Agrisains, **13**(1): 55–65.
- Usman, B.I.; Amin, S.M.N.; Arshad, A. and Aminur Rahman, M. (2013).** Fecundity and egg size of grey-eel catfish, *Plotosus canius* (Hamilton, 1822) from the coastal waters of Kampong Telok, Negeri Sembilan Peninsular Malaysia. Asian Journal of Animal and Veterinary Advances, **8**(2): 348–354.
- Villéger, S.; Brosse, S.; Mouchet, M.; Mouillot, D. and Vanni, M.J. (2017).** Functional ecology of fish: current approaches and future challenges. Aquatic Sciences, **79**(4): 783–801.
- Whitten, A.J.; Bishop, K.D.; Nash, S.V. and Clayton, L. (1987).** One or more extinctions born Sulawesi, Indonesia? Conservation Biology, **1**(1): 42–48.