Contents:

Letter from the Editor	2
Bilateral large ovarian lymphoma presenting by abdominal enlargement (Case Report) Ashraf Sobhi Abolouz, Ahmed Hussein Mohamed	3
Obstetric outcome of cervical cerclage among pregnant women in Mansoura University Hospital Asmaa Mohamed Gomaa, Elsaied Mohamed Abdelhady, Ahmed Fathy Gibreel, Mohamed Sayed Abdelhafez.	6
Fetal middle cerebral artery Doppler in post term pregnancy: a predicting factor for induction of labor outcome Doaa Shafie Belal, Maged Al Mohammady, Somaya Hisham Mohamed, Mostafa Mahmoud	14
COVID-19 disease effect on ovarian reserve in women of reproductive age: an analytical before-and-after COVID-19 study Mohamed K. Etman, Marwa S. Sofy, Abd El-Samie Abd El-Mneim Abd El-Samie.	23
Breastfed Impact on Intelligence Quotient in Children Attended To Pediatric Clinic in Al-Hussien Hospital at Cairo, Egypt Mohammed Mahmoud Abd Alrahman Hamouda.	32
The Correlation between Duration of Fetal Extraction during Cesarean Section and Development of Transient Tachypnea of Newborn Mohamed A. Hussein, Wesam K.M Hassan, Amal Kotb Abdallah.	38

Trans Obturator Tape (TOT) Procedure in cases of Stress Urinary Incontinenque: Patients' Satisfaction and Cure Rates (Retrospective Analysis)	
Dina Yahia Mansour, Mohammed Ahmed Elkadi, Mohammed Elsayed Bassam, Alhassan Mohammed Khadr.	_49
Third Trimestric Ultrasound Diagnosis of Placenta Accreta Spectrum and Correlation of The Finding with Figo Grading System Mona Fouad, Mona M. AboulGhar, Ghada Abouseriae, Ahmed Elminawy.	64
The impact of self-administered vaginal isonicotinic acid hydrazide (INH) administration 12 hours prior to levonorgestrel-releasing intrauterine system in women delivered only by elective cesarean section: A randomized double blinded clinical trial Hany F. Sallam, Nahla W. Shady, Ahmed A. Taha.	72
Impact Of Positive GCT With Negative Oral GTT On Perinatal Morbidity: Prospective Cohort Study Dina Yahia Mansour, Noha Emam Youness, Snosi MS.	81
Patient satisfaction and Quality of life after dienogest treatment versus surgical excision of ovarian endometrioma Khaled A. Atwa, Zakia M. Ibrahim, Eman M. El Bassuony, Omima T. Taha, Asmaa M. Elgedawy.	89
Inhibin A may be the Black Horse for Determination of the Optimal Triggering Time and Decision-making for Oocyte Retrieval Youssef Abdel Zaher, Omnia Youssef Habashy, Hamasat A Alnoury, Wagdy Megahed Amer.	98

Letter from the Editor:

Dear esteemed colleagues,

Warm greetings

We welcome your comments as well as the scientific activity to be incorporated in the upcoming issues. Very important subjects are included in this issue: Planned placement of cervical cerclage in is mainly either history-indicated and/or prophylactic, especially in multifetal gestations. It appears to be effective in reducing the incidence of mid trimester miscarriage and preterm birth in high-risk cases.

MCA-PI evaluation prior to induction of labor (IOL) is a useful tool in prediction of IOL outcome. Lower PI values may predict successful outcome. CL assessment and EFW are other factors that may predict the outcome of IOL

It was found that COVID-19 has no effect on ovarian reserve.

A study compared intelligence level between breastfed and non-breastfed children, concluded that there was significant relation between high Wechsler Intelligence Scale score and breastfed.

The surgical outcome and intraoperative FIGO grading are strongly correlated with the presence of placental lacunae, loss of clear zone and myometrial thinning and moderately correlated with Placental bulge

A positive oral glucose challenge test only without gestational diabetes is a risk factor for perinatal morbidity like LGA and NICU admission, so early screening for GDM is advisable.

Best regards.

Aboubakr Elnashar MD Chief Editor of EFSSJ Prof. obs Gyn. Benha university, Egypt elnashar53@hotmail.com