



Microbes and Infectious Diseases

Journal homepage: <https://mid.journals.ekb.eg/>

Original article

Bacterial respiratory infection epidemiology and antibiogram profile in patients attending in regional hospital, Meknes, Morocco

Najia El Hamzaoui* ^{1,2}, Fatima El Kamari ³, Anjoud Harmouzi ⁴, Samira Jaouhar ⁵

1- Laboratory of medical analyses, Meknes's Hospital, Health Regional Direction of Fez-Meknes

2- Faculty of Sciences, Moulay Ismail University, Meknes, Morocco

3- Laboratory of Natural Substances, Pharmacology, Environment, Modeling, Health and Quality of Life (SNAMOPEQ), Faculty of Sciences Dhar El Mahraz, Sidi Mohamed Ben Abdellah University, Fez, Morocco

4- Laboratory of Agrophysiology, Biotechnology, Environment and Quality, Faculty of Sciences, Ibn Tofail University, Kenitra, Morocco

5- Laboratory of Microbial Biotechnology & Bioactive Molecules, Faculty of Sciences and Techniques, Sidi Mohamed Ben Abdellah University, PB 2202, Fez, Morocco

ARTICLE INFO

Article history:

Received 16 February 2024

Received in revised form 19 March 2024

Accepted 23 March 2024

Keywords:

Respiratory system
Microbial infection
Hospital
Retrospective study

ABSTRACT

Objective: The main goal of this current study was to assess the prevalence of these infections in the outpatient setting as well as in hospitalized patients at Mohammed V hospital in Meknes, Morocco. **Method:** The epidemiological and microbiological characteristics of respiratory infections mentioned in the registers of the laboratory of microbiology at Mohammed V hospital, Meknes, were collected on pre-established exploitation sheets for each case throughout a 3-year period (2017-2019). **Results:** A total of 7863 patients diagnosed with respiratory infections were admitted to various hospital departments. During 2017, the percentage of hospitalized patients was equivalent to 33.6% of cases, and during the year 2018, this percentage increased by roughly 4.61% before decreasing by 10.03% in 2019. Males were the most affected with a proportion of 61%, while females did not exceed 31%. The pulmonology department had the highest proportion of hospitalized patients (62.60%), followed by the medicine department (14.63%), and gastroenterology (13.55%). The seasonal distribution peaks in spring with a percentage of 37.6%. According to the bacteriological examination, 20 patients (5.42%) had a bacterial respiratory infection. The biochemical classification of the isolated bacterial species revealed the presence of three bacterial species with a high isolation frequency of *Streptococcus pneumoniae* (85%), followed by *Escherichia coli* (10%) and *Staphylococcus aureus* (5%). **Conclusion:** Respiratory infections are widespread throughout the spring season, with male being the most affected. Monitoring and characterization of epidemiological and microbiological data are still required to prevent and mitigate the burden of these infections on patients, healthcare systems, and society.

Introduction

Bacterial infections constitute a serious challenge that threatens the public health and healthcare systems [1-3]. Conventional treatment of

bacterial infections using chemotherapeutics accentuates the emergence and development of bacterial resistance [4]. Bacterial infections may affect different parts of the respiratory tract. They

DOI: 10.21608/MID.2024.270547.1803

* Corresponding author: Najia El Hamzaoui

E-mail address: najia.elhamzaoui@gmail.com

© 2020 The author (s). Published by Zagazig University. This is an open access article under the CC BY 4.0 license <https://creativecommons.org/licenses/by/4.0/>.

are ubiquitous and commonly affected both adults and children, causing mortality and morbidity [4,5]. Infections are most usually in the respiratory tract and according to the current estimates; pneumonia accounts for approximately 15 % of global fatalities in children under the age of five [6].

Furthermore, the healthcare acquired infections (HAIs) is higher in developing countries than European and American countries. In fact, these infections can arise at any age and in various clinical situations, including immunosuppression, genetic background (cystic fibrosis), intubated-ventilated patients, and healthy individuals. Particularly, the prevalence of HAIs is highly widespread in intensive care units [7]. As a result, they have a significant economic influence and carry a significant weight in our society. The frequency of consultations by general practitioners for respiratory tract infections outnumbers all other reasons combined [8-10].

The direct contact with the environment and permanent exposure to microorganisms suspended in the air can explain the frequency of respiratory tract infections. Pathogenic bacteria can affect healthy subjects, but infections occur in case of defense system damaged. The infection can be located at any level of the respiratory tract and the topography determines the clinical manifestations [11]. The clinical syndromes vary according to their location. The clinical manifestations of the respiratory tract infections also depend on the causative agent. The viruses are particularly important in the upper airway infections and are responsible for the most cases of the pharyngitis [12]. The bacteria are mainly responsible for the otitis media, the pharyngitis, the epiglottitis, the bronchitis and the pneumonia. Presumably, the scarcity of data on the real burden of bacterial infections and its prevalence and epidemiology in Morocco is a key stone of the preparation of the present study.

The main purpose of the present study was to establish the bacterial infection distribution according to sex, department and season, as well as the resistance of isolated strains against different used antibiotics. It is worth to be noted that this is the first study designed to examine the respiratory infections and antibiotic resistance of strains isolated from outpatients and hospitalized patients in Mohamed V Hospital of Meknes over a three-year period (2017-2019). The results of the present study

will be of paramount importance to take the right and preventive actions to avoid HAIs.

Material and methods

Study setting

According to the last Moroccan territorial division of 2015, Fez-Meknes is one of the twelve new regions of Morocco with 4236892 inhabitants, which represents approximately 13% of the overall population of the country [13]. The region is spread over an area of order of 40075 km², which corresponds to 5.7% of the total area of the country.

Hospital Mohamed V is a provincial reference center for care and consultation. It is one of the most important hospital structures in the Fez-Meknes region. It has a functioning bed capacity of 378 beds and provides diagnostic services and specialized care in 22 disciplines.

Methodology

The current study focuses on cases of respiratory infections collected in the Mohamed V hospital laboratory over a three-year period (2017-2019). For this purpose, different clinical (acute infections, angina, acute bronchitis ...), biochemical and microbiological variables that are noted in the registers of Microbiology laboratories of Hospital of Meknes are collected on pre-established sheets for each case.

Bacterial isolation and bacterial resistance frequency

7863 patients attending the outpatient of different departments of Mohamed V hospital diagnosed with respiratory infections were involved in the preparation of the current study.

Clinical sampling

Respiratory infection was carefully diagnosed. Pleural fluid, nasal sections, throat and protected distal sampling were collected from different patients and sent for further microbiological analysis [14].

The germs were isolated using several culture mediums, including chapman agar, cysteine-lactose-electrolyte-deficient agar and chocolate agar. The microorganisms were identified based on cultural, morphological, and biochemical characteristics, supplemented with other assays, including Gram coloration, catalase, cytochrome oxidase and emission of gas bubbles [15].

Antibiogram

The antimicrobial susceptibility assays were conducted on Muller-Hinton agar according to

the method of the Kirby-Bauer disk diffusion [16]. The antibiotic agents tested were: Amoxicilline (AML) 25 µg, Amoxicillin+clavulanic acid 25/20µg (AMC), cefoxitin 10µg (FOX), ceftazidime 30µg(CAZ), cefotaxime 30µg (CTX), imipenem 10µg (IMP), trimethoprim/sulfamethoxazole 1.25/23.75µg(SXT), ciprofloxacin 5µg (CIP), amikacin 30µg (AK), penicillin G 6µg (PenG), erythromycin 30µg (ERY), tobramycin 10µg (TOB), tetracycline 30 µg(TET), colistin 50µg (CL), vancomycin 30µg (VAN), antibody antibiotic conjugate (TAC), fusidic acid 10µg (FA), levofloxacin 5µg (LE), gentamicin 10µg (GEN), rifampicin 30µg (RA), and netilmicin 30µg (NET). Resistance results were evaluated in accordance with Clinical and Laboratory Standards Institute [17].

Ethics and confidentiality

The protocol for the study was approved by the ethics commission of the Regional Delegation of Public Health of the Littoral. The confidentiality of the information was respected since the names of the patients did not appear on the study documents.

Statistical analysis

The statistical analysis was performed by multiple comparison (Tuckey test) at a significant level of 95% ($p < 0.05$). The findings were executed using Graph Pad Prism 5 software.

Results

Distribution of diagnosed patients according to gender

The current study included 7863 patients who were hospitalized and tested for respiratory infections during 2017-2019. During 2017, the percentage of hospitalized patients was equivalent to 33.6% of cases, and during the year 2018, this percentage increased by roughly 4.61% before decreasing by 10.03% in 2019.

It is clearly seen that men represent the highest proportion of patients with a percentage of 61%, whereas females represent 39% of all hospitalized patients.

Distribution of diagnosed patients according to the hospital department

Figure 1 displays the obtained results of frequency of distribution of respiratory infections among all patients. The treatment of the obtained results revealed that the highest frequency was

recorded in the department of pulmonology with a percentage of 62.60% followed by department of internal medicine (14.63%) and the department of gastroenterology (13.55%). Whereas, the lowest frequencies were registered in both departments of neurology and emergency with proportions of 0.27% for each.

The results of the seasonal distribution are presented in **Figure 2**. The results are expressed as the average infection proportions of each season during the three-years. The analysis of results showed that the highest percentage of patients affected was registered during spring season with a value of 37.6% followed by winter season with a percentage of 27.1%. Whereas, the lowest percentage of patients affected according season was registered during autumn with a value of 15.4%.

Bacteriological analysis

The results of bacteriological analysis are depicted in **Figure 3**. Microbial identification assays revealed that 90% of isolates are Gram-positive bacteria. It is clearly seen that *Streptococcus pneumoniae* represented 85% of all bacterial strains identified followed by *Escherichia coli* and *Staphylococcus aureus* with proportions of 10% and 5%, respectively.

During the three years and among the 10 hospital departments where the infections were identified, we remark that 6 departments (60%) had bacterial infections. Most of these infections were concentrated in the pulmonology department at a frequency of 65% (**Table 1**).

S. pneumoniae is more frequently isolated in pulmonology department. *E. coli* is isolated only in two departments (Pulmonology and gastroenterology). Whereas, *S. aureus* was isolated only in the pediatrics department.

Bacterial resistance frequency

The sensitivity of bacterial strains isolated from different patients diagnosed at different Mohamed V hospital departments is presented in **Table 2**. The results showed that all three bacterial strains isolated from patients were resistant to at least five antibiotics. *Escherichia coli* was the most resistant among the isolated bacterial strains with frequencies of 100%, 85%, 84%, 75%, 45%, and 43% against AML, CIP, AMC, FOX, CAZ, and CTX, respectively. However, *S. pneumoniae* and *S. aureus* showed mild resistance against the different antibiotics tested with a percentage that did not exceed 59%.

Table 1. Frequency of bacterial species according to the hospital services.

Department	Number of bacterial species			
	<i>S. pneumoniae</i>	<i>E. coli</i>	<i>S. aureus</i>	Total (%)
Pulmonology	12	1	0	13 (65%)
Pediatrics	1	0	1	2 (10%)
Surgery	2	0	0	2 (10%)
Cardiology	1	0	0	1 (5%)
Medicine	1	0	0	1 (5%)
Gastroenterology	0	1	0	1 (5%)

Table 2. The resistance frequency of different bacterial strains isolated from patients diagnosed with bacterial respiratory infections.

	AML	AMC	FOX	CAZ	CTX	IMP	SXT	CIP	AK	PenG	ERY	TOB	TET	CL	VAN	TAC	FA	LE	GEN	RA	NET
<i>Escherchia coli</i>	100	84	75	45	43	0	57	85	42	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
<i>S. pneumoniae</i>	28	ND	ND	ND	19	ND	ND	ND	ND	52	59	31	17	8	ND	ND	ND	ND	ND	ND	ND
<i>S. aureus</i>	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	0	0	54	50	45	17	11

ND: Not determined




Figure 1. Distribution of the patients according to the hospital department.**Figure 2.** Seasonal distribution of respiratory infections according to the seasons.

Figure 3. Distribution of identified bacterial species.

Discussion

Respiratory infections are a serious challenge of human health and healthcare systems. They can have very varied clinical manifestations depending on the site of infection, patient dietary status, pathophysiology, and immunological system. Respiratory infections are one of the top causes of death worldwide. It is the primary reason for a consultation and prescription in the practice of general medicine. The prevalence of these infections in populations varies according to different factors, including the year, the country, the monitoring systems, and the level of health prevention [18]. Within this framework, the present study was undertaken for the first time to study the bacterial respiratory infection epidemiology in Mohamed V hospital during three years (2017-2019). The findings revealed that the males were the most individuals affected by bacterial respiratory infections. Furthermore, *S. pneumoniae* was the main responsible pathogenic bacteria of bacterial respiratory infections, which presents mild resistance against numerous antibiotic agents. The ministry of health reported that respiratory infections are a serious health challenge and represented approximately 5.5 million medical consultations [19].

The monthly and seasonal distribution of these infections at Mohammed V Hospital in Meknes revealed that respiratory infections are particularly common in the spring, especially in March, April and May. Furthermore, Elhella and Lahrach demonstrated that respiratory infections are

extremely common throughout the winter season, particularly January, February and March [20]. The obtained results from this study are in accordance with those of a retrospective study conducted at the military hospital, Rabat. The sampling method used to examine respiratory infections is always determined by the site of infection and the patient's terrain. The authors reported that more than 90% of the proven microbiological infections are caused by *S. pneumoniae*, *H. influenzae*, *M. pneumoniae*, *L. pneumophila*, *C. pneumonia* and *S. pneumonia*, which are the most common cause of hospitalization [21,22]. In fact, the results of the present study demonstrated that the *S. pneumoniae* is the most frequently isolated (85%). It has been found that *S. pneumoniae* is pathogenic bacterial strain causing pneumonia, otitis media, sepsis, and meningitis [23]. In the same context, numerous studies reported the same findings reporting that *S. pneumoniae* is the main reason of bacterial respiratory infections with a proportion of 50.5% [24,25]. Interestingly, the bacterial strains detected and identified showed high variability of resistance against several antibiotics, particularly *Escherichia coli* (Table 2). Pathogenic bacteria develop antibiotic resistance throughout different strategies, including target modification and efflux transporters [23]. Due to the occurrence of negative side effects and elevating risk of resistance to the antibacterial agents, there has been a real need to in the quest of natural alternatives.

Conclusion

The overall outcomes of the current study revealed that the respiratory infections are common throughout the spring season, with men being the

most impacted. Respiratory infections are primarily caused by *S. pneumoniae*, which affects the respiratory tract. The current study investigated the progression and prevalence of the respiratory infections at Mohamed V hospital, Morocco. These infections are widespread in the spring months, when the majority of the patients have been admitted to the pulmonology department.

Author contributions

All authors have read and approved the final version of the manuscript for publication.

Financial support and sponsorship

The study received no financial support or sponsorship.

Conflicts of interest

The authors declare no conflicts of interest related to this research.

References

- 1- Jroundi I, Mahraoui C, Benmessaoud R, Moraleda C, Benjelloun B, Bassat Q. Knowledge gaps on paediatric respiratory infections in Morocco, Northern Africa. Arch Pub Health 2015;73:28.
- 2- Aguilar GR, Swetschinski LR, Weaver ND, Ikuta KS, Mestrovic T, Gray AP, et al. The burden of antimicrobial resistance in the Americas in 2019: a cross-country systematic analysis. The Lancet Regional Health–Americas 2023;25.
- 3- Timpau AS, Miftode RS, Costache II, Petris AO, Miftode IL, Gheorghe L, et al. An Overview of the Impact of Bacterial Infections and the Associated Mortality Predictors in Patients with COVID-19 Admitted to a Tertiary Center from Eastern Europe. Antibiotics 2023; 12: 144.
- 4- Li T, Wang Z, Guo J, de la Fuente-Nunez C, Wang J, Han B, et al. Bacterial resistance to antibacterial agents: Mechanisms, control strategies, and implications for global health. Sci. Total Environ 2023; 860: 160461.
- 5- Ighid N, El Akil S, Aghrouch M, Hassaini N, Izaabel EH. Acute Respiratory Infections Epidemiology and Etiology in Hospitalized Moroccan Children under 15 Years. IJMS 2020; 7.
- 6- Liu L, Johnson HL, Cousens S, Perin J, Scott S, Lawn JE, et al. Global, regional, and national causes of child mortality: an updated systematic analysis for 2010 with time trends since 2000. The lancet 2012; 379: 2151–61.
- 7- El mekes A, Zahlane K, Ait said L, Tadlaoui Ouafi A, Barakate M. The clinical and epidemiological risk factors of infections due to multi-drug resistant bacteria in an adult intensive care unit of University Hospital Center in Marrakesh-Morocco. J Infect Public Health 2020; 13: 637–43.
- 8- Passali D, Arezzo MF, De Rose A, De Simone G, Forte G, Jablko-Musial M, et al. Benzylamine hydrochloride for the treatment of sore throat and irritative/inflammatory conditions of the oropharynx: a cross-national survey among pharmacists and general practitioners. BMC Prim Care 2022; 23: 154.
- 9- Tataru AM. The Infectious Diseases Consult Handbook: Common Questions and Answers. Springer Nature 2023.
- 10- Turner E, Johnson E, Levin K, Gingles S, Mackay E, Roux C, et al. Multi-disciplinary community respiratory team management of patients with chronic respiratory illness during the COVID-19 pandemic. NPJ Prim Care Respir Med 2022; 32: 26.
- 11- Reis JGC, Reis CSM, Costa DCSD, Lucena MM, Schubach ADO, Oliveira R

- DVC, et al. Factors associated with clinical and topographical features of laryngeal tuberculosis. *PloS one* 2016; 11: e0153450.
- 12- Lyczak JB, Cannon CL, Pier GB. Lung Infections Associated with Cystic Fibrosis. *Clin Microbiol Rev* 2002; 15: 194–222.
- 13- HCP. Glossaire | Site institutionnel du Haut-Commissariat au Plan du Royaume du Maroc. Available at: https://www.hcp.ma/glossary/_gw.html. Accessed March 13, 2024.
- 14- Loens K, Van Heirstraeten L, Malhotra-Kumar S, Goossens H, Ieven M. Optimal Sampling Sites and Methods for Detection of Pathogens Possibly Causing Community-Acquired Lower Respiratory Tract Infections. *J Clin Microbiol* 2009; 47: 21–31.
- 15- Hindi NK. In vitro antibacterial activity of aquatic garlic extract, apple vinegar and apple vinegar-garlic extract combination. *American J Phytomed Clin Therap* 2013; 1: 42–51.
- 16- Bauer AW, Kirby WMM, Sherris JC, Turck M. Antibiotic Susceptibility Testing by a Standardized Single Disk Method. *American J Clin Pathol* 1966; 45(4-ts): 493–6.
- 17- Kibret M, and Abera B. Prevalence and antibiogram of bacterial isolates from urinary tract infections at Dessie Health Research Laboratory, Ethiopia. *Asian Pac J Trop Biomed* 2014; 4: 164–8.
- 18- Couret NC. Impact d'un protocole d'observation de pneumonies aiguës communautaires hospitalisées: à propos de 144 patients hospitalisés au CH de Saint-Gaudens. PhD Thesis 2007.
- 19- Initiative ST. Prise en charge combinée des maladies respiratoires et de la tuberculose au Maroc. Organisation mondiale de la Santé 2001.
- 20- El Hella N and Lahrach S. Etiologie bactérienne des infections respiratoires aiguës chez les enfants. FSSI 2007.
- 21- Lin SH, Lai CC, Tan CK, Liao WH, Hsueh PR. Outcomes of hospitalized patients with bacteraemic and non-bacteraemic community-acquired pneumonia caused by *Streptococcus pneumoniae*. *Epidemiol Infect* 2011; 139: 1307–16.
- 22- Tan TQ, Mason Jr EO, Wald ER, Barson WJ, Schutze GE, Bradley JS, et al. Clinical characteristics of children with complicated pneumonia caused by *Streptococcus pneumoniae*. *Pediatrics* 2002; 110: 1–6.
- 23- Li L, Ma J, Yu Z, Li M, Zhang W, Sun H. Epidemiological characteristics and antibiotic resistance mechanisms of *Streptococcus pneumoniae*: An updated review. *Microbiol Res* 2023; 266: 127221.
- 24- Ben-Shimol S, Greenberg D, Hazan G, Shemer-Avni Y, Givon-Lavi N, Dagan R. Seasonality of both bacteremic and nonbacteremic pneumonia coincides with viral lower respiratory tract infections in early childhood, in contrast to nonpneumonia invasive pneumococcal disease, in the pre-pneumococcal conjugate vaccine era. *Clin Infect Dis* 2015; 60: 1384–7.
- 25- Murdoch KM, Mitra B, Lambert S, Erbas B. What is the seasonal distribution of community acquired pneumonia over time? A systematic review. *Aust Emerg Nurs J* 2014; 17(1): 30-42.