



## **Effect of Vitamin D Deficiency on Dental Caries Among School Aged Children of The Local Area of District Mardan**

Muhammad Iqbal Khan Rahman<sup>1</sup>, Usman Saeed<sup>2</sup>, Shakir Ullah<sup>3</sup>, Dr.Umair Islam<sup>4</sup>, Hamza Khan<sup>3</sup>, Said Ullah<sup>3</sup>, Abbas Khan<sup>3</sup>, Basit Ali<sup>3</sup>, Iqbal Muhammad<sup>5</sup>, Noor Muhammad<sup>5</sup>, Kabir Ahmad<sup>3</sup>, Muhammad Yaseen<sup>2</sup>

<sup>1</sup> Department of Microbiology University of Swat, Pakistan

<sup>2</sup> Department of Zoology GC University Lahore, Pakistan

<sup>3</sup> Department of Pathology Takht Bhai Institute of Health and Management Sciences Pakistan

<sup>4</sup> Department of Internal Medicine Khyber Teaching Hospital Peshawar Pakistan

<sup>5</sup> Department of Microbiology Kohat University of Science & Technology (KUST), Kohat, Pakistan

\*Corresponding Author: [shakirullah1992@gmail.com](mailto:shakirullah1992@gmail.com)

+923469058726

### **ABSTRACT**

Dental caries and Vitamin D deficiency are a worldwide phenomenon. Therefore the study aims to explore the effect of vitamin D deficiency on dental caries among school aged children of the local area of district Mardan. A total of 800 children between the ages of 4-15 years were studied. A systematic random sampling technique was applied for sample collection. In the current research work, 800 samples were estimated for the exploration of vitamin D deficiency in school-aged children. A high percentage of Vitamin D deficiency was found in males 90(22.5%) while a very low ratio was found in females 50(12.5%). The ratios of insufficiency in the school-aged children were also high in male 110(27.5%) children as compared to female 50(12.5%) students were respectively. Aged-wise exploration of vitamin D deficiency shows that very low frequency was showed in low aged group 4-7 years 40 (10%), similarly in 8-11 years the frequency of vitamin D deficiency was 60(15%) while in the aged group 12-15 years the frequency of Vitamin D deficiency were found very high 80 (20%). Residency-wise results show that the frequency ratio of urban children was high 80(20%) as compared to rural area children 60(15%) respectively. This emerging health problem of Vitamin D deficiency and its connection with the academic career of school-going children has long-term health impacts. The present study highlighted the association between vitamin D deficiency in different age groups, genders, and socioeconomic status-wise exploration of vitamin D deficiency.

**Keywords:** Vitamin D, Health, Deficiency, Exploration.

*Received: 22-5-2024*

*Accepted: 14-6-2024*

*Published: 1-6-2024*

## INTRODUCTION

Vitamin D deficiency is an important public health problem in both developed and developing countries, with a reported worldwide prevalence of 30-80% in children and adults. The role of vitamin D in bone mineralization is well-documented **Calvo et al 2005**. Vitamin D [25(OH) D] started to gain importance worldwide for its important role in healthy bone structure and calcium and phosphate metabolism. There are many studies showing 25(OH) D deficiency and insufficiency in children worldwide (**Cediel 2018**). In the presence of 25(OH) D deficiency and insufficiency, absorption of both calcium and phosphorus is impaired resulting in reduced bone mineral density (**Adam et al 2010**). Low levels of 25(OH) D affects an individual's present and future health status, triggering multiple systemic responses reducing bone density and the level of immune response since there are 25(OH) D receptors in a wide range of tissues, and are related with retarded growth, skeletal deformities and secondary hyperthyroidism in the childhood, whereas hip fracture in the elderly is observed in individuals with impaired bone structure (**DeLuca et al 2004**). Also, there are increasing data explaining the relationship between low levels of 25(OH) D and different types of non-skeletal diseases including some types of cancer, autoimmune, infectious, cardiovascular and psychiatric diseases (**Holick et al 2007**). Risk factors for 25(OH)D deficiency in children were defined as obesity, intestinal malabsorption syndromes, usage of anticonvulsant agents such as Phenytoin, phenobarbital, and carbamazepine, low levels of sun exposure, clothing habits, climatization and seasonality, nutritional choices, dark skin color (**Autier et al 2010**). In order to determine an individual's vitamin D status, serum 25 (OH) D levels is measured. There are different threshold points used to determine 25 (OH) D statuses of individuals as suggested by different organizations and in guidelines (**Alyahya 2017**). Regular measurement of 25(OH) D levels in the childhood and replace the low levels with vitamin D fortification or supplementation is essential and a public health matter in order to acquire healthy generations with robust bone structure. The aim of our study was to assess serum 25(OH)D levels in elementary school children aged between 6-9 years old within a year duration and determine 25(OH)D status between different seasons(**Mansbach et al 2009**)

### Amis of the study

The aims of the study to explore gender and aged wise deficiency of vitamin D and Dental caries among school aged children of the local area of district Mardan

## MATERIALS AND METHODS

### METHODS

This study was undertaken for the Exploration of Dental caries and vitamin D deficiency in school aged children of the local area of District Mardan. We retrospectively studied the records of a total of 800 children of aged between 4-15 June 2023 to December 2023. All patients were subjected to a careful physical examination. Weights were measured using a calibrated digital scale. Height measurements were done intriplicate to the nearest millimeter using a calibrated stadiometer. Body mass index (BMI) were calculated according to the formula [weight (kg)/height (m)<sup>2</sup>]. Patients with a history of a chronic disorder or on any medication that may alter vitamin D metabolism were excluded from the study. Serum calcium (Ca), phosphorus (P), magnesium (Mg), alkaline phosphatase (ALP) and glucose levels were measured using the enzymatic colorimetric method (Roche Integra 800), while serum 25 hydroxy vitamin D

[25(OH) D] levels were measured by highperformance liquid chromatography (Shimadzu UFLC).

According to WHO formula  $BMI = \frac{\text{weight}}{\text{Height}^2(M^2)}$

s.no	status	Student BMI	Normal BMI	HB level
1	Anemia	20BMI	18.5-24.9	>12,>14g/dL
2	Underweight	18BMI	18.5-24.9	<12,14g/dL
3	Malnutrition	17.3BMI	18.5-24.9	<12,14g/dL
4	Stunting	11.2BMI	18.5-24.9	<12,14g/dL
5	wasting	9.5BMI	18.5-24.9	<12,14g/dL

### Study Area and Period

The study was conducted in Takht Bhai Mardan, which is located Khyber Pakhtunkhwa.

### Study Design

Institutional based cross-sectional study designs were used.

### Study Population

All secondary and primary school students (age group wise the source population, whereas sampled or selected students were the study population of this study.

### Sample Size Determination

A Total of 800 children between the ages of 4-15 years were studied. A systematic random sampling technique was applied for sample collection.

### Lab investigations and Tests: All patients were subjected to:

**Complete Blood count and reticulocyte count.** Patients with microcytosis underwent the following:

#### • Serum Iron and ferritin

An automatic hematological analyzer and Biochemistry analyzer were used for **Clinical Examination**

#### Routine blood tests

This is done to assess anemia and other vitamin and mineral deficiencies. There may be dehydration, low blood sugar and signs of severe infection as is evident by raised while blood cell counts.

#### Diagnosis of malnutrition in children

In children weight and height is measured and compared with the charts showing the expected average height and weight for a child of that age. Some children are persistently smaller for age and may be genetically so.

#### Blood tests in children

Routine blood tests in children include those for blood glucose, blood counts, urine for routine examination.

Levels of iron in blood, folic acid and vitamin B 12 are also done. For protein estimation other tests including

Normal ranges of ferritin 10 to 150 ng/mL for children 4<sup>th</sup> years to 14 years.

MCV normal range is 80 to 95 for children

**ANALYSIS & INTERPRETATION**

► **Analysis and Interpretation of data**

Data will analyze and interpret by using M word, Origin8 and Excel. Frequency and Percentage were calculated for all quantitative variables.

**5. ETHICAL CONSIDERATION**

- The subjects were briefed about the study.
- Consent was taken from the subjects after explaining the purpose of study for the collection of data.
- **Questionnaire form**

<b>Demographics information:</b>	<b>General information</b>	<b>Dietary Habits</b>
Child's Name: _____	Do you have breakfast every day before going to school? (Yes/No)	What is your favorite healthy food?
Age: _____ Gender: _____	How many meals do you typically eat in a day?	How often do you drink water in a day?
Grade/Class: _____	Do you eat fruits and vegetables daily? (Yes/No)	Less than 3 glasses
School Name: _____	How often do you consume fast food? (Yes/No)	3-5 glasses
<b>Family background:</b>	<b>Anthropometric Measurements</b>	6-8 glasses
How many people live in the child's household.?	Height (cm): __	More than 8 glasses
Male and female ratio in child's household.?	Weight (kg): __	Are you aware of the importance of a balanced diet? (Yes/No)
Family income status?	BMI (Body Mass Index): __	Do you receive any nutrition education at school? (Yes/No)
Hereditary diseases in family?		

<b>S.no</b>	<b>Serum 25(OH)D</b>	<b>Status</b>
<b>1</b>	<b>Serum 25(OH)D</b>	<b>Deficient &lt; 50nmol/L</b>
		<b>Insufficient 50-75nmol/L</b>
		<b>Sufficient &lt;75nmol/L</b>
<b>2</b>	<b>Mean Serum 25(OH)D</b>	<b>55± 6nmol/L</b>

**Statistical Analysis**


Statistical analysis was performed by using Origin8 and MS office word 2010.

**Information collection**

## RESULTS AND DISCUSS


### RESULTS

In the current research work 800 samples were estimated for the exploration of vitamin D deficiency in school aged children. Out of 800 samples 400 from male students while 400 were taken from the females' student. The high percentage of Vitamin D deficiency were found in male 90(22.5%) while very low ration were found in female 50(12.5%) Figure.1. while the ratios of insufficiency in the school aged children were also high in male 110(27.5%) children as compare to female 50(12.5%) and the prevalence of Dental caries was found 50(6.25%) students were respectively.




**Figure.1** Gender wise prevalence of Dental caries and Vitamin D deficiency in School Children

Aged wise exploration of vitamin D deficiency shows that very low frequency were showed in low aged group 4-7 years 40 (10%), similarly in 8-11 years the frequency of vitamin D deficiency were 60(15%) while in aged group 12-15 years the frequency of Vitamin D deficiency were found very high 80(20%) shown in Figure.2




**Figure.2** Aged group wise prevalence of Dental caries and Vitamin D deficiency in School Children

Residency wise results shows that the frequency ratio of urban children were high 80(20%) as compare to rural area children 60(15%) respectively figure.3



**Figure.3** Residence wise Vitamin D deficiency in School Children

Our study also reveal that the vitamin D deficiency related with that of socioeconomic status which shows that the high frequency were found in children of high socioeconomic status as compare to that of the children of low socioeconomic status. In conclusion the vitamin D deficiency was found to be dependent on sunlight, diet and the other factors are socioeconomic status, residency of schoolchildren. The results explore that the need for behavior change communiqué on the significance of exposure to sunlight to create appropriate vitamin D in the schoolchildren of the local area of Mardan. This emerging health problem of Vitamin D deficiency and its connection with the academic carrier of school going children long term health consequences. As the present study only highlighted the association between vitamin D deficiency in different aged group, gender and socioeconomic status wise exploration of vitamin D deficiency.



**Figure.4** Socioeconomic status wise Vitamin D deficiency in School Children

## DISCUSSION

In the current research work 800 samples were estimated for the exploration of Dental caries and vitamin D deficiency among school aged children. Out of 800 samples 400 from male students while 400 were taken from the females' student. The high percentage of Vitamin D deficiency were found in male 90(22.5%) while very low ration were found in female 50(12.5%) Figure.1. while the ratios of insufficiency in the school aged children were also high in male 110(27.5%) children as compare to female 50(12.5%) students were respectively. Same study were also conducted by (Zhu et al 2012) a total of 6,008 children aged 1 month to 16 years partaken in this cross-sectional study. All the subjects were divided into subgroups according to their age: 0-1y, 2-5y, 6-11y and 12-16y representative infancy, preschool, school age and adolescence stages respectively. The highest mean level of serum 25(OH)D was found in the 0-1y stage (99 nmol/L) and the lowest one was found in 12-16y stage (52 nmol/L). Accordingly, the prevalence of serum

25(OH)D levels of < 75 nmol/L and < 50 nmol/L were at the lowest among infants (33.6% and 5.4% respectively) and rose to the highest among adolescents (89.6% and 46.4% respectively). The mean levels of serum 25(OH)D and the prevalence of vitamin D deficiency changed according to seasons. In winter and spring, more than 50% of school age children and adolescents had a 25(OH)D level at < 50 nmol/L. If the threshold is changed to < 75 nmol/L, all of the adolescents (100%) had low 25(OH)D levels in winter and 93.7% school age children as well.

Aged wise exploration of vitamin D deficiency shows that very low frequency were showed in low aged group 4-7 years 40 (10%), similarly in 8-11 years the frequency of vitamin D deficiency were 60(15%) while in aged group 12-15 years the frequency of Vitamin D deficiency were found very high 80(20%). Similarly a research work also performed by (Hocaoğlu *et al* 2019) Serum 25(OH)D levels ranged from 3.90 to 64.60 ng/mL, the median value was 25.95 ng/mL for all subjects. Of all the primary school children, 485 (75.78%) had adequate levels of 25(OH)D. Vitamin D deficiency was observed in 36 of children (5.62%), whereas insufficient levels of 25(OH)D were found in 119 children (18.60%). The ratio of vitamin D insufficiency and deficiency together was highest in spring (31.87%) and lowest in summer (13.12%).

Residency wise results shows that the frequency ration of urban children were high 80(20%) as compare to rural area children 60(15%) respectively. Residency wise study also conducted by (Manios *et al* 2017) a sample of 2386 schoolchildren (9–13 years old) from four distinct prefectures was examined. The prevalence of 25-hydroxyvitamin D (25(OH) D) concentration <30 and <50 nmol/l (vitamin D deficiency and insufficiency respectively) was 5.2 and 52.5 %, respectively. Girls had a higher prevalence of 25(OH) D <30 (7.2 v. 3.2 %) and 50 nmol/l (57.0 v. 48.0 %) than boys (P<0.001). The highest prevalence rates of 25(OH) D <30 and 50 nmol/l (9.1 and 73.1 %, respectively) were observed during spring (April to June), whereas the lowest (1.5 and 31.9 %, respectively) during autumn (October to December). The prevalence of 25(OH) D <50 nmol/l was higher in urban/semi-urban than rural regions, particularly during spring months (74.6 v. 47.2 %; P<0.001). Female sex, urban/semi-urban region of residence and spring months were found to increase the likelihood of vitamin D deficiency and insufficiency, with the highest OR observed for spring months (7.47; 95 % CI 3.23, 17.3 and 5.14; 95 % CI 3.84, 6.89 for 25(OH)D <30 and 50 nmol/l respectively).

In the current research work the prevalence of Dental caries was 50(6.25%) in vitamin D deficient students in the local area of district Mardan. A study was also conducted by [32] The dental index of decayed, missing and filled teeth (DMFT) in the deficient and insufficient vitamin D groups was significantly higher than in the sufficient vitamin D group (P=0.03). The linear regression model on the effect of age, vitamin D, and calcium on dental indexes showed no significant statistical relationship. However, in the case of the dental index, age, and calcium level affected dental caries (d), but vitamin D did not affect this index.

## CONCLUSION

The results explore that the need for behavior change communiqué on the significance of exposure to sunlight to create appropriate vitamin D in the schoolchildren of the local area of Mardan. This emerging health problem of Vitamin D deficiency and Dental caries its connection with the academic carrier of school going children long term health significances. As the present



study only highlighted the association between vitamin D deficiency in different aged group, gender and socioeconomic status wise exploration of vitamin D deficiency.

### ACKNOWLEDGEMENT

Authors feel grateful to the Microbiology Department of Takht Bhai Institute of Health and Management Sciences, Microbiology Department of Abasyn University Peshawar and Pathology Department of THQ Takht Bhai Mardan for providing facilities during research activities

### REFERENCES

- Autier P, Boniol M, Pizot C, et al.** Vitamin D status and ill health: a systematic review. *Lancet Diabetes Endocrinol* 2014;2(1):76-89
- Adams JS, Hewison M.** Update in vitamin D. *J Clin Endocrinol Metab* 2010;95:471-478.
- Autier P, Boniol M, Pizot C, Mullie P.** Vitamin D status and ill health: a systematic review. *Lancet Diabetes Endocrinol* 2014;2:76-89. Epub 2013 Dec 6
- Adams JS, Hewison M.** Update in vitamin D. *J Clin Endocrinol Metab* 2010;95:471-478.
- Autier P, Boniol M, Pizot C, Mullie P.** Vitamin D status and ill health: a systematic review. *Lancet Diabetes Endocrinol* 2014;2:76-89. Epub 2013 Dec 6
- Alyahya KO.** Vitamin D levels in schoolchildren: a crosssectional study in Kuwait. *BMC Pediatr* 2017;17:213
- Autier P, Boniol M, Pizot C, Mullie P.** Vitamin D status and ill health: a systematic review. *Lancet Diabetes Endocrinol* 2014;2:76-89. Epub 2013 Dec 6
- Brito A, Cori H, Olivares M, et al.** Less than adequate vitamin D status and intake in Latin America and the Caribbean: a problem of unknown magnitude. *Food Nutr Bull* 2013;34(1):52-64.
- Braegger C, Campoy C, Colomb V, et al.** (2013) Vitamin D in the healthy paediatric population: a position paper by the ESPGHAN Committee on Nutrition. *J Pediatr Gastroenterol Nutr* 56, 692–701.
- Calvo MS, Whiting SJ, Barton CN.** Vitamin D intake: a global perspective of current status. *J Nutr* 2005;135:310-316.
- Cediel G, Pacheco-Acosta J, CastiUo-Durdn C.** Vitamin D deficiency in pediatric clinical practice. *Arch Argent Pediatr* 2018;116:75-81.
- Carlberg C, Molnár F.** Vitamin D receptor signaling and its therapeutic implications: Genome-wide and structural view. *Can J Physiol Pharmacol* 2015;93(5):311-8.
- Cashman KD & Kiely M (2011)** Towards prevention of vitamin D deficiency and beyond: knowledge gaps and research needs in vitamin D nutrition and public health. *Br J Nutr* 106, 1617–1627
- Cediel G, Pacheco-Acosta J, CastiUo-Durdn C.** Vitamin D deficiency in pediatric clinical practice. *Arch Argent Pediatr* 2018;116:75-81
- Durán P, Mangialavori G, Biglieri A, et al.** Estudio descriptivo de la situación nutricional en niños de 6-72 meses de la República Argentina. Resultados de la Encuesta Nacional de Nutrición y Salud (ENNyS). *Arch Argent Pediatr* 2009;107(5):397-404
- DeLuca HF.** Overview of general physiologic features and functions of vitamin D. *Am J Clin Nutr* 2004;80(Suppl 6):1689-1696

- DeLuca HF.** Overview of general physiologic features and functions of vitamin D. *Am J Clin Nutr* 2004;80(Suppl 6):1689-1696
- Elder CJ, Bishop NJ.** Rickets. *Lancet* 2014;383(9929): 1665-76.
- Heaney RP.** Functional indices of vitamin D status and ramifications of vitamin D deficiency. *Am J Clin Nutr* 2004;80(Suppl 6):1706- 1709
- Hilger J, Friedel A, Herr R, et al.** A systematic review of vitamin D status in populations worldwide. *Br J Nutr* 2014;111(1):23-45.
- Holick MF.** Vitamin D Deficiency. *N Engl J Med* 2007;357:266-281.
- Hocaoğlu-Emre, F. S., Sarıbal, D., & Oğuz, O. (2019).** Vitamin D deficiency and insufficiency according to the current criteria for children: vitamin D status of elementary school children in Turkey. *Journal of clinical research in pediatric endocrinology*, 11(2), 181.
- Lindqvist PG, Epstein E, Nielsen K, et al.** Avoidance of sun exposure as a risk factor for major causes of death: a competing risk analysis of the Melanoma in Southern Sweden cohort. *J Intern Med* 2016;280(4):375-87.
- Le Roy C, Reyes M, González JM, et al.** Estado nutricional de vitamina D en pre escolares chilenos de zonas australes. *Rev Med Chil* 2013;141(4):435-41.
- Maestro B, Molero S, Bajo S, et al.** Transcriptional activation of the human insulin receptor gene by 1,25-dihydroxyvitamin D(3). *Cell Biochem Funct* 2002;20(3):227-32.
- Mansbach JM, Ginde AA, Camargo CA Jr.** Serum 25-Hydroxyvitamin D Levels Among US Children Aged 1 to 11 Years: Do Children Need More Vitamin D? *Pediatrics* 2009;124:1404-1410 .
- Manios, Y., Moschonis, G., Hulshof, T., Bourhis, A. S., Hull, G. L., Dowling, K. G., & Cashman, K. D. (2017).** Prevalence of vitamin D deficiency and insufficiency among schoolchildren in Greece: the role of sex, degree of urbanisation and seasonality. *British journal of nutrition*, 118(7), 550-558.
- Misra M, Pacaud D, Petryk A, et al. (2008)** Vitamin D deficiency in children and its management: review of current knowledge and recommendations. *Pediatrics* 122, 398–417.
- Society for Adolescent Health and Medicine.** Recommended vitamin D intake and management of low vitamin D status in adolescents: a position statement of the society for adolescent health and medicine. *J Adolesc Health* 2013;52(6):801-3.
- Taylor CL, Thomas PR, Aloia JF, et al. (2015)** Questions about vitamin D for primary care practice: input from an NIH conference. *Am J Med* 128, 1167–1170.
- Theodoratou E, Tzoulaki I, Zgaga L, et al. (2014)** Vitamin D and multiple health outcomes: umbrella review of systematic reviews and meta-analyses of observational studies and randomised trials. *BMJ* 348, g2035.
- Zhu, Z., Zhan, J., Shao, J., Chen, W., Chen, L., Li, W., ... & Zhao, Z. (2012).** High prevalence of vitamin D deficiency among children aged 1 month to 16 years in Hangzhou, China. *BMC public health*, 12, 1.