



## Cytotoxic Activity of the Green Alga *Caulerpa racemosa* Nanoparticles on Breast Cancer Cells

Sri Haryanti, Mutmainah, Dewi Ramonah, Lia Kusmita\*

STIFAR Yayasan Pharmasi Semarang Letjend Sarwo Edhie Wibobo Km 1, Semarang, Indonesia

\*Corresponding Author: [lia\\_kusmita@yahoo.com](mailto:lia_kusmita@yahoo.com)

### ARTICLE INFO

#### Article History:

Received: March 5, 2024

Accepted: May 24, 2024

Online: June 17, 2024

#### Keywords:

Cytotoxic,  
*Caulerpa racemosa*,  
Nanoparticles,  
Breast cancer

### ABSTRACT

Indonesia possesses abundant marine biological resources, making it an ideal source for medicinal materials and treatments. Among these resources is *Caulerpa racemosa*, a species of green algae commonly known as the sea grape, which harbors a diverse array of secondary metabolites containing bioactive compounds with cytotoxic properties against cancer cells. Cancer remains a formidable global health challenge, particularly impacting women, with factors such as uncontrolled cell division and metastasis contributing to its severity. *Caulerpa racemosa* is one type of green algae that can be utilized as anti-cancer. The caulerpenin content in *Caulerpa racemosa* shows bioactivity against human cell lines and has anticancer, antitumor, and antiproliferation properties. This study aimed to determine the anticancer effect of methanol extract and *Caulerpa racemosa* nanoparticles on MCF-7 breast cancer cells. The formulation of extract nanoparticles utilized the ionic gelation technique. The results showed that the concentrations of nanoparticle preparations used were successively: 15; 30; 60; 120; 240, and 480 $\mu$ g/ mL, with an incubation time for 48 hours. Additionally, the results showed that the methanol extract has a cytotoxic activity, with IC<sub>50</sub> of 38.29 $\pm$  3.2 $\mu$ g /mL in the active category. Moreover, nanoparticle preparations have a cytotoxic activity against MCF-7 cancer cells, with an IC<sub>50</sub> value of 12.35 $\pm$  2.8 $\mu$ g/ mL in the very active category. Based on these results, it appears that *Caulerpa racemosa* nanoparticle preparations have a higher cytotoxic activity than methanol extracts against MCF-7 breast cancer cells.

### INTRODUCTION

Breast cancer is one of the most common cancers affecting women, with an incidence rate of approximately one in 8- 10 women (Fard *et al.*, 2018). Based on World Cancer Report data, it is predicted that there will be an annual increase of 22 million cases (Chia *et al.*, 2015). This disease ranks among the deadliest illnesses for women globally, driven by uncontrollable factors such as cell division and metastasis (Amalina *et al.*, 2021). Presently, the medical approach to cancer treatment involves chemotherapy, radiotherapy, and surgery. However, these methods can induce hypoxia and cell death, potentially harming healthy non-cancerous cells (Zeichner *et al.*, 2015).

Therefore, it is necessary to develop effective anticancer drugs as potential alternatives to chemotherapy. One of them is with natural plant bioactive compounds that increase effectiveness but have mild side effects. Nanotechnology is known as a branch of engineering that deals with the identification and control of materials in the range of 1 to 100nm, thus providing unusual physical, chemical, and biological properties of nanoparticles (Stoica *et al.*, 2013). This technology provides benefits by increasing the bioavailability of active ingredients, controlling the release of active ingredients and possibly improving sensory properties. With a nanometer size, the active ingredient particles are more easily absorbed by the small intestinal wall thus increasing their bioavailability. Absorption of active ingredients is increased due to the increased particle solubility and large particle surface area.

*Caulerpa racemosa* is one type of green algae living in some Indonesian waters. Based on the research of Uddin *et al.* (2020), the cytotoxic activity of *Caulerpa racemosa* has an IC<sub>50</sub> value of 119.62µg/ mL. The content contained in *Caulerpa racemosa* include alkaloids, flavonoids, glycosides, phenols, saponins, steroids, and tannins (Uddin *et al.*, 2020). *Caulerpa racemosa* also contains caulerpenin which shows bioactivity against human cell lines and has anticancer, antitumor, and antiproliferation properties (Chew *et al.*, 2008). Afftan *et al* (2020) reported that the cytotoxicity of ethanol extract of green seaweed *C. racemosa* is less cytotoxic with a value (LC<sub>50</sub> = 929 µg/ml) (Villegas Vilchez *et al.*, 2020). According to Chew *et al* (2008), hexane extract showed the highest cytotoxicity against breast cancer cells, followed by ethyl acetate and ethanol extracts (IC<sub>50</sub> 23.7± 2.0, 66.7± 5.8 and 182.7± 14.3µg/ mL). In this study, the formulation of extract nanoparticles with an ionic gelation technique, using chitosan & tripolyphosphate, was carried out in the hope of improving the physical characteristics of the nanoparticles formed so as to increase the bioavailability of their active compounds.

## MATERIALS AND METHODS

### Equipment and materials

Samples of *Caulerpa racemosa* were collected in July 2023 during bright daylight hours from the Panjang Island, Jepara, Central Java, Indonesia . The materials used in this study were chitosan (pharmaceutical grade), ethanol p.a. (Merck), tripolyphosphate (technical), CH<sub>3</sub>COONa (technical), glacial acetic acid (technical), ethyl acetate (Merck), ethanol 96% (technical), HCl 37% (Merck), NaCl (Sigma-Aldrich), NaOH (Merck), K<sub>2</sub>HPO<sub>4</sub> (Merck) and distilled water. Equipments used in this study included UV-Vis spectrophotometer (Jenway 6800), shaker incubator (Stuart SI500), centrifuge (Boeco Zentrifugen D-78532), analytical balance (Precisa XB 220A), vortex mixer (Stuart SA8), hotplate, and magnetic stirrer (Stuart CB162), pH meter (Jenway 370), micropipette (Smart Gen-nex) and glassware (Pyrex). Formulation was carried out at the Pharmaceutical Technology Laboratory of Stifar Yaphar Semarang. Determination of

particle size and zeta potential were measured using particle size analyzer (PSA) (Beckman Coulter).

### **Extraction of *Caulerpa racemosa***

Fresh *Caulerpa racemosa* was selected for its vibrant green color, it is shaped like seaweed, with small, round, and slightly flattened structures. It was washed using sea water (salt water), followed by rinsing with fresh water until clean, removing dirt that sticks. Then, it was cut into small pieces and pounded. A total of 1kg of fresh *Caulerpa racemosa* was cut into small pieces then pounded and put into a maceration vessel. Subsequently, 5 liters of methanol solvent were added. Then, it was mixed homogeneously while occasionally being stirred. The extract was macerated for a total duration of 3 x 24 hours. The results were filtered using kola cloth. Then, the extract was concentrated with a rotary evaporator with a temperature of 40° C at 50rpm (**Hainil *et al.*, 2022**).

### **Phytochemical screening of *Caulerpa racemosa* extracts**

Phytochemical screening was conducted to determine the class of compounds contained in *Caulerpa racemosa* extract. The screening process was initiated by using color reaction and precipitation. Screening was carried out on alkaloids, flavonoids, saponins, tannins, and terpenoids.

### **Preparation of nanoparticles**

A total of 2.5mL of *Caulerpa racemosa* extract solution of 100mg/ mL concentration variation was then added with 2.5mL of tween 80 (a). The next step was mixing using a magnetic stirrer for 5 minutes at 1200rpm. A solution of 0.3% chitosan in acetate buffer pH 4 was added to a solution of tripolyphosphate (TPP) in distilled water (concentration 00.1%) with a ratio of 1:5 and then mixed using a magnetic stirrer again for 5 minutes at 1200rpm (b). 3mL of mixture a and 3mL of mixture b were mixed and homogenized using a magnetic stirrer for 5 minutes at 1200rpm (**Hussain & Sahudin, 2016**).

### **Evaluation of physical characteristics of nanoparticles**

The particle size, zeta potential and polydispersity index obtained after ultracentrifugation were resuspended in distilled water. The average particle size, zeta potential and PDI were then measured using a particle size analyzer (PSA) (Beckman Coulter) (**Hussain & Sahudin, 2016**).

## Cytotoxic test of *Caulerpa racemose* nanoparticles with MTT method

### a. Cell culture

This study used test cells, namely MCF-7 (breast) cells. MCF-7 cells were grown using Dulbecco's modified eagle medium (DMEM) containing 5% fetal bovine serum (FBS); 1% penicillin-streptomycin; and 0.5% amphotericin A, and then incubated at 37°C in a 5% CO<sub>2</sub> incubator. Cell harvesting was performed by the addition of trypsin-EDTA after cell confluence.

### b. Cytotoxic test


The cytotoxic test of nano preparations of seaweed extracts was carried out by the MTT method as described by Pakki *et al.* (2019) and Tanumihardja *et al.* (2020), with slight modifications (Pakki *et al.*, 2019; Tanumihardja *et al.*, 2020). A total of 100µL of cell suspension (104 cells/ mL) was put into a 96-wellplate and incubated for 24 hours, then 100µL of nano-preparation with various concentrations (6.25- 500µg/ mL) was added. A solution of 3-(4,5-dimethylthiazol-2-yl)-2,5- diphenyltetrazolium bromide (MTT) with a concentration of 0.5mg/ mL was added to each well and incubated for 4 hours. The reaction was stopped by the addition of 10% sodium dodecyl sulfate (SDS). The absorbance of each well was measured using a microplate reader (Thermo) at a lambda of 595nm. Percent inhibition was calculated by the formula:

$$\text{Inhibition (\%)} = \frac{\text{OD of cell control} - \text{OD of sample}}{\text{OD of cell control}} \times 100$$

## RESULTS AND DISCUSSION

The extraction process in this research uses methanol. The choice of methanol solvent was based on the fact that methanol is universal (can attract all compounds). The extract obtained from the extraction process underwent a methanol-free test first. Methanol-free test results were carried out to ensure that the concentrated extract obtained is free from methanol solvents. Moreover, the results of the ethanol-free test are shown in Table (1).

**Table 1.** Methanol free test results of *Caulerpa racemose* extract

Reagent	Positive result (literatur)	Research	Keterangan
+ potassium dichromate	The color of the solution changes		(-) negative
+ H <sub>2</sub> SO <sub>4</sub>		Color does not change	

The test results showed that *Caulerpa racemose* extract was free from methanol solvents since there was no color change after the addition of potassium dichromate and sulfuric acid. After the methanol-free test, the extract obtained was then subjected to the phytochemical screening test. The results of the screening test are shown in Table (2).

**Table 2.** Phytochemical screening of *Caulerpa racemose* extract

Uji	Reagent	Literature (Harbone, 1987)	Research	Conclusion
Flavonoid	Magnesium powder + HCl <sub>(p)</sub> + amyl alcohol	The solution is red, yellow, or orange in the amyl alcohol layer	The solution is red	Positive
Tannin	Gelatin	White precipitate	Brown solution	Negative
Alkaloid	HCl 2N + Dragendorff	Orange precipitate	no precipitate is formed	Negative
	HCl 2N + Mayer	White precipitate	no precipitate is formed	Negative
	HCl 2N + Wagner	Brown precipitate	no precipitate is formed	Negative
Saponin	shaken + HCl <sub>(p)</sub>	No foam	Stable foam	Negative
Terpenoid	Ether + acetat acid + H <sub>2</sub> SO <sub>4(p)</sub>	Green	Blue or green (Steroids), purple (Terpenoids), 2007)	Positive (steroid)

Screening results showed that *Caulerpa racemose* extract was positive for flavonoids and terpenoids. The test was confirmed by KLT test; the results of KLT test of *Caulerpa racemose* extract for terpenoid group are depicted in Fig. (1).



**Fig. 1.** TLC results of confirmatory tests on (a) flavonoids and (b) steroids

The KLT results were positive for flavonoids with yellow stains. After ammonia evaporation, a brownish yellow stain was formed on the extract sample. According to the literature, the sample contains flavonoids if it gives a brownish stain after ammonia evaporation (Harborne, 1987). Likewise, terpenoids showed positive stain results after being sprayed with 10% H<sub>2</sub>SO<sub>4</sub> spot. The presence of steroids is indicated by the appearance of greenish yellow, blue, purplish red, reddish blue, yellowish white, purplish blue spots (Wafa *et al.*, 2014).

The extract obtained was then made into nanoparticle preparations. In this study, several stages were carried out to obtain nanoparticle preparations. The initial stage is the preparation of a formula that refers to the research of Hussain and Sahudin (2016). The use of the ratio between chitosan and tripolyphosphate (TPP) is based on the research of Stoica *et al.*, (2013), chitosan and TPP are used because they can produce different nanoparticle sizes. Via increasing the ratio of TPP chitosan, it will produce nanoparticles with a smaller size. The results of the comparison used in this study, chitosan: TPP, is 1:5. The next stage is homogenization. One of the parameters affecting the particle size is intensity and homogeneity (Gupta, 2006). The results of nanoparticle preparation are shown in Fig. (2).



**Fig. 2.** *Caulerpa racemosa* extract nanoparticle preparation

The initial characteristics of nanoparticle preparations were carried out by physical observation. The results of physical observations in the form of clarity show that the *Caulerpa racemose* extract nanoparticle preparation is clear and has a greenish color. The results of the evaluation of *Caulerpa racemose* extract nanoparticles are further shown in Table (3).

**Table 3.** Evaluation results of *Caulerpa racemose* extract nanoparticles

Nanoparticle size (nm)	Zeta potential (mV)	Polidispersitas Index
83.7 ± 0.65	+29.15 ± 0.55	0.37 ± 0.01


The size of the resulting nanoparticles is still in the range of 10- 1000nm (Stoica *et al.*, 2013). According to Colonna *et al.* (2008), nanoparticles with TPP chitosan are produced using the ion gelation method, resulting in particles with sizes ranging from 200 to 500nm. The zeta potential value of nanoparticles is generally used to characterize the surface charge properties of these nanoparticles. The particle surface characteristics of a

nanoparticle system affect stability. Particles with potential zeta values more positive than +30 mV or more negative than -30 mV are predicted to be stable during storage, with no aggregation occurring between particles (Mohanraj & Chen, 2006). The potential zeta value obtained from the nanoparticle preparation is close to +30 mV, hence it can be predicted that the nanoparticle preparation is quite stable.


The polydispersity index value is a parameter that expresses the particle size distribution of a nanoparticle system. A polydispersity index value of less than 0.3 indicates that the size distribution is very narrow. This can be seen from the small particle size and homogeneous particle size distribution. While, the polydispersity value of more than 0.3 (PI > 0.3) indicates that the particle size distribution is very broad. The polydispersity index value states the stability of a nanoparticle system. A higher polydispersity index value suggests that more particles are aggregated, resulting in the increasing instability of the preparation (Feng Lin Yen *et al.*, 2010).

The extracts and nanoparticle preparations were then subjected to cytotoxic testing against MCF-7 cells. The ability of a substance to kill cancer cells can be measured through cytotoxic test using MTT method. The principle of this method is the change of yellow color from MTT tetrazolium to formazan which is purple. MTT will only be absorbed by living cells and reduced by the reductase enzyme in the mitochondria to formazan salt which is insoluble in water. To dissolve the formazan salt and stop the reaction, 10% SDS is added. The more purple color means the more cells are alive, and vice versa (Rai *et al.*, 2018; Benov, 2019). The results of the cytotoxic test of *Caulerpa racemose* extract nanoparticle preparations against MCF-7 cells are displayed in Fig. (3).

The cytotoxic effect of the *Caulerpa racemosa* extract nanoparticle preparation is known to increase in a dose-dependent manner, with higher concentrations used. The cytotoxic effect began to appear at a concentration of 7.5 µg/ mL and reached a maximum mortality at a concentration of 240.00 µg/ mL (Fig. 3). These results indicate that the nanoparticle preparation of *Caulerpa racemose* extract has a cytotoxic effect on cancer cells. Good cancer cell candidates must exhibit toxicity to cancer cells while remaining safe (non-toxic) to normal cells (Andreani *et al.*, 2017).



(a)



(b)

**Fig. 3.** Cytotoxic effect of (a) methanol extract and (b) *Caulerpa racemose* nanoparticle preparations on MCF-7 cells

The cytotoxic effect of a substance is assessed by the value of the half maximal inhibitory concentration or better known as  $IC_{50}$ .  $IC_{50}$  is the concentration that can kill 50% of cancer cells. The smaller the  $IC_{50}$  value, the greater the cytotoxic activity. The  $IC_{50}$  value of the methanol extract was  $38.29 \pm 3.2 \mu\text{g}/\text{mL}$ , and the nanoparticle preparation was  $12.35 \pm 2.8 \mu\text{g}/\text{mL}$ . Methanol extracts made into nanoparticle preparations have better cellular uptake ability than in the form of extracts. According to



**Nordin et al. (2018)**, the cytotoxic activity of a substance can be categorized based on the  $IC_{50}$  value. An  $IC_{50}$  value of  $\leq 20\mu\text{g/ mL}$  falls into the "very active" category, while a range of 20- 100 $\mu\text{g/ mL}$  is considered "active." A range of 100- 1000 $\mu\text{g/ mL}$  is categorized as "very weak," and an  $IC_{50}$  value of  $\geq 1000\mu\text{g/ mL}$  is classified as "inactive." Based on these results, the methanol extract is included in the active category and the nanoparticle preparation is included in the very active category against MCF-7 cancer cells.

## CONCLUSION

*Caulerpa racemose* algae has secondary metabolite compounds that have cytotoxic activity against MCF-7 cells. The results showed that the methanol extract was included in the active category with an  $IC_{50}$  of  $38.29\pm 3.2\mu\text{g/ mL}$ . For nanoparticle preparations fall in the category of "very active" against MCF-7 cancer cells, with an  $IC_{50}$  value of  $12.35\pm 2.8\mu\text{g/ mL}$ .

## CONFLICT OF INTEREST

The authors declare no conflict of interest.

## ACKNOWLEDGE

The authors would like to thank Yayasan Pharmasi Semarang for the financial support determined for this research through "HIBAH YAYASAN 2023" research grant scheme.

## REFERENCES

- Amalina A.D.; Wahyuni S. and Harjito.** (2021). Cytotoxic effects of the synthesized Citrus aurantium peels extract nanoparticles against MDA-MB-231 breast cancer cells. J. Phys. Conf. Ser., 1918 (3). doi: 10.1088/1742-6596/1918/3/032006.
- Andreani N.A.; Renzi S.; Piovani G.; Ajmone M. P.; Bomba L.; Villa R.; Ferrari M. and Dotti S.** (2017). Potential neoplastic evolution of Vero cells: In vivo and in vitro characterization. Cytotechnology., 69(5); 741-50
- Benov L.** (2019). Effect of growth media on the MTT colorimetric assay in bacteria. PLoS One., 27:14(8). doi: 10.1371/journal.pone.0219713. PMID: 31454355; PMCID: PMC6711527.
- Chew Y. L.; Lim Y.Y.; Omar M. and Khoo K.S.** (2008). Antioxidant activity of three edible seaweeds from two areas in South East Asia. LWT - Food Sci. Technol., 41 (6): 1067–1072. doi: 10.1016/J.LWT.2007.06.013.
- Chia Y.Y.; Kanthimathi M.S.; Khoo K.S.; Rajarajeswaran J.; Cheng H.M. and Yap**

- W.S.** (2015). Antioxidant and cytotoxic activities of three species of tropical seaweeds. *BMC Complement. Altern. Med.*, 15 (1). doi: 10.1186/s12906-015-0867-1.
- Colonna C.; Conti B.; Perugini P.; Pavanetto F.; Modena T.; Dorati R.; Ladarola P. and Genta I.** (2008). Ex vivo evaluation of prolidase loaded chitosan nanoparticles for the enzyme replacement therapy. *Eur. J. Pharm. Biopharm.*, 70(1): 58–65. doi: 10.1016/j.ejpb.2008.04.014.
- Fard S.E.; Tafvizi F. and Torbati M. B.** (2018). Silver nanoparticles biosynthesised using *Centella asiatica* leaf extract: Apoptosis induction in MCF-7 breast cancer cell line. *IET Nanobiotechnology*, 12(7):994–1002. doi: 10.1049/iet-nbt.2018.5069.
- Yen F.L.; Wu T.H.; Tzeng C.W.; Lin L.T. and Lin C.C.** (2010). Curcumin nanoparticles improve the physicochemical properties of curcumin and effectively enhance its antioxidant and antihepatoma activities. *J Agric Food Chem.*, 23:58(12):7376-82. doi: 10.1021/jf100135h. PMID: 20486686.
- Gupta R.B.** (2006). *Nanoparticle Technology for Drug Delivery*. Taylor&Francis Group. New York. 130pp.
- Hainil S.; Sammulia S. F. and Adella A.** (2022). Aktivitas antibakteri *Staphylococcus aureus* dan *Salmonella thypi* ekstrak metanol anggur laut (*Caulerpa racemosa*). *J. Surya Med.*, 7(2): 86–95. doi: 10.33084/jsm.v7i2.3210.
- Harborne J.B.** (1987). *Metode Fitokimia : Penuntun Cara Modern Menganalisis Tumbuhan Edisi Kedua*. ITB. Bandung.
- Hussain Z. and Sahudin S.** (2016). Preparation, characterisation and colloidal stability of chitosan-tripolyphosphate nanoparticles: Optimisation of formulation and process parameters. *Int. J. Pharm. Pharm. Sci.*, 8 (3): 297–308.
- Kammona O. and Kiparissides C.** (2012). Recent advances in nanocarrier-based mucosal delivery of biomolecules. *J. Control Release*, 161(3): 781-794. doi: 10.1016/j.jconrel.2012.05.040.
- Mohanraj V.J. and Chen Y.** (2006). Nanoparticle-A Review. *Tropical Journal Of Pharmaceutical Research.*, 5(1):561-573.
- Nordin M.L.; Abdul K. A.; Zakaria Z.A.; Abdullah R. and Abdullah M.N.H.** (2018). In vitro investigation of cytotoxic and antioxidative activities of *Ardisia crispa* against breast cancer cell lines, MCF-7 and MDA-MB231. *BMC Complement Altern Med.*, 18(1): e87

- Pakki E.; Alam G.; Usmar; Syukur R.; Muslimin L.** (2019). Anticancer activity of selected medicinal plants indigenous to Duri ethnic. *Int J Pharm Sci Res.*, 11(Supplementary 1): 602-08
- Rai Y.; Pathak R.; Kumari N.; Sah D.K.; Pandey S.; Kalra N.; Soni R.; Dwarakanath B.S. and Bhatt A.N.** (2018.) Mitochondrial biogenesis and metabolic hyperactivation limits the application of MTT assay in the estimation of radiation induced growth inhibition. *Scientific Reports.*, 8(1)
- Stoica R.; Şomoghi R. and Ion R.M.** (2013). Preparation of chitosan - Tripolyphosphate nanoparticles for the encapsulation of polyphenols extracted from rose hips. *Dig. J. Nanomater. Biostructures*, 8 (3): 955–963.
- Tanumihardja M.; Hastuti S.; Nugroho J.J.; Trilaksana A.C.; Natsir N.; Rovani C.A. and Muslimin L.** (2020). Viabilities of Odontoblast Cells Following Addition of Haruan Fish in Calcium Hydroxide. *Open Access Maced J Med Sci.* 25;8(D):58-63. DOI: <https://doi.org/10.3889/oamjms.2020.4362>
- Uddin S.A.; Akter S.; Hossen S. and Rahman M.** (2020). Antioxidant, antibacterial and cytotoxic activity of *Caulerpa racemosa* (Forsskål) J. Agardh and *Ulva* (Enteromorpha) intestinalis L. *Bangladesh J. Sci. Ind. Res.*, 55 (4): 237–244. doi: 10.3329/bjsir.v55i4.50959.
- Villegas Vélchez L.F.; Ascencios J.H. and Dooley T.P.** (2022). GlucoMedix®, an extract of *Stevia rebaudiana* and *Uncaria tomentosa*, reduces hyperglycemia, hyperlipidemia, and hypertension in rat models without toxicity: a treatment for metabolic syndrome. *BMC Complement. Med. Ther.*, 22 (1): 1–19. doi: 10.1186/S12906-022-03538-9/FIGURES/8.
- Zeichner S.B.; Ambros T.; Zaravinor J.; Montero A.J.; Mahtani R.L.; Ahn E.R.; Mani A.; Markward N.J. and Vogel C. L.** (2015). Defining the survival benchmark for breast cancer patients with systemic relapse. *Breast Cancer Basic Clin. Res.*, 9: 9–17. doi: 10.4137/BCBCr.s23794.