

Gene Action for Yield and Yield Component Traits in Bread Wheat (*Triticum aestivum* L.) Under late Sown Conditions

Ibrahim, E. M. A.

Department of Genetic, Faculty of Agriculture, South Valley University, Quna, Egypt.



ABSTRACT

The present work was conducted during the two wheat growing seasons of 2013/2014 and 2014/2015 respectively, at the Experimental Farm of South Valley University to study the gene action under normal sowing date (30th November) and late sowing dates (10th January). Five parents were crossed in a half diallel cross mating design. The F₁'s hybrids and their parents were raised under normal and late sowing dates. The analysis of the F₁ generations and their parents revealed that additive gene effects was manifested for all traits studies in the first sowing date, but the reverse was true in the second sowing date. However, over-dominance type of gene action was recorded for grain yield in the first sowing date and partial dominance in the second sowing date. As to the number of kernels/spike partial dominance was displayed in the first sowing date and over dominance in the second sowing date. Narrow sense heritability estimates ranged from 0.49 in the first sowing date to 0.69 in the second sowing date for grain yield. Meanwhile, narrow sense heritability was 0.58 for 1000-grain weight and 0.69 for grain yield in the second sowing date.

INTRODUCTION

Wheat (*Triticum aestivum* L) is the most widely consumed cereal crop worldwide especially in Egypt and is the world's most widely adapted crop, being grown on wide range of environmental condition. Eighty percent wheat crop is sown late, which is ultimately exposed to high temperature combined with lack of water during filling and hence grain yield are reduced substantially (Anonymous, (2007), Hussain *et al.*, (2010) and Mahmood *et al.*, (2010). Heat stress reducing grain quality and grain was reported by (Stone and Nicolas, 1994). Also, heat stress is major yield limiting factor and genetic management is the most apposite solution. Diallel analysis of the genetic traits is a valuable aid in selection and development of high yielding and wheat varieties. Grain yield is highly affecting by environmental stress like drought stress and heat stress because over 50% of the total wheat is sown late which results in terminal heat stress (Kaur and Behi, 2010). Breeder's evaluate many lines during selection for heat tolerance because identification of a plant with all the required genes is difficult (Ortiz-Ferrera *et al.*, 1993). Twelve lines and four testers were crossed in an L X T mating design by (Punia *et al.*, 2011). The 44 crosses and their parents were raised under normal (21 November 2002) and late sown (1, January 2003) seasons at the Experimental Farm of Ragathan College of Agriculture. Their results indicated that the parents like C306, k'sama, K9708, DWR195 and Kailash could be utilized in multiple crossing programs and further biparental mating for selection of high yielding progenies for heat tolerance. Additive gene action was found to be involved in controlling spike length (Chowdhry *et al.*, 2005). Over dominance was reported by (Khan and Dar, 2010) for number of grains per spike and grain yield per plant. Dominance was observed for spike length and grain yield. It showed potential for availability of transgressive segregates in late generation (Sadia *et al.*, 2013). Muhammad *et al.*, (2012a) analyzed F₁ progenies of 7 X 7 diallel fashion crosses comprising

four high temperatures and three susceptible spring wheat parental genotypes were evaluated under normal and heat stress conditions. Analysis of variance under both conditions indicated additive gene action with partial dominance. A full diallel cross study comprising of five bread wheat (*Triticum aestivum* L.) cultivars was carried out with parents and their F₂ progeny to determine gene action for grain yield, plant height and their complements under irrigated and water deficient stress was investigated by (Behnam *et al.*, (2015). Estimates of the genetic components of variation as well as ratio of GCA/SCA showed that all the characters were predominantly controlled by additive gene action. The results of Natasa *et al.*, (2014) indicated significant differences among the parents. However, highly significant differences for general (GCA) and specific (SCA) combining ability in F₁ generation denoted that spike length had resulted from the genes with additive and non-additive. Heritability is one of the determinants for measuring percent improvement to be expected for in a segregating population. Broad sense heritability estimates for days to heading, plant height and 1000-grain weight varied from low to moderate.

Information used in this study on various morphological traits and genetic aspects play an important role for the development of thermo-tolerant varieties and yield stability. Increasing the genetic potential for grain yield of wheat is the main goal of breeding. Yield is a complex trait and is the result of many quantitative traits which are controlled by numerous genes each having small effects. Improving direct and some other indirect components, grain yield can be improved (Zečević *et al.*, (2005). Therefore, the aim of this work is to study the genetic system controlling grain yield, number of kernels, 1000-grain weight and spike length under normal and late sowing dates which represent heat stress in order to elucidate the influence of the environmental on the type of gene action.

MATERIALS AND METHODS

Plant Materials: Five parental genotypes of wheat ie. Giza-168, Sids-112, Shandaweel-1, Qena-25 and Sakha-

94 were used in this study. The seeds were grown in the Experimental Farm of South Valley University.

Field Experiment condition: In 30th November, 2013/2014 wheat growing season the five parental genotypes of wheat were planted into the Field of the Experimental Farm South Valley University Experimental and crossed in a half diallel in order to obtain a total of 10 F₁ crosses. In 2014/2015 the five parents and their 10 F₁ hybrids of the five-parent half diallel cross were sown in the field in the two sowing dates, the first sowing date 30th November, were the sowing condition are favorable and the second sowing date 10th January which allow the plants to be subjected to the heat stress resulting from the rise of temperature late in the growing season. The experimental layout in each date was a complete Randomized Block Design with three replications. The parents were represented by one row of plants per block each row contain 40 plants, while four rows per block were used for each of the 10 F₁ hybrids. Each row was 4.0 meter long, spaced 10 cm apart with plants spaced 20 cm within rows.

Yield and other measurements: Measurements were recorded on a random sample of ten guarded plants for parents and their F₁ hybrids in each replicate in the two experiments. The following characters were recorded for each plant: grain yield, 1000 grain weight, number of kernels/spike and spike length.

Statistical analysis: Means of ten plants for parents and their F₁ hybrids for each of the characters studied were used for statistical analysis (Steel and Torrie, 1980). The collected data were analyzed using diallel analysis as developed by Hayman [1954] and Jinks [1954].

RESULTS AND DISCUSSION

F₁ performance: The data in Table 1 showed that the highly significant among the different entries of F₁ diallel of the four studied traits. Similar results were obtained by Messias and Antonio (2001) and Muhammad *et al.*, (2012b). The means of grain yield/plant ranged from 1.2 (g) for P₅ to 2.73 (g) for P₃ in the first season, while in the second season ranged from 0.45 (g) for P₅ to 1.87 (g) for P₄. Regard to F₁'s the means ranged from 1.41 (g) for (P₃ X P₅) to 3.18 (g) for (P₃ X P₄) in the first sowing date and ranged from 0.39 (g) for (P₃ X P₅) to 2.08 (g) for (P₃ X P₄) in the second sowing date, (Table 2). As to the number of kernels/spike the means of parents ranged from 45 for P₅ to 61.67 for P₃ in the first sowing date, in contrast the means ranged from 38.33 for P₃ and P₅ to 51.33 for P₄. However, the means of F₁'s crosses ranged from 41.67 for (P₂ X P₅) to 59.33 for (P₃ X P₄) in first sewing date, but in the second sowing date the means ranged from 31.67 for (P₂ X P₅) to 46.33 for (P₃ X P₄), (Table 3). Here too, the means of 1000-grain weight ranged from 26.73 (g) for P₅ to 44.44 (g) for P₃ in early sowing date, also the mean of parents ranged from 11.17 (g) for P₅ to 39.95 (g) for P₃ at the late sewing date which represent heat stress. Finally, for spike length the means of parents ranged from 8.67 (cm) for P₅ to 12 (cm) for P₃, but in the second sowing date the means ranged from 5.67 (cm) for P₅ to 8.67 (cm) for P₁ (Table 5). (P₁ X P₃) F₁'s gave the highest mean performance and P₂ X P₄ gave the lowest mean of spike length in the first sowing date. In contrast, the P₃ X P₄ gave the highest mean (10.0), while the (P₁ X P₃) displayed the lowest mean value of spike length.

Table 1. The analysis of variance of four studied traits of 2014/2015 wheat growing season among the different entries of diallel Table.

| Item | sowing dates | d.f | grain yield | 1000 grain weight | number of kernel/spike | spike lenght |
|-----------|-----------------|-----|-------------|-------------------|------------------------|--------------|
| Blocks | 1 st | 2 | 0.0484 | 143.555 | 90.508 | 1.120 |
| | 2 nd | 2 | 0.0539 | 4.9726 | 22.807 | 4.004 |
| Genotypes | 1 st | 14 | 0.7522** | 114.291** | 155.050** | 4.10**8 |
| | 2 nd | 14 | 0.7712** | 109.47** | 317.37**4 | 3.396** |
| Error | 1 st | 28 | 0.0478 | 19.546 | 17.581 | 0.429 |
| | 2 nd | 28 | 0.00705 | 6.4733 | 3.476 | 0.155 |

1st: first sowing date 30th November and 2nd : the second sowing date 10th January

Table 2. The means of grain yield (g) of F₁ diallel cross grown of 2014/2015 wheat growing season.

| Parents | sowing dates | P ₁ | P ₂ | P ₃ | P ₄ | P ₅ |
|------------------------------|-----------------|----------------|----------------|----------------|----------------|----------------|
| P ₁ (Giza-168) | 1 st | 2.00 | 2.08 | 2.49 | 2.21 | 1.75 |
| | 2 nd | 1.22 | 0.95 | 1.53 | 0.86 | 0.49 |
| P ₂ (Sids-112) | 1 st | | 1.43 | 1.60 | 1.95 | 2.05 |
| | 2 nd | | 0.59 | 0.95 | 1.22 | 0.62 |
| P ₃ (Shandweel 1) | 1 st | | | 2.73 | 3.18 | 1.41 |
| | 2 nd | | | 1.53 | 2.08 | 0.39 |
| P ₄ (Qena 25) | 1 st | | | | 2.23 | 2.04 |
| | 2 nd | | | | 1.87 | 1.20 |
| P ₅ (Sakha-94) | 1 st | | | | | 1.20 |
| | 2 nd | | | | | 0.45 |

1st: first sowing date 30th November and 2nd : the second sowing date 10th January

Table 3. the means of Number of Kernels / spike of F1 diallel cross of 2014/2015 wheat growing season.

| Parents | sowing dates | P ₁ | P ₂ | P ₃ | P ₄ | P ₅ |
|-------------------------------|-----------------|----------------|----------------|----------------|----------------|----------------|
| P ₁ (Giza-168) | 1 st | 52.00 | 43.67 | 58.67 | 55.33 | 48.33 |
| | 2 nd | 44.00 | 26.33 | 43.33 | 37.00 | 33.67 |
| P ₂ (Sids-112) | 1 st | | 46.33 | 43.00 | 50.33 | 41.67 |
| | 2 nd | | 37.00 | 36.33 | 40.33 | 31.67 |
| P ₃ (Shandweel -1) | 1 st | | | 61.67 | 59.33 | 47.33 |
| | 2 nd | | | 38.33 | 46.33 | 32.67 |
| P ₄ (Qena-25) | 1 st | | | | 53.00 | 52.33 |
| | 2 nd | | | | 51.33 | 39.00 |
| P ₅ (Sakha-94) | 1 st | | | | | 45.00 |
| | 2 nd | | | | | 38.33 |

1st: first sowing date 30th November and 2nd : the second sowing date 10th January

Table 4. the means of 1000 grain weight of F1 diallel cross of 2014/2015 wheat growing season.

| Parents | sowing dates | P ₁ | P ₂ | P ₃ | P ₄ | P ₅ |
|-------------------------------|-----------------|----------------|----------------|----------------|----------------|----------------|
| P ₁ (Giza-168) | 1 st | 38.53 | 48.47 | 42.88 | 39.78 | 36.18 |
| | 2 nd | 27.78 | 36.16 | 35.21 | 23.12 | 14.50 |
| P ₂ (Sids-112) | 1 st | | 31.11 | 37.26 | 38.61 | 49.62 |
| | 2 nd | | 16.01 | 26.18 | 30.41 | 19.51 |
| P ₃ (Shandweel -1) | 1 st | | | 44.44 | 53.68 | 30.04 |
| | 2 nd | | | 39.95 | 44.89 | 11.82 |
| P ₄ (Qena-25) | 1 st | | | | 42.19 | 39.25 |
| | 2 nd | | | | 36.44 | 30.67 |
| P ₅ (Sakha-94) | 1 st | | | | | 26.73 |
| | 2 nd | | | | | 11.17 |

1st: first sowing date 30th November and 2nd : the second sowing date 10th January

The F₁ diallel analysis: The diallel analysis of variance of the grain yield, number of kernels/spike, 1000-grain weight and spike length per plant for each of the two sowing dates are given in Table 6. The analysis revealed highly significant additive and non additive gene effects over the two sowing dates as indicated by the significance of "a" and "b" items. The additive mean square was greater in magnitude than non additive for the first sowing date for all studied traits, but the reverse was true for the second sowing date for 1000-grain weight. These results are in harmony with results of Muhammad *et al.*, (2012b) who indicated gene action with partial dominance for grain yield and number of kernels/spike suggesting these traits might be useful for the development of terminal heat tolerant Varieties by pedigree selection. Sadia *et al.*, (2013) found that dominance controlling on grain yield and spike length. Also, generation means analysis by Muhammad *et al.*, (2012c) revealed additive, dominance and epistatic genetic effects were operating of the plant character under both conditions. Ahmadi *et al.* (2003) found that the general combining ability (GCA) were highly significant for number of grains/spike, 1000-kernels

weight and grain yield, while Specific combining ability (SCA) effects were highly significant for all studied traits. On partitioning the non-additive effects item "b" into its component, it was evident from the significance of item "b" that F₁ hybrids exhibited dominance for higher grain yield/plant, number of kernels/spike and 1000 grain weight in the first sowing date, but the reverse was true in the second sowing date "heat stress". The average of the F₁ exceeded that of the parents by 7% for grain yield/plant, 12.02% for number of kernels/spike and by 11.98 for 1000-grain weight in the first sowing date, but in the second sowing date the F₁ exceeded that of parents by 0.05 for spike length. The significant "b2" item indicated assymetrical gene distribution of genes affecting grain yield, number of kernels/spike, 1000-grain weight and spike length in the two sowing dates at loci showing dominance, while the significant of "b3" item indicated further dominance effects due to specific combinations. These results were in same line with these obtained by Pumia *et al.*, (2011) who showed that the crosses K9708 X PBN-51 for proline content and heat injury and C309 with kaitash for heat injury had desirable significant SCA effects.

Table 5. the means of spike length of F1 diallel cross of 2014/2015 wheat growing season.

| Parents | sowing dates | P ₁ | P ₂ | P ₃ | P ₄ | P ₅ |
|-------------------------------|-----------------|----------------|----------------|----------------|----------------|----------------|
| P ₁ (Giza-168) | 1 st | 12.67 | 11.00 | 12.50 | 11.50 | 11.33 |
| | 2 nd | 8.67 | 8.67 | 6.50 | 7.83 | 8.17 |
| P ₂ (Sids-112) | 1 st | | 11.00 | 10.33 | 9.33 | 9.33 |
| | 2 nd | | 7.00 | 7.33 | 7.50 | 6.67 |
| P ₃ (Shandweel -1) | 1 st | | | 12.00 | 12.33 | 10.17 |
| | 2 nd | | | 8.17 | 10.00 | 7.17 |
| P ₄ (Qena-25) | 1 st | | | | 11.67 | 10.17 |
| | 2 nd | | | | 8.17 | 8.83 |
| P ₅ (Sakha-94) | 1 st | | | | | 8.67 |
| | 2 nd | | | | | 5.67 |

1st: first sowing date 30th November and 2nd : the second sowing date 10th January

Table 6. The diallel analysis of variance of four studied traits of the F₁ diallel cross grown in the two wheat sowing dates of 2014/2015 season.

| Item | d.f | grain yield | | Number of kernel/spike | | 1000 -grain weight | | spike length | |
|-------------------|-----|------------------------------|------------------------------|------------------------------|------------------------------|------------------------------|------------------------------|------------------------------|------------------------------|
| | | 1 st growing data | 2 nd growing data | 1 st growing data | 2 nd growing data | 1 st growing data | 2 nd growing data | 1 st growing data | 2 nd growing data |
| | | a | 4 | 2.348114** | 3.214518** | 517.4297** | 358.9727** | 179.7305** | 1125.711** |
| b | 10 | 0.8660584** | 0.5652695** | 67.32813** | 119.1391** | 300.2281** | 311.4125** | 2.167969** | 5.57666** |
| b1 | 1 | 0.2845923** | 0.1331413** | 30.72 | 316.2134 | 297.6842** | 11.50134* | 1.92** | 1.3333** |
| b2 | 4 | 0.2168675** | 0.1214447** | 97.52832** | 86.9043** | 96.35547** | 120.7388** | 2.60669** | 2.802765** |
| b3 | 5 | 1.501704** | 1.006755** | 50.4896* | 105.512** | 463.8351** | 523.9337** | 1.866586** | 8.644442** |
| B x a | 8 | 0.0748** | 0.0214** | 31.586 | 11.41 | 22.49 | 7.077 | 0.9567 | 0.16333 |
| B x b | 20 | 0.0296 | 0.00835 | 34.28 | 10.97 | 33.20 | 5.51 | 0.647 | 0.3057 |
| B x b1 | 2 | 0.0341 | 0.00660 | 3.64 | 0.693 | 10.44 | 5.90 | 0.28 | 0.243 |
| B x b2 | 8 | 0.0428 | 0.00254 | 13.59 | 7.86 | 9.62 | 1.69 | 0.547 | 0.169 |
| B x b3 | 10 | 0.024 | 0.0133 | 56.96* | 15.51 | 56.61 | 8.49 | 0.800 | 0.428* |
| Block interaction | 28 | 0.0248 | 0.00705 | 19.5459 | 6.4733 | 17.58 | 3.48 | 0.429 | 0.1545868 |

** : P < 0.01

All items were tested against the block interaction M.S




Fig.1. Wr, Vr graphs of F₁ diallel cross for grain yield in first sowing date (A), for grain yield in second sowing date (B), number of kernels/spike in first sowing date (C) and 1000-grain weight in second sowing date (D)

The analysis of variance of (Wr + Vr) and (Wr - Vr) are shown in Table 7, which revealed highly significant differences for (Wr + Vr) and (Wr - Vr) for grain yield in the two sowing dates indicated the presence of non-additive gene and epistatic effects. While, the analysis of (Wr + Vr) and (Wr - Vr) for

number of kernels/spike revealed non significant for the (Wr + Vr) in the second sowing date and highly significant (Wr - Vr) in the second sowing date. Evidently, additive gene and epistatic effects were presented in the second sowing date. As to the 1000-grain weight the analysis of variance revealed highly

significant of (Wr + Vr) in the second sowing date indicate the presence of non additive gene effects and also highly significant (Wr - Vr) in the two sowing dates. Regards spike length/plant the analysis of (Wr + Vr) and (Wr - Vr) showed significant of (Wr + Vr) in the second sowing date confirming the non-additive gene effect. The slope of the Wr/Vr regression line [Fig. 1 (A and B)] was significantly deviated from zero, but not from unity for grain yield, in the two sowing dates. While, for number of kernels/spike the slope of Wr/Vr was significantly deviated from zero, but not from unity. and in the second sowing date for 1000-grain weight. Evidently additive - dominance model of gene action was operating. The regression coefficient of Wr/Vr was not significantly different from zero for number of kernels/spike in the second sowing date, 1000-grain

weight in the first sowing date indicating that non-allelic gene interaction was manifested. Array No. 3 as common parent represented the extreme recessive genotype which was located at the end of the regression line for grain yield in the two sowing dates, also for number of kernels/spike in the first sowing date and 1000-grain weight in the second sowing date (Fig 1: A, B and C). Meanwhile, array No. 2 represent the extreme dominant gene type which was located near the origin point of regression line for grain yield in the two sowing dates (Fig 1,A and B). For number of kernels/spike array No. 4 represented the extreme dominant genotype which was located near the origin point of regression line. Similar results were also reported by Farooq *et al.*, (2010) and Nazeer *et al.*, (2011).

Table 7. Analyses of variance of Wr + Vr and Wr - Vr values for all studied traits of F₁ hybrids.

| Source of variation | d.f | sowing dates | Grain yield | | Number of kernel/spike | | 1000 grain weight | | Spike length | |
|---------------------|-----|-----------------|-------------|----------|------------------------|----------|-------------------|-----------|--------------|---------|
| | | | Wr+Vr | Wr -Vr | Wr+Vr | Wr-Vr | Wr + Vr | Wr - Vr | Wr + Vr | Wr - Vr |
| Blocks | 2 | 1 st | 0.0126 | 0.0026 | 179.004 | 1605.25 | 3591.73 | 7578.11** | 1.402 | 1.063** |
| | | 2 nd | 0.0205 | 0.00007 | 528.164 | 299.114* | 2411.63 | 93.437 | 0.137 | 0.1313 |
| Array | 4 | 1 st | 0.4094** | 0.0284** | 6357.95* | 950.145 | 4349.63** | 3774.12 | 1.362 | 0.242 |
| | | 2 nd | 0.1765** | 0.0101** | 843.018 | 747.22** | 24790.3** | 885.909** | 2.951* | 1.1036 |
| Error | 8 | 1 st | 0.0136 | 0.0026 | 1871.71 | 508.194 | 1332.59 | 562.462 | 0.476 | 0.177 |
| | | 2 nd | 0.0039 | 0.00032 | 739.145 | 67.152 | 863.31 | 100.500 | 0.514 | 9.3876 |

1st: first sowing date 30th November and 2nd: the second sowing date 10th January

Genetic parameters: The estimates of various components of the genetic variation for grain yield/plant, number of kernels/spike and 1000-grain weight in the first and second sowing dates are given in Table 8. The "D" parameter estimating the additive effect was smaller than "H₁" confirming that over dominance was operating in the first sowing date and partial dominance in the second sowing date for grain yield/plant. As to the number of kernels/plant the "D" parameter estimating the additive effect was much greater than the "H₁" confirming the partial dominance was operating in the first sowing date, but over-dominance was manifested in the second sowing date for 1000-grain weight. The results of Farhad *et al.*, (2011) revealed that significant additive "D" and dominant genetic variations for number of grains/spike, 1000 grain weight and grain yield/plant under early planting. Similarly, under the late planting date significant additive (D) and dominant (H) genetic variation were observed for number of grains/spike and grain yield/plant, which for 1000-grain weight only additive "D" genetic component was significant. Rashid *et al.*, (2012) using 6 × 6 wheat diallel crosses, indicated that grain yield was controlled by additive gene effects. They also emphasized that selection of genotypes in early generations has higher efficiency in breeding programs. The same results were reported for wheat spike length (SL), number of spikelets/spike (SN) and grain yield /plant (GY) (Anwar *et al.*, 2011, Nazir *et al.* 2005, Mahmood and Chowdhry 2002). Also, similar results were obtained by Mahdiyeh and Bahram (2014) who pointed that SCA revealed that selection among progenies of cross ad1 X Darab2 cross would be efficient for increasing grain yield/plant (Gy). The GCA estimates revealed that the cross ad1 and Marvdasht for

grain yield /plant (Gy) were the best combiner. The average degree of dominance measured by (H1/D)^{1/2} ratio reached 1.20 and 0.88 for grain yield in the first and second sowing dates, respectively. As to the number of kernels/spike the (H1/D)^{1/2} reached 0.11 in the first sowing date, but it reached 1.15 for 1000-grain weight in the second sowing date. The "F" value was positive and significant for grain yield in the first and second sowing dates indicating unequal of dominant and recessive alleles among parents. In contrast, The F value was negative in the first sowing date for number of kernels/spike indicating an excess of dominant than recessive alleles. The UV values were less than 0.25 indicating unequal distribution of the dominant and recessive alleles among the five parents which has been indicated before from the significant "b₂". Narrow sense heritability estimates ranged from 0.58 for 1000-grain weight and number of kernel to 0.69 for grain yield in the second sowing date. High estimates of narrow sense heritability for these traits suggest that genetic improvement for heat tolerance (second sowing date) may be achieved through selection breeding. The results of Abd-Allah *et al.*, (2007) indicated that Narrow sense heritability estimates ranged from 0.0% for number of spikes/plant in the third cross to 0.71% for 1000 kernel weight in the first cross under optimum sowing date, and from 31.14% for number of spike/plant in the second cross to 78.21% before maturity date in the first cross under late sowing date. Also Amein, (2007) found that the parent of spring regression heritability, under optimum sowing date ranged from 31.99 for number of spikes/plant in the third cross to 77.19 for grain yield/plant in the first cross and grain yield/plant in the third cross under late sewing date. The results revealed that selection among progenies of the F₁ hybrids in late

generations would be efficient in breeding programs to improve these studied traits.

Table 8. Components of the genetic variation for grain yield, number of kernels/spike and 1000-grain weight of 2014/2015 of wheat growing season.

| Character | Grain yield | | Number of kernels/spike | | 1000-grain weight | |
|-----------------------|------------------------------|------------------------------|------------------------------|------------------------------|------------------------------|------------------------------|
| | X ± S. E | | X ± S. E | | X ± S. E | |
| Item | 1 st growing data | 2 nd growing data | 1 st growing data | 2 nd growing data | 1 st growing data | 2 nd growing data |
| D | 0.36 ± 0.032 | 0.36 ± 0.013 | 24.14 ± 14.26 | 29.23 ± 5.18 | 38.37 ± 14.99 | 153.04 ± 6.34 |
| F | 0.08 ± 0.081 | 0.05 ± 0.028 | -29.25 ± 35.6 | -2.55 ± 12.95 | 30.16 ± 37.47 | 26.40 ± 15.83 |
| H ₁ | 0.56 ± 0.087 | 0.38 ± 0.030 | 13.57 ± 38.50 | 79.98 ± 13.99 | 173.71 ± 40.51 | 222.71 ± 17.12 |
| H ₂ | 0.53 ± 0.079 | 0.36 ± 0.028 | 7.36 ± 34.92 | 66.99 ± 12.69 | 166.39 ± 36.74 | 200.93 ± 15.53 |
| E | 0.02 ± 0.013 | 0.007 ± 0.005 | 19.55 ± 5.82 | 6.47 ± 2.12 | 17.58 ± 6.12 | 3.48 ± 2.59 |
| (H1/D) ^{1/2} | 1.20 | 0.88 | 0.11 | 0.95 | 1.45 | 1.15 |
| UV | 0.24 | 0.24 | 0.14 | 0.21 | 0.24 | 0.23 |
| Broad- sense hb.s | 0.92 | 0.98 | 0.62 | 0.86 | 0.12 | 0.97 |
| Narrow-sense hn.s | 0.49 | 0.69 | 0.58 | 0.49 | 0.74 | 0.58 |

REFERENCES

- Abd-allah M., H. Soheir and L. A. Amin (2013). Genotypic differences for heat tolerance traits in bread wheat using five parameters genetic model Alex. J. Agric. Res., 58 (2): 83-96.
- Ahmadi J., A. Zali, B. Y. Samadi, A. Talaie, M. R. Ghannadha, A. Saiedi (2003). A study of combining ability and gene effect in bread wheat under drought stress conditions by diallel method. Iranian J. Agri. Sci., 34: 1-8.
- Amein K. A. (2007). Genetic improvement of grain quality in some cereal crops, protein quality and quality in wheat (*Triticum aestivum* L.) Ph.D thesis Fac. Of Agric. Assiut Univ Egypt.
- Anonymous, (2007). Recommendations of international wheat seminar. P. 372-274 In: M. A. Khan (Ed) Proc. Int. Wheat Seminar, Wheat Research Institute, Faisalabad, Pakistan, 20 -21 Feb. 2006.
- Anwar J., Akbar M., Hussain M., Asghar S., Ahmad J., and M. Owais (2011). Combining ability estimates for grain yield in wheat. J Agric. Res., 49(4): 437-445.
- Behnam N. K., Saeid A. and S. A. Mohammadi (2015). Genetic analysis of grain yield and plant height in full diallel crosses of bread wheat. Biological forum an International Journal, 7 (1): 1164 -1172.
- Chowdhry M. A., M. S. Saeed, I. Khaliq and M. Ahsan (2005) Combining ability analysis for some polygenic traits in a 5 X 5 diallel cross of bread wheat. Assian J. PI. Sci., 4 (4): 405 – 408.
- Farhad A., K. Saleem, S. Q. Ahmed, K. Hakim, K. Ayub and M. Fida (2011). Genetic analysis of some quantitative traits in bread wheat across environments. African Journal of Agricultural Research, 6(3): 686 – 692.
- Farooq J., I. Khaliq, A. S. Khan and M. A. Pervez (2010). Studying the genetic mechanism of some yield contributing traits in wheat (*Triticum aestivum* L.). Int. J. Agric. Biol. 12: 241–246.
- Hayman, B. I. (1954). The theory and analysis of diallel crosses. *Genetics*, 39: 789–809.
- Jinks J. L. (1954). The analysis of continuous variation in a diallel cross of *Nicotiana rustica* varieties. *Genetics*, 39: 767-788.
- Hussain I., A. W. M. Ashraf and S. M. A. Basra (2010). Changes in growth and yield of maize grown in the glasshouse. Int. J. Agric. Biol., 12: 9-12.
- Kahn M. H. and A. N. Dar (2010). Correlation and path coefficient analysis of some quantitative traits in wheat. Afr. Crop. Sci. J., 18 (1): 9-14.
- Kaur V. and R. K. Behi (2010). Grain yield in wheat as affected by short periods of high temperature, drought and their interaction during pre- and post- anthesis stages. Cereal Res. Commun, 35: 514 – 520.
- Mahdiyeh Z. K. and B. Heider (2014). Diallel cross study for estimating genetic component underling wheat grain yield. J. Biol. Environ. Sci., 8 (22): 37 -51.
- Mahmood N. and M. A. Chowdhry (2002). Ability of bread wheat genotypes to combine for high yield under varying sowing conditions. J. Genet. Breed., 56: 119–25.
- Mahmood S. A., W. F. Javed and S. M. A. Basra (2010). Heat stress effects on forage quality characteristics of maize (*Zea mays* L.) cultivars. Int. J. Agric. Biol., 12: 701-706.
- Messias G. P. and T. A. Antonio (2001). Estimation of genetic components in popcorn based on the nested design. Crop breeding and Applied Biotechnology, 1 (1): 3 -10.
- Muhammad A. R., A. Salam and I. R. Iftikhar (2012b). Genetic Studies for yield and yield related parameters in bread wheat. American-Eurasian J. Agric. & Environ. Sci., 12 (12): 1579-1583.
- Muhammad I., K. A. Ihsan, and A. Asghar (2012a). Genetic studies for some agronomic traits in spring wheat under heat stress. Pak. J. Agri. Sci., 49 (1): 11 – 21.
- Muhammad S., K. A. Iftkar, M. A. Faqir and A. Asghar (2012c). Generation Means Analysis in cotton (*Gossypium hirsutum* L.) for drought tolerance. Pakistan Journal of Nutrition, 11 (10): 843 -847.
- Natasa U., P. Sofiji, D. Miodrag, H. Nikola and M. Vukosavuev (2014). Diallel analysis for spike length in winter wheat. Turkish J. of Agric. and Natural Sci., 2: 1455-1459.

- Nazeer W., J. Farooq, M. Tauseef, S. Ahmed, M. A. Khan, K. Mahmood, A. Hussain, M. Iqbal, and H. M. Nasrullah (2011). Diallel analysis to study the genetic makeup of spike and yield contributing traits in wheat (*Triticum aestivum* L.). Afr. J. Biotechnol., 10(63):13735-13743.
- Nazir S., A. S. Khan, and Z. Ali (2005). Combining ability analysis for yield and yield contributing traits in bread wheat. J. Agric. Social Sci., 1(2): 129-132.
- Ortiz-Ferrera G., R. Rajaram and M. G. Mosaad (1993). Breeding strategies for improving wheat in heat-stressed environments p. 24-32. In D. A. Saunders and G. P. Hettel(ed). Wheat in heat stressed environments: Irrigated, Dry Areas and Rice-wheat farming system. UNDP/ ARC/ BARI/ CIMMYT, Mexico.
- Punia S. S., A. M. Shah and B. R. Ranwaha (2011). Genetic analysis for high temperature tolerance in bread wheat. African Crop Science Journal, 19 (3): 149 – 163.
- Rashid M. A. R., S. Khan A., and R. Iftikhar (2012). Genetic studies for yield and yield related parameters in bread wheat. American-Eurasian J. Agric. Environ. Sci., 12: 1579-1583.
- Sadia K., I. Sobia, K. Ihsan and K. Mahmood (2013). Genetic analysis for yield and some yield related traits in spring wheat (*Triticum aestivum* L.). J. Agric. Res., 51 (3): 272 – 277.
- Steel, R. G. D. and J. H. Torrie (1980). Principles and Procedures of Statistics with Special Reference to Biological Species. 2nd ed. McGraw Hill Book Co., Inc., New York.
- Stone P. J. and M. E. Nicolas (1994). Wheat cultivars vary widely in their responses of grain yield and quality to short periods of post-anthesis heat stress. Australian Journal of plant physiology, 21: 887 – 900.
- Zečević, V. K., D. Mićanović, D. M. Pavlović and D. Urošević, 2005. The inheritance of plant height in winter wheat (*Triticum aestivum* L.). Genetika 37(2):173-179.

فعل الجين لصفات المحصول ومكوناته تحت ظروف الزراعة المتأخرة في قمح الخبز

عبيد محمد أحمد إبراهيم

قسم الوراثة – كلية الزراعة – جامعة جنوب الوادي – مصر

أجريت هذه الدراسة خلال موسمي زراعة القمح 2014/2013 و2015/2014 في تجربة حقلية بمزرعة كلية الزراعة بجامعة جنوب الوادي لدراسة طبيعة فعل الجين لصفات المحصول ومكوناته تحت ظروف الزراعة الطبيعية (30 نوفمبر) والزراعة المتأخرة (10 يناير). واستخدمت في هذه التجربة خمس أصناف من القمح المصري حيث تم زراعة هذه الأصناف في موسم زراعة القمح 2014/2013 وتم إجراء التهجين بين الآباء في اتجاه واحد للحصول على بذور 10 هجن منها في الجيل الأول. وفي موسم 2015/2014 تم زراعة الآباء وهجن الجيل الأول في تجربة قطاعات كاملة العشوائية وتم تحليل بيانات الآباء والجيل الأول لصفات محصول القمح وهي وزن المحصول وعدد الحبوب في كل سنبل ووزن الألف حبة وطول السنبل. أظهرت النتائج أن جميع الصفات محل الدراسة محكومة وراثيا بالطراز المضيف والسيادي لفعل الجين كما كان تأثير فعل الجين المضيف أكثر أهمية لجميع الصفات الوراثية محل الدراسة في الميعاد الطبيعي (30 نوفمبر). وبالنسبة لصفة المحصول ظهر تأثير السيادة الفائقة من فعل الجين في الميعاد الطبيعي بينما ظهر تأثير السيادة الجزئية في الميعاد الثاني. بالإضافة إلى ذلك كانت درجة التوريث بمعناها الضيق والواسع عالية لجميع الصفات محل الدراسة مما يشير إلى إمكانية التربية والتحسين لهذه الصفات من خلال برامج الانتخاب في الأجيال اللاحقة المتقدمة.