

EFFECT OF CULTIVARS AND YEAST EXTRACT ON KEY PRIMARY AND SECONDARY METABOLITES IN *Catharanthus roseus*(L) G. DON.

Mohamed, Z.A.; M.M.M. Darwish and Najme A. Said

Agricultural Botany Department, Faculty of Agriculture, Mansoura University, Mansoura, Egypt



ABSTRACT

A pot experiment was conducted during the growing seasons 2013/2014 and 2014/2015 at The Experimental Farm and laboratories of the Agricultural Botany Department, Faculty of Agriculture, Mansoura University to study the effects of yeast extract either at 0, 4 or 8 gL⁻¹ on key primary and secondary metabolites of *Catharanthus roseus*, cultivars cv. *rosea* and cv. *alba*. Results indicated that total alkaloids, total soluble phenols and flavonoids were significantly higher in *cvrosea* compared with *cvalba*. Application of yeast extract (YE), generally, enhanced the accumulation of alkaloids, phenols and flavonoids. In addition, treatment with YE increased the concentration of kinetin, indole acetic acid, gibberellic acid and benzyladenine whereas decreased that of abscisic acid. Moreover, the concentration of the essential elements N, P, K, Ca and Mg were higher in YE-treated plants, and, the effect of the higher level was more effective. It was concluded that YE could be utilized as an elicitor to enhance the accumulation of the medicinally-important secondary metabolites in *C. roseus* plants.

INTRODUCTION

Catharanthus roseus(L) G. Don is a tropical and subtropical species belonging to Apocyanaceae. *C. roseus* is a renowned medicinal plant due to the presence of alkaloids which is distributed in all parts of the plant. The plant species contains about 130 alkaloids of indole group and many other secondary metabolites including monoterpenoids, glucosides, steroids, phenolics, flavonoids and anthocyanins. The plant is used for the treatment of diabetes, fever, malaria, throat infections, and chest complaints, for the regulation of menstrual cycles and as euphoriant (Ambusta, 1992). The medicinal value of the plant is mostly due to its alkaloids, though this content is considerably low. So, many strategies have been attempted to enhance its alkaloids content. Elicitation have been a widely adopted approach of enhancing secondary metabolite production in general and specifically for inducing the biosynthesis of *C. roseus* alkaloids. Elicitors are either biotic or abiotic. Abiotic elicitors include heavy metal ions, inhibition of some metabolic steps, certain antibiotics, UV radiation, stress factors and growth substances. Treatment with lead was reported to increase content of alkaloids, flavonoids and phenols in *C. roseus* callus cultures (Amirjaniet al., 2015). Tryptophan and Putescine applied as a foliar spray increased total alkaloids content in *C. roseus* leaves (Talaat et al., 2005). In addition, salinity stress (Jaleel et al., 2008, b) and various growth substances (Jaleel et al., 2008, a; Alam et al., 2012) have been used to augment alkaloids biosynthesis. Biotic elicitors are complex biological compounds with unknown composition like microbial cell wall preparations and yeast extract (YE). YE is used as a biotic elicitor for the

induction and enhancement of secondary metabolites (Abraham et al., 2011). YE was reported to enhance silymarin production in cell cultures of *Silybum marianum* (Sanchez-Sampedro et al., 2005; Hasanlooet al., 2008), the alkaloid Mitragnine content in *Mitragynaspeciosa* suspension cultures (Zuldinet al., 2013), Isoflavone content in soybean (Al-Tawaha, 2011), and noradrenaline production in hairy root culture of *Portulacaeoleracea*(Pirian and Piri, 2013). However, the effect of YE on elicitation of secondary metabolites in *C. roseus* *in vivo* is less investigated and poorly understood. So, the aim of the present study was to investigate the effect of YE on key metabolites in *C. roseus* *in vivo*.

MATERIALS AND METHODS

Two pot experiments were conducted during the two successive growing seasons 2013/2014, and 2014/2015 at The Experimental Farm and Laboratories of the Agric. Bot. Dept., Fac. of Agric., Mansoura University, to study the effects of yeast extract (YE) either at 0, 4 or 8 g L⁻¹ on certain secondary metabolites, growth substances and some macro-elements of two cultivars of *Catharanthus roseus* namely cv *rosea* and *cvalba*. *Catharanthus roseus* seeds were collected from the Campus gardens of Mansoura University and were surface sterilized in a 0.2% HgCl₂ solution for 5 min, then thoroughly washed with tap water, and sown on 10th of March and 28 th March in the two growing seasons respectively, in plastic pots, 25 cm in diameter containing 7 kg of a soil, 20 seeds/ pot. Main physical and chemical characteristics of the experimental soil shown in Table (1) were estimated according to Hoddinott and Lamb (1990).

Table (1). Physical and chemical characteristics of the experimental soil.

Sand %	Silt %	Clay	Organic matter %	Total N %	Available K ppm	Available P ppm	TSS
38.6	26.2	35.2	2.1	0.13	226	13	0.24

Pots were irrigated to maintain field capacity and arranged in a complete randomized block design with

four replications. Thirty days after sowing, seedlings were thinned to leave four uniform seedlings per pot. 45

days after sowing (DAS), yeast extract was sprayed onto foliage till leaves dripping using tween 20, 0.05%, as a wetting agent. Control plants were sprayed with deionized water. One week after the first YE application, a second application was done using the same concentrations. Each pot received as calcium superphosphate (15.5 % P₂O₅) at the rate of 3 g/ Pot mixed with the soil before sowing. 2 g ammonium sulphate (20.5 % N) and 2 g of potassium sulfate (50 % K₂O) were also added to each pot in two equal doses, 25, 40 DAS. 65 DAS, shoot samples were collected to determine the following biochemical constituents.

Total alkaloids:

Concentration of total alkaloids was determined according to the method of Afaqet *et al.* (1994). Five hundred mg of shoots powder were transferred to a 100 ml round bottom reflux flask and refluxed for 6 h in a known volume of ethyl alcohol. The extract was then filtered using whatman filter paper No. 1 and 50 ml of dilute HCl was added. Afterward, it was transferred to a separating funnel to which 50 ml of diethyl ether was added. The mixture was shaken for 15 min, the upper diethyl ether layer was discarded whereas the lower water layer was decanted in a beaker and made slightly basic using ammonia solution. The decanted content was again fractionated in a separating funnel using 50 ml of diethyl ether. To the second decant, anhydrous sodium carbonate was added. The mixture was decanted in a pre-weighed dry porcelain dish and was heat-evaporated until dry and weighed again. Total alkaloids concentration was expressed as mg g⁻¹ dry matter.

Total flavonoids:

Total flavonoids concentration was determined by aluminium chloride colorimetric method as described by Lin and Tang (2007). One gram of the methanolic extract was mixed with 0.1 ml of 10% aluminium chloride hexahydrate, 0.1 ml of 1 M potassium acetate and 2.8 ml of deionized water. After incubation for 40 min at room temperature, absorbance of the reaction mixture was recorded at 415 nm. Total flavonoids concentration was estimated from a standard curve established using quercetin and expressed as mg (quercetin equivalent) per g dry matter.

Total Soluble phenols:

Total Soluble phenols concentration was determined using Folin-Ciocalteu reagent according to Singleton *et al.* (1999). The reaction mixture contained 1 ml of the methanolic extract, 9 ml of distilled water, 1 ml of Folin-Ciocalteu reagent and 10 ml of 7% (w/v) sodium carbonate, and incubated for 90 min at room temperature. Absorbance was recorded at 765 nm, and total phenols concentration was estimated using a standard curve established with Gallic acid and expressed as mg (Gallic acid equivalent) per gram dry matter.

Endogenous plant hormones:

Extraction of Endogenous plant hormones were carried out according to the method of Shindy and Smith (1975) and determined using HPLC procedures as described by Baydar and Ulger (1998). For extraction, 6 g of the fresh shoot samples were homogenized and extracted in cold methanol (80% v/v).

The extract was evaporated to the aqueous phase in a rotary evaporator. The aqueous phase was adjusted to pH 8.6 with 1% NaOH and partitioned three times with equal volumes of ethyl acetate. After removal of the ethyl acetate phase, the aqueous phase was adjusted to pH 2.8 with 1% HCl and partitioned three times with equal volumes of ethyl acetate. The combined acidic ethyl acetate phase was used for HPLC determination of acidic endogenous plant hormones. The dried basic ethyl acetate fraction was dissolved in 80% methanol which is then evaporated under vacuum, leaving an aqueous phase which was adjusted to pH 2.8 with 1% HCl and partitioned three times with 50 ml of ethyl acetate. The ethyl acetate phases were combined, reduced to 5 ml volume and used for the determination of neutral auxins. The remaining aqueous phase was adjusted to pH 5.5 with 1% NaOH and partitioned three times with 50 ml of water-saturated L-butanol. Butanol phases were combined, reduced to 5 ml volume and used for the determination of cytokinins.

For HPLC determination of plant growth substances, 20 µl of sample was injected into HPLC (Waters U6K HPLC). Separation and determination were performed on a C₁₈ column (3.9 x 300 mm, silica-based packing material). The elution system consisted of 100% methanol, 2% acetic acid and was run at a flow rate of 1.0 ml min⁻¹.

Determination of elements concentration:

200 mg of the dried shoot powder was wet digested in a mixture of sulfuric and perchloric acids (2:1 v/v). Total nitrogen was determined using modified Microkjeldahl's method according to Pregl (1945). Phosphorus was determined by the molybdenum blue method according to Murphy and Riley (1962). Potassium concentration was determined by the flame photometric method, whereas Ca and Mg concentration were determined using versenate method in the wet digested plant material according to Richards (1954).

Statistical analysis:

ANOVA was performed using SPSS (version 16.0) as a combined analysis of the two growing seasons. Duncan's Multiple Range list was applied to determine significant difference between means when ANOVA was significant at P ≤ 0.05.

RESULTS

Alkaloids, Total soluble phenols and flavonoids concentrations were significantly higher in cv rosea compared with cv alba. Rosea CV contained 46.5, 54.2, 34.5% higher alkaloids, total soluble phenols and flavonoids, respectively compared with cv alba. YE increased alkaloids, total soluble phenols and flavonoids concentration, though the increase was not significant in the case of alkaloids and total soluble phenols (Table 2). In its enhancing effect on the concentration of flavonoids, there was no significant difference between the two tested levels of YE. The interaction between cultivars and YE was not significant regarding the

concentration of alkaloids, Total soluble phenols and flavonoids.

The concentration from endogenous plant hormones differed significantly between the two cultivars. Roseacv contained higher concentration from IAA, GA₃, kinetin (kin) and Benzyl Adenine (BA). On the other hand, cvrosea had lower concentration from abscisic acid (ABA). YE at both applied levels increased Kin whereas decreased abscisic acid concentration (Table 3). On the other hand, IAA, GA₃ and BA concentration were increased only at the higher adopted level. Though concentration of IAA, GA₃, Kin, BA were higher at the higher YE level (8gL⁻¹) compared with those at the lower level (4 gL⁻¹), there was no significant difference between the two levels. This is also true in the case of ABA, though its concentration was lower at the higher level. The interaction between cultivars and YE was significant in case of Kin and BA concentration. The highest Kin concentration was recorded in cvrosea treated with YE at the higher level, whereas the lowest concentration was recorded in cv

albauntreated with YE. Similar trend was recorded regarding BA concentration, where cvrosea treated with YE at its higher level contained the highest BA concentration whereas cv alba not treated with YE contained the lowest level.

Nitrogen (N), phosphorus (P), potassium (K), calcium (Ca), magnesium (Mg) concentration were higher in cv rosea compared with cv alba, though the effect was insignificant in case of nitrogen concentration. YE increased N, P, K, Ca and Mg concentration though the effect was insignificant at the lower level (Table 4). YE at the higher level led to a significant higher concentration from both N, P compared with the lower level. The difference between the two levels regarding K, Ca and Mg was insignificant. The effect of the interaction between cultivars and YE was insignificant regarding the concentration of all elements except Mg. Cultivar rosea treated with the higher level of YE contained the highest Mg concentration whereas the lowest Mg concentration was recorded in cv alba not treated with YE.

Table (2). Effects of yeast extract on total alkaloids, phenols and flavonoids in *C. roseus* shoots.

Character	Treatment	Alkaloids mg g ⁻¹ D. wt.	Phenols mg g ⁻¹ D. wt.	Flavonoids mg g ⁻¹ D. wt
A	A ₁	2.73	14.82	81.10
	A ₂	4.00	22.85	109.08
Significance at 0.05		*	*	*
B	B ₀	3.03	18.25	80.66 ^b
	B ₁	3.31	18.35	97.20 ^b
	B ₂	3.76	19.90	107.41 ^a
Significance at 0.05		N.S.	N.S.	*
A x B	A ₁ B ₀	2.48	14.75	70.75
	A ₁ B ₁	2.78	14.55	82.25
	A ₁ B ₂	2.95	15.15	90.30
	A ₂ B ₀	3.58	21.75	90.85
	A ₂ B ₁	3.85	22.15	112.15
	A ₂ B ₂	4.58	24.65	124.53
Significance at 0.05		N.S.	N.S.	N.S.

* Means followed by the same letter are not significantly different at P = 0.05; NS = not significant

Table (3). Effects of yeast extract on plant growth substances in *C. roseus* shoots.

Treatment	Character	IAA µg 100 g ⁻¹ F.wt	GA ₃ µg 100 g ⁻¹ F.wt	Kinetin	BA	ABS µg 100 g ⁻¹ F.wt
A	A ₁	13.92	18.68	11.15	0.41	7.63
	A ₂	21.57	27.09	16.46	0.70	4.53
Significance at 0.05		*	*	*	*	*
B	B ₀	14.39 ^b	20.25 ^b	9.25 ^b	0.42 ^b	7.76 ^a
	B ₁	17.45 ^{ab}	22.96 ^{ab}	14.58 ^a	0.56 ^{ab}	5.55 ^b
	B ₂	21.39 ^a	23.44 ^a	17.59 ^a	0.69 ^a	4.91 ^b
Significance at 0.05		*	*	*	*	*
A x B	A ₁ B ₀	10.15	16.15	8.75 ^c	0.35 ^c	9.78
	A ₁ B ₁	13.65	18.85	11.23 ^c	0.42 ^c	6.95
	A ₁ B ₂	17.95	21.03	13.48 ^{bc}	0.48 ^c	6.15
	A ₂ B ₀	18.63	24.35	9.75 ^c	0.49 ^c	5.75
	A ₂ B ₁	21.25	27.08	17.93 ^{ab}	0.71 ^b	4.15
	A ₂ B ₂	24.83	29.85	21.70 ^a	0.90 ^a	3.68
Significance at 0.05		N.S.	N.S.	*	*	N.S.

* Means followed by the same letter are not significantly different at P = 0.05; NS = not significant

Table (4). Effects of yeast extract on macroelements in *C. roseus* shoots.

Treatment	Character	Nitrogen mg g ⁻¹ D. wt.	Phosphorus mg g ⁻¹ D. wt.	Potassium mg g ⁻¹ D. wt.	Calcium mg g ⁻¹ D. wt.	Magnesium mg g ⁻¹ D. wt.
A	A ₁	30.80	3.54	30.39	2.74	0.51
	A ₂	33.68	5.04	35.96	3.25	0.70
Significance at 0.05		N.S.	*	*	*	*
B	B ₀	26.24 ^b	3.21 ^b	29.51 ^b	2.30 ^b	0.47 ^b
	B ₁	29.20 ^b	4.23 ^b	33.46 ^{ab}	3.03 ^{ab}	0.55 ^{ab}
	B ₂	41.38 ^a	5.44 ^a	36.55 ^a	3.68 ^a	0.78 ^a
Significance at 0.05		*	*	*	*	*
A x B	A ₁ B ₀	24.80	2.78	26.33	2.08	0.42 ^b
	A ₂ B ₁	28.33	3.33	30.63	2.88	0.51 ^b
	A ₁ B ₂	39.28	4.53	34.33	3.28	0.59 ^b
	A ₂ B ₀	27.48	3.65	32.80	2.53	0.53 ^b
	A ₂ B ₁	30.08	5.13	36.30	3.18	0.60 ^b
	A ₂ B ₂	43.48	6.35	38.78	4.08	0.98 ^a
Significance at 0.05		N.S.	N.S.	N.S.	N.S.	*

* Means followed by the same letter are not significantly different at P = 0.05; NS = not significant

DISCUSSION

The higher alkaloids concentration in *cv rosea* compared with *cv alba* may be attributed to higher activities of carbonic anhydrase and nitrate reductase as well as higher leaf nitrogen concentration in *cv rosea* (Idreeset al., 2010). Dutta et al. (2005) pointed out that cultivar variations in *C. roseus* on the basis of alkaloids concentration is regulated at the level of gene expression.

The use of elicitors is an effective approach for inducing the production of alkaloids in plant tissues (Gautomet al., 2011). Elicitors may be biotic or abiotic. Various biotic elicitors have been employed to enhance alkaloids biosynthesis in *C. roseus*. Within this class of elicitors cell wall filtrates of the fungus *Protomycesgravidus* (Bhagwath and Hjortso(2000), *Pseudomonas fluorescens* (Jaleel et al., 2009), a combination of *P. fluorescens* and *Azospirillumbrasilense* (Karthikeyan et al., 2009), arbuscularmycorrhizal fungi, AMF (Karthikeyan et al., 2009) and cell wall fragments of *Aspergillusniger*, *Fusariummoniliforme* and *Trichoderma viride* (Namedoet al., 2002).

Low yield of *C. roseus* alkaloids versus their high demand worldwide led researchers to try diverse approaches to increase their production. Among these approaches utilization of endophytes (Koulet al., 2013), exogenous application of tryptophan and putrescine (Talaat et al., 2005), induction of stress through salinization (Jaleel et al., 2008b), application of plant growth regulators and fungicides (Jaleel et al., 2008a), application of diverse growth regulators (Alamet al., 2012), and induction of stress through application of heavy metals (Amirjaniet al., 2015).

Application of YE enhanced secondary metabolites production in *C. roseus* (Table 2). Similar results were reported in different plant species (Sanchez-Sampedro et al., 2005; Hasanloo et al., 2008; Al-Tawaha, 2011; Zuldinet al., 2013; Pirian and Piri, 2013). In *Curcuma mangga* cultures, YE induced the accumulation of secondary metabolites (Abrahim et al., 2011). YE enhanced the accumulation of the alkaloid 6-

methoxymellein in carrot cells (Guo and Ohta, 1994). In addition, YE increased alkaloids yield in *Hyoscyamusmuticus* callus (Ibrahim et al., 2009). In *Mitragynaspeciosa*, YE inoculation to cell suspension cultures increased the production of the alkaloid mitragynine and the highest content was achieved at 250 mg l⁻¹ YE (Zuldinet al., 2013). Likewise, YE increased hoscycamine 2.5-fold when added to *Daturastramonium* cultures (Zabetakis et al., 1999).

Simone (2010) reported that the elicitation of terpeneindole alkaloids (TIA) through the application of YE was accompanied with the induction of reactive oxygen species (ROS). Also, the author ascribed YE-induced TIA to enhanced expression of TIA-biosynthetic genes. The generation of ROS through oxidative burst was also reported in tobacco cultures elicited with YE (Baier et al., 1999). A transient increase in cytosolic calcium levels in *C. roseus* cells was reported by Memelink et al. (2001), in harmony with the results of the present investigation (Table 4). Induction of Ca²⁺ is necessary for the induction of Jasmonate accumulation as well as for strictosidine synthase (STR) and tryptophan decarboxylase (TDC) gene expression (Memelink et al., 2001), both effects lead to enhanced TIA biosynthesis. Methyl Jasmonate (MJ) is a general secondary metabolism inducer and enhanced tabersonine biosynthesis in hairy root cultures of *C. roseus* (Rodriguez et al., 2003). In addition, the perception of YE in *C. roseus* suspension cultures leads to the induction of TIA biosynthesis genes including those encoding for STR and TDC (Pauwet al., 2004).

It is worth mentioning that the involvement of the elicitor in the course of essential events in secondary metabolism proceeds as follows (Siddiqui et al., 2013 as follows Siddiqui et al 2013):

- 1) The elicitor binds to plasma membrane receptors.
- 2) Influx of Ca²⁺ to the cytoplasm.
- 3) Cytoplasm pH decreases whereas protein phosphorylation patterns increases and the activity of NADPH oxidases and protein kinase is increased.
- 4) Cell wall structure changes towards enhanced lignification and ROS generation enhances.

- 5) Synthesis of JA and salicylic acid as secondary messengers is enhanced.
- 6) Genes that produce defence-related proteins, plant defense molecules like phytoalexins and other secondary compounds including alkaloids are enhanced.

REFERENCES

- Abraham, F.; A. Bhatt; C.L. Keng; G. Indrayanto and S.F. Sulaiman (2011). Effect of yeast extract and chitosan on shoot proliferation, morphology and antioxidant activity of *Curcuma mangga* *in vitro* Plantlets. African J. Biotech., 10: 7787-7795.
- Afaq, S.H. and Tajuddin Siddiqui, M.M. (1994). Standardization of herbal drugs. Aligarh. Muslim University Publ., Aligarh, India.
- Alam, M.M.;Naeem, M; Idrees, M.; Masroor, M. and Khan, A.(2012). Augmentation of photosynthesis, crop productivity, enzyme Activities and alkaloid production in Sadabahar(*Catharanthus roseus* L.) through Application of Diverse Plant Growth Regulators. J. Crop Sci. Biotech.,15 (2) :117 – 129.
- Al-Tawaha, A.M.(2011). Effect of soil type and exogenous application of yeast extract on soybean seed is flavone concentration.Int. J. Agric. Biol., 3: 275-278.
- Ambusta, C.S.(1992).The Wealth of India. Raw Materials (Revised Edition), Publication and Information Directorate, CSIR, New Delhi; 3: 117.
- Amirjani, M.R.; Abnosi, M.H.; Mahdiyeh, M. and Gharehsheykhlou, S. (2015). Study of protein profile and induction of alkaloids, flavonoids and enzymatic antioxidants in callus of *Catharanthus roseus* L. treated with lead. Science Road J., 3: 12-21.
- Baier, R., Schiene, K., Kohring, B., Flaschel, E., Niehaus, K. (1999). Alfalfa and tobacco cells react differently to chitin ligosaccharides and *Sinorhizobiummeliloti* nodulation factors. Planta, 210: 157-164.
- Baydar H, Ulger S (1998) Correlations between changes in the amount of endogenous phytohormones and flowering in the safflower (*Carthamus tinctorius*L.). Turk. J. Biol. 22: 421-425.
- Bhagwath S.G. and Hjortsø, M.A. (2000). Statistical analysis of elicitation strategies for thiarubrine A production in hairy root cultures of *Ambrosia artemisiifolia*. Journal of Biotechnology,80: 159–167.
- Jaleel, C.A.;Ragupathi Gopi,Rajaram Panneerselvam.(2009). Alterations in non-enzymatic antioxidant components of *Catharanthus roseus* exposed to paclobutrazol, gibberellic acid and *Pseudomonas fluorescens*. Plant Omics Journal, 2:30-40.
- Dutta, A.; Batra, J.; Pandey-Rai, S. *et al.* (2005). Expression of terpenoid indole alkaloid biosynthetic pathway genes corresponds to accumulation of related alkaloids in *Catharanthus roseus*(L.) G. Doss. Planta, 220: 376-383.
- Gautam, S .; Mishra, and Tiwari, A.(2011). Catharanthus alkaloids and their enhanced production using elicitors a review. Int. J. pharm.Techn., 3:371-724.
- Guo, Z. and Y. Ohta (1994). Effect of ethylene biosynthesis on the accumulation of 6-methoxymellein induced by elicitors in carrot cells. J. Plant Physiol., 144: 700-704.
- Hasanloo, T.; Rohnama,;Sepehrifor, R. and Shams, M.R. (2008). The Influence of yeast extract on the production of flavonolignans in hairy root cultures of *Sealyham marionum* (L.) Gaertn. In Abu Osman, H.S. (Ed.) : Biomed 2008 proceedings z1, pp 358-361. Springer- Verlag, Berlin, Heidelberg.
- Hoddinott, K.B. and R.O. Lamb (1990). Physico-Chemical Aspects of Soil and Related Materials. ASTM (ASTM special technical publication;1095) Philadelphia, PA.
- Ibrahim A.I., Abd El Kawi M, Nower A, Abdel Motaal A, Abd El Aal A (2009) Alkaloid Production and Organogenesis from Callus of *Hyoscyamus muticus* L. In vitro Journal of Applied Sciences Research 5,82–92.
- Idrees, M., Naeem, M. and Khon, M.A.(2010). The superiority of cv roseus over cv alba of periwinkle (*Catharanthus roseus* L.) in alkaloid production and other physiological attributes.Turk J.Biol.,43:81-88.
- Jaleel, C.A.; Gopi, R.; Chang-Xirg, Z.; Azooz, M.M. and pannier Selvam, R.(2008,a) plant growth regulators and fungicides alters Growth characteristics in *Catharanthus roseus*; comparative study Global J. Mol. Sic., 3:93-99.
- Jaleel, C.A.; Sankar, B.; Sridharan, R. and PanneerSelvam, R. (2008,b). Soil salinity alters growth, chlorophyll content and Secondary metabolite accumulation in *Catharanthus roseus*. Turk. J. Biol., 32:79-83.
- Karthiketan, B., C. Abdul Jaleel, Zhao Changxing, M.M. Jeo, J. Srimannarayan and M. Deiveekasundaram, 2008. AM fungi and phosphorus levels enhances the biomass yield and ajmalicine production in *Catharanthus roseus*. Eur Asian Journal of Biosciences, 2: 26-33.
- Karthikeyan, Cheruth Abdul Jaleel and M.M. Azooz (2009). Individual and Combined Effects of *Azospirillum brasilense* and *pseudomonas fluorescens* on Biomass Yield and Ajmalicine Production in *Catharanthus roseus*. Academic Journal of Plant Sciences, 2:69-73.
- Koul, M.; Lakra, N.S.; Chandra, R. and Chandra, S.(2013). *Catharanthus roseus* and prospects of its endophytes: a new avenue for Production of bioactive metabolites. Inten. J. pharm. Sci. Res., 4:2705-2716.
- Lin, J.Y. and Tang, C.Y. (2007). Determination of total phenolic and flavonoid contents in selected fruits and vegetables, as well as their stimulatory effects on mouse splenocyte proliferation. Food Chem., 101: 140.

- Memelink J, Verpoorte R, Kijne W (2001). Organization of Jasmonate responsive gene expression in alkaloid metabolism. Trends Plant Sci., 6: 212-219
- Murphy, J. and Riley, J.P. (1962). A modified single solution method for the determination of phosphate in natural waters. Anal. Chem. Acta, 27: 31-36.
- Namdeo. A.G.(2007). Plant cell elicitation for production of secondary metabolites: A review. Pharmacognosy Reviews 1:69-79.
- Namdeo A, Patil S, Fulzele DP. (2002). Influence of fungal elicitors on production of ajmalicine by cell cultures of *Catharanthus roseus*. Biotechnology Progress; 18:159-162.
- Pauw B, Hilliou FAO, Sandonis Martin V, Chatel G, de Wolf CJF, Champion A, Pre´ M, van Duijn B, Kijne JW, van der Fits L, Memelink J (2004). Zinc finger proteins act as transcriptional repressors of alkaloid biosynthesis genes in *Catharanthus roseus*. J Biol Chem 279:52940-52948
- Pirian, K. and Piri, K. (2013). Influence of yeast extract as a biotic elicitor on noradrenaline production in hairy root culture of *Portulaca oleracea* L. Inter. J. Agron. Plant Prod., 4: 2960-2964.
- Pregl, F. (1945). Quantitative organic microanalysis. 4th Ed., J. and A. Churchill Ltd., London.
- Namdeo A, Patil S, Fulzele DP. Influence of fungal elicitors on production of ajmalicine by cell cultures of *Catharanthus roseus*. Biotechnology Progress 2002; 18:159-162.
- Richards, L.A. (1954). Diagnosis and improvement of saline and alkali Soils. U.S. Dept. of Agric.; Handbook No.60.
- Rodrigues S, Compagnon V, Crouch NP, St-Pierre B, De Luca V (2003). Jasmonate – induced epoxidation of tabersonine by cytochrome P-450 in hairy root cultures of *Catharanthus roseus* Phytchemistry, 46: 401-409.
- Sanchez-Sampedro ,M.A.; Fernandez-Tarrago, J. and Corchet, P. (2005). Yeast extract and methyl jasmonate induced silymarin production in cell cultures of *Silybum marianum* (L.) Gaertn. J. Biotech., 119:60-69.
- Shindy, W.W. and Smith, O.E. (1975). Identification of plant hormones from cotton ovules. Plant Physiol., 55: 550-554.
- Siddiqui, Z.H.; Mujib, A.; Zafar, M.; Aslam, J.; Hakeem, K.R. and Parween, T. (2013). In- vitro production of secondary Metabolites using elicitor in *Catharanthus roseus*: a case study. In K.R. Hakeem et al. (eds). Crop Improvement. Springer Science and Business Media.
- Simone, F. (2010). Biological elicitors of plant secondary metabolites: mode of action and use in the production of nutraceuticals. Bio-Farms for nutraceuticals: functional food and safety control by biosensors, Chapter 12.
- Singleton, V.L., Orthofer, R. and Lamuela-Raventos, R.M. (1999). Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin - Ciocalteu reagent. Meth. Enzymol., 299:152.
- Talaat, I. M.; Bekheta, MA. And Mahgoub, M.H. (2005). Physiological response of periwinkle plants (*Catharanthus roseus* L.) to tryptophan and putrescine. Int. J. Agric. Biol., 7:210-213.
- Zabetakis, I.; Edwards, R. and Hagan, D. (1999). Elicitation of tropane alkaloid biosynthesis in transformed root cultures of *Datura stramonium* Phytochemistry, 50: 53-56.
- Zuldin, N.N.M.; Said, I.M.; Noor, N.M.; Zainal, Z.; Kial, C.J. and Ismail, I. (2013). Induction and analysis the alkaloid Mitragynine content of a *Mitragyna speciosa* suspension culture system upon elicitation and precursor feeding. The Sci. world J., 2013:1-11. (Article ID 209434).

تأثير اصناف ومستخلص الخميرة على بعض مركبات التمثيل الحيوى الأولية و الثانوية الهامة فى نبات الونكا زين العابدين عبد الحميد محمد، محمود محمد مصطفى درويش و نجمة عبد السلام سعيد السائح قسم النبات الزراعى، كلية الزراعة، جامعة المنصورة، المنصورة، ج.م.ع

اجريت التجربة خلال الموسمين ٢٠١٣/٢٠١٤ و ٢٠١٤/٢٠١٥ في المزرعة التجريبية ومعامل قسم النبات الزراعى، كلية الزراعة جامعة المنصورة لدراسة تأثير مستخلص الخميرة بتركيز ٤ و ٨ جم/لتر على بعض مركبات التمثيل الحيوى الأولية و الثانوية الهامة فى نبات الونكا ، صنفى Rosea و صنف Alba. وأشارت النتائج إلى ان القلويدات الكلية، الفينولات الذاتية الكلية والفلافونيدات كانت أعلى تركيزاً في صنف Rosea مقارنة مع صنف Alba. وأشارت النتائج أيضاً إلى ان المعاملة بمستخلص الخميرة، بصفة عامة، تؤدي إلى زيادة تراكم القلويدات الكلية، الفينولات الذاتية الكلية والفلافونيدات وكذا زيادة تركيز الكينتين، إندول حمض الخليك ، حمض الجبريليك و البنزويلادينين بينما تؤدي إلى نقص فى تركيز حمض الأبسيسيك. وعلاوة على ذلك، فإن تركيز العناصر الأساسية النيتروجين، الفوسفور، كانت اعلى في النباتات المعاملة بمستخلص الخميرة. وكان تأثير المستوى الاعلى من المستخلص (٨ جم/لتر) هو الأكثر فعالية فى هذا الشأن.

وفي ضوء النتائج المتحصل عليها يمكن إستنتاج أن مستخلص الخميرة يمكن ان يستخدم لتحسين تراكم المركبات الثانوية ذات الأهمية الطبية فى نبات الونكا.