



---

## RESEARCH ARTICLE

### Cord Blood Interleukin-6 as a Predictor of Early Onset Sepsis in High Risk Neonates

Gehan L. Abdel-Hakeem\*<sup>1</sup>, Nageh M. Shehata<sup>1</sup>, Waleed M. Abdel-Hameed<sup>2</sup>, Nashwa M. Abdel- Wahab<sup>1</sup>

\*Correspondence: Gehan L. Abdel-Hakeem. Department of Pediatrics, Faculty of Medicine, Minia University, Egypt

Email: [gehanlotfy72@yahoo.com](mailto:gehanlotfy72@yahoo.com)

Full list of author information is available at the end of the article.

---

#### Abstract

**Introduction:** The purpose of this study was to evaluate accuracy of cord blood interleukin-6 (IL-6) as an early marker of early onset neonatal sepsis (EOS) in neonates with maternal risk factors..

**Objectives:** To evaluate accuracy of cord blood IL-6 as a predictor of early onset neonatal sepsis (EONS) in neonates with prenatal risk factors for sepsis.

**Patients and methods:** The study included 80 neonates divided into two groups. Group(I) included 25 full-term and 25 preterm neonates. Group (II) included thirty healthy age and sex matched controls with mean gestational age  $35.2 \pm 3.8$ . All patients were subjected to adequate history taking, full clinical examination, complete blood count, blood culture, cord blood interleukin-6 and serum high sensitive CRP for elevated IL-6 level.

**Results:** IL-6 level was significantly higher in the group (I) compared with group(II). No significant correlation between gestational age and level of IL-6. The best cutoff value for serum interleukin-6 as a marker for early detection of neonatal sepsis is  $\geq 50$  pg/ml with sensitivity 96 % and specificity 100%.

**Conclusion:** Cord blood IL-6 can be a sensitive predictor of early onset sepsis for neonates with high risk of sepsis.

**Key words:** cord blood, interleukin-6, maternal risk factors, neonatal sepsis.

## Background

Early-onset neonatal sepsis (EONS) is a serious complication with a mortality rate ranging from 1.5% in term to almost 40% in very-low-birth weight infants.[1] Neonatal sepsis may be categorized as early-onset or late-onset. Of newborns with early-onset sepsis, 85% present within 24 hours, 5% present at 24-48 hours, and a smaller percentage present within 48-72 hours.[2] EONS is more rapid in premature neonates and it is associated with acquisition of microorganisms from the mother. Trans placental infection or an ascending infection from the cervix may be caused by organisms that colonize the mother's genitourinary (GU) tract and pass through the colonized birth canal at delivery. The microorganisms most commonly associated with early-onset infection include Group B Streptococcus (GBS), Escherichia coli, Coagulase-negative Staphylococcus, Haemophilus influenzae and Listeria monocytogenes. Different biomarkers have been used to identify patients early in the course of neonatal sepsis. Fetal inflammatory response syndrome (FIRS) has been defined by high levels of pro-inflammatory cytokines in fetal blood.[3]

Interleukin-6(IL-6)(an early marker of infection) is a rapid response inflammatory protein with a short plasma half life, in EOS, infection is commonly acquired in utero resulting in the development of chorioamnionitis. [4] Chorioamnionitis activates the fetal immune system leading to fetal inflammatory response syndrome (FIRS). In some neonates, FIRS is amplified and these neonates become symptomatic after birth with clinical manifestations suggestive of EONS. FIRS can be diagnosed by an increased number of pro-inflammatory cytokines in cord blood, of which interleukin-6 (IL-6) is one of the early mediators released in the circulation.[4,5]

## Patients and Methods

This study was carried out from December, 2012 and August, 2013. Eighty patients were selected at the delivery time from obstetric emergency department in Minia University hospital for children. Neonates involved in this study were classified into two groups:

*Group I:* Fifty newborn infants with prenatal risk factor for EONS. Twenty five were full term while 25 were preterm. Twenty nine (58%) were males while 21(42%) were females. The prenatal risk factors include any of the following: Premature rupture of membrane more than 18h, unclean foul smelly turbid vaginal secretion, foul smelling liquor, maternal pyrexia (temp.>38C), maternal leukocytosis (leukocytes>14000), dysuria and prolonged labor more than 24h.

*Group II:* Thirty newborn infants without prenatal risk factor for EONS included as control. Fifteen were full term while 15 were preterm; 12(40%) were males and 18(60%) were females.

Both groups (I&II) were subjected to thorough history taking including: gestational age, sex, obstetric history (maternal temperature, duration of labor, amount and color of vaginal secretions). General examination including anthropometric measures, vital signs, chest, heart, abdominal, neurological examination. Umbilical stump status (infected or not) and neonatal activity (doing or not doing well). Criteria of neonatal sepsis included alteration in at least 2 of the following clinical signs:

- 1-Respiratory distress: apnea, tachypnea, or hypoxemia.
- 2-Cardiological: tachycardia or bradycardia.
- 3-Hemodynamics: bad color, poor peripheral hypo perfusion, hypotension.
- 4-Neurological: irritability, lethargy, hypotonia, hypo activity, seizures.

5-Gastrointestinal: poor feeding, abdominal distension, feeding intolerance.

6-Temperature: fever  $>38^{\circ}\text{C}$  or hypothermia  $<36^{\circ}\text{C}$ .

7-Metabolic: metabolic acidosis or hyperglycemia. [6]

Plus elevated immature/ total neutrophils ratio (I/T ratio) $>0.2$  or presence of toxic granules or elevated white blood cell count  $> 25000 \text{ mm}^3$  at the time of evaluation.

Early onset neonatal sepsis is defined as: Sepsis occurs at  $<72$  hours after delivery. [7]

**Laboratory investigations carried out to neonates were included** complete blood count (CBC), blood culture, cord blood interleukin-6 level and cord blood high sensitive C reactive protein(HsCRP) were assayed immediately after collection at labor by ELISA technique using (Ani Biotech Oy, Orgenium laboratories, Finland). Cases with elevated cord IL-6levels or CRP were followed up by serum HsCRP using (ACCUBIND ELISA Microwells (Monobind inc.100 North Pointe Drive Lakeforest, CA 92630 USA).

The study was approved by the local research ethics committee of the faculty of medicine in Minia University and a written informed consent was obtained from the parents of all neonates to share in the study.

### Statistical analysis:

Data were statistically analyzed using the SPSS software package, version 16 (SPSS Inc., Chicago, IL, USA) on a personal computer. Numerical data were expressed as range, mean $\pm$ SD, median, and percentiles. Non numerical data were expressed as frequencies. Comparative studies were done using Student t test and chi square test. (p value  $< 0.05$  was considered significant). Pearson correlation test was used to detect correlation between different parameters. Receiver operating characteristic (ROC)

Curve analyses with measurement of area under the curve (AUC) were performed to identify the appropriate cut-off values.  $P<0.05$  was considered statistically significant.

## Results

Our study included 80 neonates who further subdivided into 2 groups. Group I included 50neonates with prenatal risk factors for infection and group II included 30 healthy neonates as a control. There were significant differences between both groups regarding length, Apgar score, history of prolonged labor, hospital admission ( $p<0.001$ ); birth weight( $p<0.01$ )and H.C ( $p=0.004$ ). While, there was no significant difference between both groups regarding gestational age, sex, mode of delivery.(table 1)

Regarding to clinical data, 33(66%) of group(I) neonates had respiratory distress, chest retraction (68%), cyanosis (26%), grunting (58.1%),while all neonates included were normal in group(II).There were significant differences between both groups regarding chest retraction, respiratory distress , cyanosis and grunting ( $p<0.001$ ) while no significant difference between both groups regarding convulsion, skin mottling, pallor, jaundice, abdominal distention and organomegaly (fig.1).

Regarding obstetric and maternal signs of groups I, (100%) of mothers of the neonates had PROM. Fever was found in30 mothers (60%), and lower abdominal tenderness were detected in 17 mothers (34.7%).Twenty four (48%) of the mothers had history of ante partum antibiotic, while in 48 of them (96%)there was history of intra partum antibiotics.(Figure 2).Table 2shows some laboratory data of group I neonates. Significant difference between group I and II regarding Cord blood IL-6and HsCRP ( $p<0.001$  for both ) (Table 3) Out of the 50 neonates with prenatal risk factors 20 neonates (40%) developed sepsis and blood culture were positive in 14 of these 20 neonates. ( 65%)

were Streptococci positive, (15%) were E. coli and klebsella detected in (20%) of these septicemia neonates. Table 4 shows the difference between cord IL-6 and HsCRP and some laboratory data in the infected and non infected group I neonates. In the infected group I patients, mean cord IL-6 was  $118 \pm 32$ , cord HsCRP was  $30.1 \pm 10$ , WBC was  $16.8 \pm 8.3$  and serum HsCRP after 72h was  $54.9 \pm 22.5$ . While in non infected group I patients, cord IL-6 was  $56.5 \pm 24$ , cord HsCRP was  $25.4 \pm 19.5$ , WBC was  $12 \pm 4.3$  and serum HsCRP after 72h was  $37.2 \pm 23.5$ . There were significant differences between infected and non infected group I patients regarding cord IL-6 ( $P=0.001$ ), WBC ( $P=0.001$ ) and HsCRP after 72h ( $P=0.01$ ) while no significant difference between them as regards cord blood HsCRP ( $p=0.33$ ). (table 4)

Studying the correlation between cord IL-6 level and some lab data among group I

## Discussion

IL-6 is an early and sensitive marker for neonatal sepsis.[8] In the present study, cord IL-6 was significantly higher in the case group compared with the control group. Same results were found by Cernada et al who found that serum interleukin-6 was higher in the case group than the control group.[9] Emphasizing the role of this useful inflammatory marker in detection of neonatal sepsis.[10] The inflammatory response is mediated by cytokines that are used as neonatal infection markers, especially interleukin-6 (IL-6). IL-6 is an inducer of hepatic protein synthesis, promotes production and liberation of C-reactive protein, and can be detected early when there is bacterial bloodstream invasion. It acts as a signal for T-cell activation, promotes antibody secretion by B cells and differentiation of cytotoxic T cells, and stimulates liberation of other cytokines, particularly TNF- $\alpha$ . [11]

IL-6 is an important neonatal inflammatory mediator. There is several studies demonstrating

neonates, There was significant positive correlation between IL-6 and WBCs ( $p=0.01$ ,  $r=0.33$ ), toxic granules ( $P=0.001$ ,  $r=0.46$ ) and serum HsCRP after 72h ( $p=0.002$ ,  $r=0.44$ ) while there was no correlation between cord IL-6 and Hb ( $p=0.8$ ,  $r=0.03$ ), platelets ( $p=0.9$ ,  $r=-0.01$ ), neutrophils ( $p=0.5$ ,  $r=0.1$ ) and cord blood HsCRP ( $p=0.54$ ,  $r=0.09$ ). (table 5, figure 3,4)

Cut off point of cord IL-6 was  $\geq 84$  while of cord CRP was  $\geq 21$  sensitivity of cord IL-6 was 90% and specificity was 96.7% while for cord CRP sensitivity was 85% and specificity was 56.7% and for serum HsCRP after 72h 65% and specificity was 90%. (complete term before abbreviation) NPV of IL-6 was 93.5% and (complete term before abbreviation) PPV was 94.7% while NPV of CRP was 85% and PPV 56.7%. For serum HsCRP after 72h, PPV was 81.2 and NPV 79.4 (Fig 5).

an elevation of plasma IL-6 levels in septic newborn infants. Balance between pro and anti-inflammatory cytokines is determinant for their blood levels. Low IL-10 plasma level in infected newborn infants contribute to an increase of IL-6.[12]

Significant positive correlations were found between interleukin-6 and gestational age. Greater IL-6 elevations in term infants compared to preterm infants until the inflammation becomes severe.[13] Significant differences in IL-6 levels were seen in term versus preterm infants when the acute chorioamnionitis was mild or moderate may be related to the relative immaturity of the preterm immune system, as has been demonstrated in vivo and in vitro. While others [3,5] reported that immature infants seem to be capable of synthesizing this cytokine.

HsCRP was high in the studied group compared with control group and no significant correlation between IL-6 and cord HsCRP was found while a significant positive correlation was obtained

between cord IL-6 and CRP after 72 hours. After a trauma or the invasion of microorganisms an acute localized inflammatory reaction is initiated by activation of local resident cells. The contact with bacterial endo- or exotoxins initiates the release of prostaglandins, leukotriene, and histamine, which results in vasodilatation, elevated vascular permeability, and attraction and activation of further inflammatory cells.[14,15] Activated fibroblasts, leukocytes, and endothelial cells produce pro-inflammatory cytokines including IL-1, TNF- $\alpha$ , and IL-6 are responsible for the development of fever, lethargy, arthralgia, and headache, they activate the vascular endothelial cells, regulate proliferation of T- and B-lymphocytes, activate macrophages, have pro-coagulatory effects on endothelial cells, and they induce the production of acute-phase-proteins in the hepatocytes of the liver.

Acute-phase-proteins form a heterogeneous group and include components of the complement system, coagulation factors, protease inhibitors, metal binding proteins, HsCRP, and other proteins that increase or decrease by more than 25% during an inflammatory reaction.[14,16]

HsCRP, the major acute-phase reactant in humans, derives mainly from hepatocytes in response to interleukin-6 (IL-6) and is then secreted into the systemic circulation.[17]

HsCRP level is not recommended as a sole indicator of neonatal sepsis but may be used as part of a sepsis workup or as a serial study during infection to determine response to antibiotics, duration of therapy, and/or relapse of infection.[18]

Serum concentrations of HsCRP increase several hundredfold in response to bacterial infection, making it an attractive diagnostic test for neonatal sepsis. Several hours are needed for HsCRP levels to increase in serum (including activation of neutrophils, elaboration of interleukin-6, and induction of hepatic synthesis of HsCRP) therefore limiting the sensitivity of

this test in diagnosing EONS. HsCRP levels are consistently 48 hours after the onset of infection.[19]

Various factors apart from sepsis can cause a rise in CRP levels especially Meconium aspiration syndrome, perinatal asphyxia, surgery, etc.. but a negative CRP rules out infection with high certainty i.e. it has a high Negative Predictive Value.[20]

Based on sensitivity and specificity of the results, cord blood interleukin-6 at cut off point of  $>84\text{pg/ml}$  or more, can be used as a marker for the diagnosis of EOS in neonates with suspicion of infection with.[21]

IL-6 can be a valid marker for predicting NS. It may be considered for early diagnosis and control of sepsis in neonatal care units.[22] Also, IL-6 had high sensitivity and specificity for early detection of neonatal sepsis comparing with the sensitivity of HsCRP.[21]

## Conclusions

Cord blood interleukin-6 increases significantly in neonates with risk factors for early onset neonatal sepsis can be used as an early marker to diagnose early onset sepsis. The combination of IL-6 (early and sensitive marker) with serum HsCRP (late and specific marker) can elicit better sensitivity than either marker alone.

**Conflict of interest:** The authors declared no conflict of interest.

## Author's contributions

GA and SN conceived the study. AW was responsible for laboratory investigations and data interpretation. AN revised the patients medical reports and the final manuscript. All authors revised the final draft of the manuscript.

## Acknowledgment

The author would like to thank the nurse staff of neonatal intensive care unit in El Minia university hospital for children for their assistance in the cord blood and blood cultures sampling.

#### Author's details

<sup>1</sup>*Pediatric Department, Faculty of Medicine, Minia University, Egypt.*

<sup>2</sup>*Clinical pathology Department, Faculty of Medicine, Minia University, Egypt*

**Date received:** 11<sup>th</sup> March 2019. Accepted 9<sup>th</sup> April 2019. Published 29<sup>st</sup> June 2019.

## References

1. Weston EJ, Pondo T, Lewis MM, Martell-Cleary P, Morin C, Jewell B, Daily P, Apostol M, Petit S, Farley M, Lynfield R, Reingold A, Hansen NI, Stoll BJ, Shane AL, Zell E, Schrag SJ The burden of invasive early-onset neonatal sepsis in the United States. *Pediatr Infect Dis J* 2011; 30(11): 937–41.
2. Klinger G, Levy I, Sirota L, Boyko V, Reichman B, Lerner-Geva L Epidemiology and risk factors for early onset sepsis among very-low-birthweight infants. *Am J ObstetGynecol* 2009; 201(1):38.e1-6.
3. Messer J, Eyer D, Donato L, Gallati H, Matis J, Simeoni U Evaluation of interleukin-6 and soluble receptors of tumour necrosis factor for early diagnosis of neonatal infection. *Pediatr J* 1996; 129: 574–580.
4. Malik A, Hui CP, Pennie RA, Kirpalani H. Beyond the complete blood cell count and C-reactive protein: a systematic review of modern diagnostic tests for neonatal sepsis *Arch Pediatr Adolesc Med* 2003; 157:511-6.
5. Shigeru Nishimaki, Miho Sato, Hiromi An, Yoshio Shima, Toru Akaike, Utako Yokoyama and Shumpei Yokota Comparison of markers for fetal inflammatory response syndrome: fetal blood interleukin-6 and neonatal urinary beta(2)-microglobulin. *J ObstetGynaecol Res* 2009; 35:472-6.
6. Alexander JM1, Gilstrap LC, Cox SM, McIntire DM, Leveno KJ Clinical chorioamnionitis and the prognosis for very low birth weight infants. *Obstet Gynecol* 1998; 91:725-9.
7. Van den Hoogen A, Gerards LJ, Verboon-Macielek MA, Fleer A, Krediet TG Long-Term Trends in the Epidemiology of Neonatal Sepsis and Antibiotic Susceptibility of Causative Agents. *Neonatology* 2009; 97(1):22-28.
8. Renato S. Procianny1, Rita C. Silveira2 The role of sample collection timing on interleukin-6 levels in early-onset neonatal sepsis *Jornal de Pediatria* 2004; 0021-7557/04/80-05/407.
9. María Cernada, Natalia Badía, Vicente Modesto, Ricardo Alonso, Asunción Mejías, Sergio Golombek5, Máximo Vent Cord blood interleukin-6 as a predictor of early-onset neonatal sepsis *Acta Pædiatrica* 2013; 0803–5253.
10. Marzie Heidarzadeh Arani, Amir hosein Movahedian, Mohammad Ghannae Arani Mojtaba Adinah , Ziba Mosayeb Predictive Value of Interleukin-6 (IL6) in Term Neonates with Early Sepsis during 2010-2011 *Jundishapur J Microbiol* 2013; 6(10): e8580.
11. Buck C, Bundschu J, Gallati H, Bartmann P, PohlandtF Interleukin-6: a sensitive parameter for the early diagnosis of neonatal bacterial infection. *Pediatrics* 1994; 93(1):54-8.
12. Damman O, Leviton A Brain damage in preterm newborns: Biological response modification as a strategy to reduce disabilities *J Pediatr* 2013; 2000.136:433-8.
13. Rogers BB, Alexander JM, Head J, McIntire D, Leveno KJ Umbilical vein

- interleukin-6 levels correlate with the severity of placental inflammation and gestational ageHumPathol (2002); 33(3):335-40.
14. Jaye DL and WaitesKB Clinical applications of C-reactive protein in pediatrics. *Pediatr Infect Dis* 2002; 16: 735–746.
  15. Du Clos TW Function of C-reactive protein. *Annals of Medicine* 2003; 24:2748.
  16. Volanakis JE Human C-reactive protein: expression, structure, and function. *Molecular Immunology* 2001; 38(2-3):189-97.
  17. Rattazzi M, Puato M, Faggini E, Bertipaglia B, Zambon A, Pauletto P C-reactive protein and interleukin-6 in vascular disease: culprits or passive bystanders? *J Hypertens* 2003; 21:1787–1803.
  18. Anderson-Berry AL, Bellig LL and Ohning BL Neonatal sepsis. *eMedicine* 2010; 188: 1157-59.
  19. Ehl S, Gering B, Bartmann P, Hogel J, Pohlandt F C-reactive protein is a useful marker for guiding duration of antibiotic therapy in suspected neonatal bacterial infection. *Pediatrics* 1997; 99(2):216-21.
  20. M Himayun, S Ahmad, A Rasool *Role Of C-Reactive Protein In Early Onset Neonatal sepsis*. *The Internet Journal of Pediatrics and Neonatology* 2009; (11) : 2.
  21. BeceiroMosquera J, SiveraMonzo CL, Oria de Rueda Salguero O, Olivas López de Soria C, HerbozoNory C Usefulness of a rapid serum interleukin-6 test combined with C-reactive protein to predict sepsis in newborns with suspicion of infection. *An Pediatr (Barc)* 2009; 71(6):483–8.
  22. .LobatShahkar , AbbasaliKeshtkar, Arezou Mirfazeli<sup>2</sup>, Ali Ahani<sup>2</sup> and Gholamreza Roshandel The Role of IL-6 for Predicting Neonatal Sepsis: A Systematic Review and Meta-Analysis. *Iran J Pediatr* 2011; 21 (4): 411-41.

**Citation:** El-Mazary MA. Cord Blood Hemoglobin A1c and MCP-1 as predictors for Cardiomyopathy in infants Mothers with gestational diabetes. *Ann Neo J*. 2019, 1(2): 59-71.



**Copyright:** El-Mazary MA et al, 2019. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (4).

