

## Genetic Evaluation for Some Productive and Reproductive Traits by Using Animal Model in A Commercial Friesian Herd in Egypt

El-Awady, H. G.; A. E. Abd el-Khalek and M. Abo Elreesh

<sup>1</sup>Animal Production Department, Faculty of Agriculture, Kafrelsheikh University, PC: 33516, Kafrelsheikh, Egypt, E-mail hassanelawady63@yahoo.com

<sup>2</sup>Animal Production Department, Faculty of Agriculture, Mansoura University, PC: 35516, Mansoura, Egypt.



### ABSTRACT

Data used in this investigation were collected from 4370 records for 874 purebred Friesian cows' daughters of 44 sires from 1985 to 2003, which belong to Shobratana Animal Production Society, located in the north part of Delta region near Tanta city in Egypt. Variance and covariance components for 305 day milk yield (305- dMY, kg), lactation length (LL, day), dry period (DP, day), days open (DO, day) and calving interval, (CI, day) were estimated with REML using an animal model (Boldman, 1955), that included the fixed effects of month and year of calving and parity of cow. Random effects were animal, direct and maternal genetic effects, maternal permanent environmental effect and random effect. Overall means of (305-dMY), LL, DP, DO and CI were 4897 kg, 327 day, 120 day, 66.3 day and 401.16 day. Month and year of calving were highly significant effect ( $P < 0.01$ ) for all studied traits. Direct heritability ( $h^2_a$ ) of the mentioned traits was 0.38, 0.13, 0.14, 0.11 and 0.16, respectively. Heritabilities of maternal effects ( $h^2_m$ ) of (305-dMY) (0.13), LL (0.18), DP (0.23), DO (0.14) and CI (0.21), respectively. Estimates of direct-maternal genetic correlations in all traits studied ranging from -0.60 to 0.74. Genetic correlations among different studied traits ranged from -0.80 to 1.00 while the phenotypic correlations among investigated traits were positive, ranging from 0.07 to 0.99. Range of predicted breeding value (PBV's) of sires were 462 kg, 1.05 d, 0.24 d, 5.93 d and 2.26 d, for 305dMY, LL, DP, DO and CI, respectively. PBV's of cows of the mentioned traits was 1123 kg, 2.92 d, 1.64 d, 5.93 d and 2.92 d respectively. Corresponding PBV's of dams for the above same traits were 510 kg, 1.42 d, 0.80 d, 7.20 d and 3.00 d, respectively. Accuracy of (PBV's) ranged from 38 to 88, 73 to 88 and 39 to 89% for sires, cows and dams, respectively; indicating the genetic improvement could be attained through any bath of them. Estimates of temporary environmental variance as a proportion of phenotypic variance for LL, DP, DO and CI ranged from 35 to 71% indicating the improvement in these traits could be achieved by improving for these temporary factors.

**Keywords:** Direct and maternal genetic parameters, breeding value, Friesian cows and Animal model.

### INTRODUCTION

Friesian cattle are the most popular dairy cattle and also represent a major source of red meat in Egypt. The success of selection for milk production has contributed to the domination of the Friesian breed around the world. In Egypt, dairy industry represents 35% of the total animal production sector. During the last two decades, considerable emphasis had been placed upon the importance of Friesian cattle in Egypt for milk production, accordingly the number of large Friesian herds had increased either in the governmental or commercial farms through importation from Europe and USA (Shalaby *et al.*, 2001). Main reasons of low productivity of farm animals are; non-descript breed, poor management, lack of nutrition, lack of resources, low inputs; inadequate artificial insemination service and diseases. These causes lead to low average milk production, late age at first calving, delayed conception, impaired fertility, long calving intervals (Khan *et al.*, 2008). The performance of Holstein-Friesian cattle for milk production affected by different factors such as animal genetic merit, calving season, administrative, feeding and diseases had reported by (Epaphras *et al.*, 2004). Aziz *et al.*, (2014) estimate the heritability for lactation milk yield (0.26), lactation length (0.10), dry period (0.06), calving interval (0.07), gestation length (0.06) and open days (0.03). They added that the genetic and phenotypic correlations between milk yield and fertility traits were positive except between LL and DP. Sharma, (1982), Roy *et al.*, (1985) and Abubakar *et al.*, (1987) reported that the reproductive traits could improved by sire selection. Calving interval is one of the most important measures of reproductive efficiency in dairy cattle, early postpartum breeding in dairy

animals, high fertility, short dry period and early maturity are resulted in more calves and high milk yield per unit of time throughout the herd life (Britt, 1975). With the help of heritability one can predict the breeding value of the individual. The magnitude of heritability dictates the choice of the selection method and breeding system (Paul *et al.*, 2003). Addition, a goal of dairy cattle breeder is to increase genetic merit of milk production to determine the effectiveness of breeding programs. Falconer and Mackay, (1996) estimation of  $h^2$  for any trait can be calculated from sire model or animal model. Genetic variance might be underestimated if selection intensity is higher for males than for females because analyses with sire models accounted only for genetic variance of sires (Albuquerque *et al.*, 1995). Animal model take into account differential selection of males and females and might provide more accurate estimates of parameters than do sire model. Gutierrez *et al.*, (1995) with Spanish Holstein Friesian cows, comparison between sire and animal model, suggested that a sire model based estimating procedure for genetic parameters may be preferred when a small number of individuals, little pedigree information and highly unbalanced distribution of effects. Also, the same authors concluded that the cheapest in terms of computing costs was based on a sire model and the most expensive on an animal model.

The objectives of the present study were estimate some non-genetic factors affecting some productive (305-dMY, LL and DP) and reproductive traits (DO and CI) and estimate the genetic and phenotypic parameters and breeding values for these traits by using Animal Model.

## MATERIALS AND METHODS

### Data and managements:

Data used in this investigation were collected from 4370 records relevant to 874 purebred Friesian cows which belong to Shobratana Animal Production Society, located in the North Part of Delta region near Tanta city in Egypt. These cows were daughters of 44 sires. The records used covered the period from 1985 to 2003. The following five productive and reproductive traits were analyzed, 305 day milk yield (305-dMY, kg), lactation length (LL, day), dry period (DP, day), days open (DO, day) and calving interval (CI, day). Animals were housed free in shaded open yards covered with 3.5 - 4 meters high roofs, grouped according to their average daily milk yield and fed ad libitum on berseem (*Trifolium alexandrinum*) and rice straw in addition concentrates feed mixture from December to April (green season) and mature cows fed daily on ration consisting of cotton seed cakes, barley wheat and rice bran (not less than 18% protein from May to November (dry season). Mineral mixture bricks were offered ad libitum as lick salt in front animals, and on balanced ration of a concentrates according to their production and weigh. Rice straw and water were offered ad libitum. Limited amount clover hay when available. Cows were artificially inseminated during heat, which occurred after 60 days postpartum using imported frozen semen from USA and Canada. Heifers were artificially inseminated for the first time in the first two heats once they attained 350 kg or 18-22 months of live body weight. Pregnancy diagnoses were carried out routinely at 60 days after service by rectal palpation. If conception did not occur or the cows were seen in estrus, the cows were inseminated again. The cows were machine milked three times a day at 04.00h, 12.00 and 19.00h. The calves born were artificially suckled after calving to weaning excluding colostrums period.

### Statistical analysis:

Data were analyzed using the two methods, firstly, by using Linear Mixed Model Least Squares and Maximum Likelihood (LSMLMW) computer program of Harvey (1990) for obtained the main effects. The following model was used in the first analysis:

$$Y_{ijkl} = \mu + S_i + M_j + Y_k + P_l + \beta (\text{Age}) + e_{ijklm} \quad (1)$$

### Where:

$Y_{ijkl}$  = the individual observation,

$M$  = the overall mean,

$S_i$  = the random effect of the  $i^{\text{th}}$  sire,  $i = 1$  to 44,

$M_j$  = the fixed effect of the  $j^{\text{th}}$  month of calving;  $j = 1, 2, 3, \dots, \text{and } 12$  (January, February, March, .... and December),

$Y_k$  = the fixed effect of the  $k^{\text{th}}$  year of calving,  $k = (85, 86, 87, \dots, 2003)$ ,

$\beta$  = Partial linear and quadratic regression coefficients, respectively for different traits studied on age at first calving (months).

$e_{ijklm}$  = Residual term assumed to be random and distributed as a normal distribution with mean zero and variance  $\sigma^2_e$ .

Secondly, data analyzed to estimate the variance and covariance component with derivative-free restricted a maximum likelihood (REML) procedure using the MTDFREML program of Boldman *et al.*, (1995). The multiple models were included the additive

genetic effect of animal, maternal genetic effect and the maternal effect due dam (permanent environmental), which allowed for estimation of the genetic covariance between direct and maternal genetic effects as follows:

$$Y = X\beta + Za + Mm + Wpe + e$$

**Where:**  $Y$  = a vector of observations,  $\beta$  = a vector of fixed effects,  $a$  = vector of additive genetic effects,  $m$  = vector of maternal genetic effects,  $M$  = the incidence matrix relating records to maternal genetic effect,  $pe$  = vector of environmental effects contributed by dams to records of their progeny (permanent environmental),  $W$  = is the incidence matrix relating records to permanent environmental effects and  $e$  = vector of the residual effects.  $X$  and  $Z$  are incidence matrices relating records to fixed and genetic effects, respectively. The variance and covariance structure for model was as follows:

$E(y) = Xb$  and

$$V = \begin{bmatrix} a \\ m \\ pe \\ e \end{bmatrix} = \begin{bmatrix} A\sigma^2_a & A\sigma_{am} & 0 & 0 \\ A\sigma_{am} & A\sigma^2_m & 0 & 0 \\ 0 & 0 & I_d\sigma^2_{pe} & 0 \\ 0 & 0 & 0 & I_N\sigma^2_e \end{bmatrix}$$

**Where:**  $d$ , is the number of dams and  $N$  is the number of records,  $A$  is the number relationship matrix among animals,  $\sigma^2_a$  is the additive genetic variance,  $\sigma^2_m$  is the maternal genetic variance,  $\sigma_{am}^2$  is the additive and maternal genetic covariance,  $\sigma^2_{pe}$  is the maternal permanent environmental variance and  $\sigma^2_e$  = the random residual effect associated with each of observation. Estimates of additive or the direct ( $h^2_a$ ) and maternal ( $h^2_m$ ) heritabilities were calculated as ratios of estimates of additive ( $\sigma^2_a$ ) and maternal genetic ( $\sigma^2_m$ ) variance, respectively to phenotypic variance ( $\sigma^2_p$ ). The direct maternal correlation ( $r_{am}$ ) was computed as the ratio of the estimates of direct maternal covariance's ( $\sigma^2_{am}$ ) to the product of the square roots of estimates of  $\sigma^2_a$  and  $\sigma^2_m$ .  $\sigma^2_{pe}$  is the ratio of estimates of maternal environmental variance ( $\sigma^2_{pe}$ ) to the total phenotypic variance ( $\sigma^2_p$ ).  $I_d, I_N$  are identity matrices of appropriate order, the number of dam and number of animals with records respectively.

Multi-trait animal model (MTAM) was used to estimate BLUP values. The mixed model equations (MME) used for best linear unbiased estimation (BLUE) of function (b) and for the BLUP of (a) and (pe) in matrix notation as follows:

$$\begin{bmatrix} X'X & X'Z & X'W \\ Z'X & Z'Z + A^{-1}\alpha_1 & Z'W \\ W'X & W'Z & W'W + I\alpha_2 \end{bmatrix} \begin{bmatrix} \hat{b} \\ \hat{a} \\ p_e \end{bmatrix} = \begin{bmatrix} X'y' \\ Z'y' \\ W'y' \end{bmatrix}$$

Where:

$$\alpha_1 = \frac{\sigma^2_e}{\sigma^2_a} \quad , \quad \alpha_2 = \frac{\sigma^2_e}{\sigma^2_{pe}}$$

Convergence reached when the simplex variance was less than 10<sup>-9</sup> and then several extra rounds of iterations were executed to ensure that a global maximum was reached. Best linear unbiased perdition (BLUP) of estimated breeding values was calculated by using MTDFREML program for animals in pedigree file for multi-traits analysis.

## RESULTS AND DISCUSION

### Statistical descriptive:

Unadjusted means and their standard deviations (SD) and coefficient variabilities (C.V) for different studied traits are presented in Table 1.

**Table 1. Unadjusted means, standard deviations (SD) and coefficient variabilities (C.V) for 305 day milk yield (305dMY), lactation length (LL), dry period (DP), days open (DO) and calving interval (CI) in a commercial herd of Friesian cows in Egypt.**

Trait	Mean	SD	CV%
305dMY (Kg)	4897	1543	31.51
LL (day)	327	69.92	21.38
DP (day)	66.30	21.04	31.73
DO (day)	120	64.71	53.93
CI (day)	401.16	80.30	20.02

The present overall mean of 305dMY was lower (4897 kg) than those estimated by Khattab *et al.*, (2000) (5076 kg) from a commercial herds of Holstein Friesian cattle in Egypt and Shalaby *et al.*, (2013) (5387 kg) working with another set of Friesian cows at the same farm and was higher than the published estimates on Friesian cattle in Egypt 2655 kg (Oudah *et al.*, 2010) and 3936 kg (El-Awady and Oudah, 2011). In addition, 3710 kg obtained by Tadesse *et al.*, (2010) for Friesian cows in Ethiopia.

The present overall mean of LL was 327 days. The same results found by Shalaby *et al.*, (2013) working with another set of Friesian cows at the same farm. However a shorter mean (291.86 days) was stated by Sattar *et al.*, (2005).

The present overall mean of DP (66.30 days) was shorter than that reported by Oudah *et al.*, (2001) on Friesian cattle in Egypt (79.3 days) and Shalaby *et al.*, (2013) was (72.9). But it was longer than that obtained by Khattab and Atil (1999) being 65 days. The overall means of DO and CI reported in the present study were 120 days and 401.16 days, respectively (Table 1). These values were lower than the estimates (141 and 422 days, respectively) reported by Shalaby *et al.*, (2001) on a similar Friesian herd in Egypt. In this study, the coefficients of variations (CV %) ranged from 20.02 for CI to 53.93% for DO. Such large coefficients of variation are indicative leaders for opportunities for improvement in these traits. The differences between our findings and other investigators may be related to genetic and environmental factors from differences statistical models.

The present results showed that the sire was highly significant ( $P < 0.01$ ) effect on all traits studied under investigation as shown in Table 2. These results are in agreement with the findings obtained by Oudah *et al.*, 2008 on DO and CI, Shalaby *et al.*, 2001 on TMY.

The present results indicate that the month and year of calving were highly significant ( $P < 0.01$ ) on all traits under investigation and shown that parity was significant ( $P < 0.05$  and  $P < 0.01$ ) on dry period, day open and calving interval (Table 2). These results are in agreement with Shalaby *et al.* (2001) found that month of calving was highly significant effect on DO and CI.

On the other hand, EL-Barbary *et al.*, (1992) found that the parity had no significant on total milk yield of Friesian cows in Egypt. Table 2 showed that the age at first calving as linear and quadratic regression coefficient had significant ( $P < 0.05$  and or  $P < 0.01$ ) on all traits under investigation except effect of quadratic regression on DO and effect of linear regression on CI.

**Table (2):Least square analysis of variance for 305 day milk yield (305dMY), lactation length (LL), dry period (DP), days open (DO) and calving interval (CI) in a commercial herd of Friesian cows in Egypt.**

S.O.V	d.f	F- Significance				
		305dMY	LL	DP	DO	CI
Sires	43	3.34**	1.74**	93.19**	10.73**	35.15**
Month of calving	11	23.14**	3.58**	10.83**	4.54**	15.85**
Year of calving	9	137.50**	25.53**	104.39**	11.25**	55.33**
Parity	4	.01 <sup>ns</sup>	.11 <sup>ns</sup>	8.69**	2.93*	1.11*
Regression						
AFC (L)	1	270.80**	283.94**	275.40**	385.76**	0.27 <sup>ns</sup>
AFC (Q)	1	109.12**	73.68**	997.70**	1.09 <sup>ns</sup>	2.72*
Remainder	2623	20942.41	51.35	71.29	81.05	22746.11

\*\* significant at  $P \leq 0.01$  \* significant at  $P \leq 0.05$  Ns not significant

### Genetic and phenotypic parameters:

Estimates of variance and covariance components are shown in Table 3 while in the Table 4 are those for other genetic parameters.

Direct heritability for 305dMY (0.38) was moderate, while for LL, DP, DO and CI were stingy, being 0.13, 0.11, 0.14 and 0.16, respectively. The estimates of maternal heritability for LL, DP, DO and CI were 0.18, 0.14, 0.23 and 0.21, respectively (Table 4) and considered reasonable value compared with  $h^2_m$  for 305dMY (0.13). This value is generally higher than to those obtained for Holstein cattle raised in Egypt (Salem *et al.*, 2006; El-Arian *et al.*, 2003 and Alhammad, 2005) which ranged for LP, DP and CI from (0.03 to 0.12 ), (0.003 to 0.03) and (0.003 to 0.09), respectively.

The present estimates indicate the direct heritability ( $h^2_a$ ) of 305dMY is approximately 3.0 times their respective maternal part, while the maternal heritability for other traits investigated almost is close with direct heritability but may be higher as shown in table 4. The present estimates of maternal heritability suggest that maternal effects need to be considered in the analysis model and the selection for productive and reproductive traits in Friesian cows in Egypt. Also, in present study, the maternal permanent environmental effect on 305dMY, LL, DP, DO and CI due to dams accounted for 19%, 21%, 4%, 28% and 9%, respectively from the phenotypic variance.

The estimates of genetic correlations between 305dMY and each of LL, DP and CI were positive and high, being 0.79, 0.92 and 0.98, respectively. In this respect, the estimates of genetic correlations between LL and each of DO and CI were 0.91 and 0.95 respectively. The present results are in the desirable direction indicating that high yielding cows are also having longer LP and shorter DP. Similar results were obtained by Salem and Abdel-Raouf (1999), Shitta *et al.* (2002) and El-Arian *et al.* (2003) on Friesian and Holstein cattle.

**Table(3)Estimates of direct and maternal genetic variance and covariance components, permanent environmental and residual of different studied traits using animal model.**

Item	Direct and maternal genetic variances and covariances										
	Direct	305dMY, a	LL, a	DP, a	DO, a	CI, a	305dMY, m	LL, m	DP, m	DO, m	CI, m
305dMY, a		460.35									
LL, a		168.95	99.35								
DP, a		-17.94	-68.76	109.26							
DO, a		155.47	71.44	7.41	62.03						
CI, a		291.53	131.29	15.94	109.20	192.23					
305dMY, m		94.90	88.18	-13.21	49.77	105.13	159.71				
LL, m		99.43	37.88	-31.97	48.18	62.97	65.60	85.90			
DP, m		-42.90	-9.38	-30.74	-25.01	-34.25	-89.19	-9.81	138.35		
DO, m		-134.53	-79.92	-20.41	-12.14	-35.63	76.64	27.63	-39.90	105.66	
CI, m		-136.47	-63.40	-9.21	-53.18	-108.54	91.51	39.91	-65.61	66.39	95.76
Permanent environmental											
			305dMY		LL		DP		DO		CI
305dMY, kg			231.28								
LL, d			13.82		101.98						
DP, d			-23.43		32.42		41.22				
DO, d			-91.86		6.91		-35.12		129.89		
CI, d			27.87		2.55		10.55		46.09		39.92
Residual											
305dMY, kg			365.18								
LL, d			23.19		230.03						
DP, d			-50.15		318.40		688.64				
DO, d			43.54		-63.35		-149.48		160.23		
CI, d			121.60		211.84		-146.62		76.63		240.86
Phenotypic*											
305dMY, kg			1217.262								
LL, d			618.65		479.233						
DP, d			119.52		88.63		969.916				
DO, d			716.65		323.19		46.65		457.808		
CI, d			729.49		342.14		124.97		411.22		446.052

**Table (4): Direct and maternal heritabilities (diagonal), direct and maternal genetic correlations and direct maternal genetic correlations (below the diagonal), residual variance (temporary variance) as the proportion from phenotypic variance and phenotypic correlations between different traits investigated using animal model.**

Item	Direct and maternal genetic variances and covariances										
	Direct	305dMY, a	LL, a	DP, a	DO, a	CI, a	305dMY, m	LL, m	DP, m	DO, m	CI, m
305dMY, a		.38±.01									
LL, a		0.79	.13±.01								
DP, a		-0.08	-0.66	.11±.03							
DO, a		0.92	0.91	0.09	.14±.02						
CI, a		0.98	0.95	0.11	1.00	.16±.02					
305dMY, m		0.35	0.70	-0.10	0.50	0.60	.13±.07				
LL, m		0.50	0.41	-0.33	0.66	0.49	0.56	.18±.02			
DP, m		-0.17	-0.08	-0.25	-0.27	-0.21	-0.60	-0.09	.14±.13		
DO, m		-0.61	-0.78	-0.19	-0.15	-0.25	0.59	0.29	-0.33	.23±.18	
CI, m		-0.65	-0.65	-0.09	-0.69	-0.80	0.74	0.44	-0.57	0.66	.21±.14
Permanent environmental											
			305dMY		LL		DP		DO		CI
305dMY, kg			0.19±0.11								
LL, d			0.09		0.21±0.09						
DP, d			-0.24		0.50		0.04±0.02				
DO, d			-0.53		0.06		-0.48		0.28±0.18		
CI, d			0.29		0.04		0.26		0.64		0.09±0.03
b- Residual											
305dMY, kg			0.30±0.02								
LL, d			0.08		0.48±0.30						
DP, d			-0.10		0.80		0.71±0.19				
DO, d			0.18		-0.33		-0.45		0.35±0.09		
CI, d			0.41		0.90		-0.36		0.39		0.54±0.16
c- Phenotypic											
305dMY, kg			1217.262								
LL, d			0.81		479.233						
DP, d			0.11		0.13		969.916				
DO, d			0.96		0.69		0.07		457.808		
CI, d			0.99		0.74		0.19		0.91		446.052

The phenotypic correlations between 305dMY and each of LL, DO and CI were 0.81, 0.96 and 0.99, respectively. In addition, the phenotypic correlation between DO and CI was 0.91 (Table 4). Estimates of genetic maternal correlations ( $r_m$ ) between MY and all traits studied except DP were moderate and positive. The estimates of correlations between direct and maternal genetic effects were negative, except among 305dMY, LL and DO were positive in Table 4. Khattab *et al.*, (2005) found the same trend.

The results in table 4 agreement with Salem *et al.*, (2006), found the genetic correlation between LP and 305d MY was positive being 0.43, and the phenotypic correlation between LP and DP were negative and high (-0.89) on Friesian and Holstein cattle

Predicted breeding value of sires, cows and dams for studied traits 305dMY, LL, DP, DO and CI are present in Table 5. The benefit of abundant and good measurement is that we may be better able to identify the genetically superior animal. This leads to more accurate selection and more genetic improvement. Accuracy of predicted breeding value, ranged from 38 to 88, 73 to 88 and 39 to 89% for sires, cows and dams, respectively, indicate that genetic improvement could be attained through each of cows or sires or dams.

**Table (5) Range of breeding values through sires, Cows and Dams and its accuracies, % for 305 day milk yield (305dMY), lactation length (LL), dry period (DP), days open (DO) and calving interval (CI) in a commercial herd of Friesian cows in Egypt.**

Traits	Breeding Values			Range
	Minimum	Maximum	Accuracy, % <sup>1</sup>	
<b>Breeding Values of Sires</b>				
305dMY (Kg)	-301±127	161±143	73-79	462
LL (day)	-0.66±1.51	0.39±1.61	38-50	1.05
DP (day)	-0.096±1.69	0.152±2.33	54-79	0.248
DO (day)	-2.58±0.88	3.35±0.96	80-81	5.93
CI (day)	-0.981±1.11	1.28±1.14	88-88	2.261
<b>Breeding Values of Cows</b>				
305dMY(Kg)	-486±135	637±189	73-76	1123
LL (day)	-1.27±1.11	1.65±1.19	73-77	2.92
DP (day)	-1.21±1.52	0.432±1.62	81-84	1.642
DO (day)	-2.46±0.91	3.47±0.94	78-79	5.93
CI (day)	-0.91±1.11	2.01±1.38	81-88	2.92
<b>Breeding Values of Dams</b>				
305dMY(Kg)	-278±192	232±172	39-57	510
LL (day)	-0.71±1.66	0.71±1.61	38-39	1.42
DP (day)	-0.48±2.43	0.32±2.46	46-48	0.80
DO (day)	-3.07±0.89	4.13±0.92	79-80	7.20
CI (day)	-0.91±1.09	2.09±1.14	88-89	3.00

<sup>1</sup>Accuracy%=

The present investigation showed that more variation of breeding values and standard errors, this increase the possibility of selection for these traits. Concerning to accuracy of breeding values, the highest accuracy was the value of CI, while the lowest accuracy was the value of LL. High accuracy levels of breeding values help breeders to select for traits in their animals and hence genetic improvement in herds. The present results with agreement found by Radwan *et al.*, (2015) working in Holstein Friesian cows raised in Egypt.

Abd El-Harith *et al.*, (2002) concluded that breeding values for 305dMY by using the multiple-traits analysis is recommended to obtain more accuracy

because it makes use of all the information about the lactation and the covariance among them as the relationship between the relative indifferent traits working in Friesian cows in Egypt.

## CONCLUSION

The present results had shown that the estimates of the mean values for traits under investigation in normal range with other studies Commercial Friesian Herd in Egypt. The present heritability estimates for 305-dMY emphasized the possibility of realizing a considerable rate of genetic improvement in this trait under investigation. Genetic enhancement for 305dMY following up improve each of LL, DP and CI. The estimates of heritabilities were low for some traits studied indicated that the major part of the variation in these traits was environmental and selection may not prove effective in bringing about genetic improvement in these traits. Therefore, better management can play a major role in improving these traits. Estimated of breeding values were high ranges indicated the existence of more genetic variation among individuals and hence increase the possibility dairy cattle selection for these traits, which reduce bias from selection and achieve the best accuracy of predictions. In the future this should translate into good herds without productive and reproductive problems which would lead to more and rapid genetic improvement. The accuracies of predicted breeding values, similarly for sires, cows and dams indicate that genetic improvement could be attained through any bath of sires or cows or dams. Also, it could be concluded that selection for traits under this investigation based on direct genetic components only may not give optimal response because of the negative genetic correlation between direct and maternal effects. Therefore, future selection plans for productive and reproductive traits need to consider maternal effects in the analysis models in order to optimize expected total response over the long run.

## REFERENCES

- Abd El-Harith, H. H.; E. A. El-Tawil and M. F. Adel Gelil (2002). Different animal models for estimating breeding values using earlier records in Friesian herd in Egypt. *Egypt J. Anim. Prod.* 39: 9-19.
- Abubakar ,B. Y., Mc Dowell , R. E. , Van Vleck , L.D. and E. Cabello.(1987). Phenotypic and genotype parameters for Holsteinsin Mexico. *Trop. Agric (Trinidad)*, 64: 23 -26.
- Albuquerque. L. G.; G. Dimov, and J. F. Feown (1995). Estimates using an animal model of (co) variances for yields of milk, fat, and protein for the first lactation of Holstein cows in Californian and New York. *J. Dairy Sci.*, 78:1591.
- Alhammad, H.O.A., (2005). Phenotypic and genetic parameters of some milk production traits of Holstein cattle in Egypt. M. Sc. Thesis Fac. Agric. Cairo Univ., Cairo, Egypt.

- Aziz S., Z. Ulutaş, A. Y. Adkinson., R. W. Adkinson (2014). Genetic parameters of first lactation milk yield and fertility traits in Brown Swiss cattle *Ann. Anim. Sci.*, 14:545–557.
- Boldman, K. G.; L. A. Kriese; L. D. Van Vleck and S. D. Kachman (1995). A manual for use of MTDFREML. A set for programs to obtain estimates of variances and covariance (DRAFT). U.S. Department of Agriculture, Agriculture Research Service, Clay Center, NE, USA
- Britt, G. H. (1975). Early postpartum breeding in dairy cows A. review. *J. dairy Sci.*, 55: 266.
- El-Arian, M.N., H.G. El-Awady and A.S. Khattab, (2003). Genetic analysis for some productive traits of Holstein Friesian cows in Egypt through MTDFREML program. *Egypt. J. Anim. Prod.*, 40: 99.
- El-Awady and Oudah (2011). Genetic and Economic Analysis for the Relationship between Udder Health and Milk Production Traits in Friesian Cows *Asian-Aust. J. Anim. Sci.* 24:1514-1524.
- El-Barbary, A. S. A.; F. F. M. Khalil and A. E. Mahady (1992). Studies on Danish Friesian cattle in Egypt. Some factors affecting reproductive traits. *J. Agric. Sci., Mansoura Univ.*, 17: 2130-2135.
- Epaphras, A., E.D. Karimuribo and S. N. Msellem (2004). Effect of season and parity on lactation of crossbred Ayrshire cows reared under coastal tropical climate in Tanzania. *Livest. Res. Rural. Dev.* 6 (16) Online: URL: <http://www.cipav.org.co/lrrd/lrrd16/6/epap16042>.
- Falconer and Mackay, (1996). Introduction to quantitative genetics Fourth edition, longman.
- Gutierrez, J. P.; J. Canon; M. Rico (1995). Comparison of two models for estimation of variance components in a sample of Spanish Holstein Friesian. *J. Anim. Breed. Genet.* 111: 169-174.
- Harvey, W.R. (1990):User's guide for LSMLMW, mixed model least squares and maximum likelihood computer program.PC-2 version 2, Ohio state Univ., Columbus(Mimeograph), USA.
- Khan, M. S; Z. M Rehman, A. Khan and S. Ahmad, (2008). Genetic resources and diversity in Pakistani cattle. *Pakistan Vet. J.*, 28: 95-1020.
- Khattab, A. S.; M. N. El-Arian and H. Atil (2000). Estimation of producing ability of Holstein Friesian cattle in a commercial herd in Egypt. *Pakistan J. Biology. Sci.*, 3:1076.
- Khattab, A. S.; H. Atil and L. Badawy (2005). Variances of direct and maternal genetic effects for milk yield and age at first calving in a herd of Friesian cattle in Egypt. *Arch. Tierz., Dummerstorf* 48:24-31.
- Khattab, A.S. and H. Atil, (1999). Genetic study of fertility traits and productive in a local born Friesian cattle in Egypt. *Pak. J. Biol. Sci.*, 2: 1178-1183.
- Oudah, E. Z. M.; M. A. Mostafa; M. N. El-Arian and R. F. Ismail. (2008). Productive and reproductive performance of Russian Black Pied cows in the first three lactations under hot climate conditions in Uzbekistan. *J. Agric. Sci., Mansoura Univ.*, 33: 4063-4075.
- Oudah, E. Z. M. and Zainab A. Khalefa (2010). Genetic evaluation for Friesian Cattle in Egypt using single-trait animal models. *J. Animal and Poultry Production, Mansoura University*, Vol. 1: 371-381
- Oudah, E. Z. M.; N. A. Shalaby and M. A. Mustafa. (2001). Genetic and non- genetic factors affecting days open, number of service per conception and age at first calving in a herd of Holstein-Friesian cattle. *Pakistan Journal Biological Sciences*, 4: 740-744.
- Paul A.K., I. Singh, and V.K. Bhatia (2003). A note on estimation of heritabilities of different dairy characteristics by different methods. *Indian J. Anim. Res.* 37: 153-154.
- Radwan H. A. A; E. A. Abo Elfadl and A. M. Fardos (2015). Estimates of Population Parameters of Some Economic Traits in Holstein Friesian Cows by Using Statistical Program. *Global Veterinaria* 14: 129-135.
- Roy , R. Tomer, S. P.; Monglik, V. P. Sharma, R. J. and Chaudlary, R. P. (1985). Factors affecting reproductive and productive traits in different crossbred dairy cattle. *Indian J. Anim. Sci.*, 55: 385- 386.
- Salem, A.Y. and E. M. Abdel-Raouf, (1999). Genetic study of fertility and milk productive traits in a commercial herd of Holstein-Friesian cattle in Egypt. *J. Agric. Res., Tanta Univ.*, 25: 204.
- Salem, M. A; H. M. Esmail, R. R. Sadek and A.A. Nigm (2006). Phenotypic and genetic parameters of milk production and reproductive performance of Holstein cattle under the intensive production system in Egypt. *Egyptian J. Anim. Prod.*, (2006) 43(1):1-10
- Sattar, A. R. H; A. A. K Mirza; Niazi and M. Latif (2005). Productive and reproductive performance of Holstein Friesian cows in Pakistan. *Pakistan Vet. J.*, 25: 75-81.
- Shalaby, N. A., E. Z. M. Oudah and M. Abdel-Momin (2001). Genetic analysis of some productive and reproductive traits and sire evaluation in imported and locally born Friesian cattle raised in Egypt. *Pakistan J. Biol. Sci.*, 4: 893-901.
- Shalaby, N. A; A. S. A. El-Barbary; E. Z. M. Oudah and M. Helmy (2013). Genetic analysis of some productive and reproductive traits of first lactation of Friesian cattle raised in Egypt. *J. Animal and Poultry Prod., Mansoura Univ.*, 4: 97-106.
- Sharma, L. D. (1982). Reproductive performance of Nagaur cattle. *Indian. J. Animal Sci.*, 53: 1019-1020.

- Shitta, A.A., M.A. Tag El-Dein and Set El-Habaeib S. Awad (2002). A study on productive and reproductive traits of Friesian cattle in Egypt. J. Agric. Sci. Mansoura Univ., 27: 7281.
- Tadesse, M. J, J. Thiengtham, A. Pinyopummin. and S. Prasanpanich., (2010). Productive and reproductive performance of Holstein Friesian dairy cows in Ethiopia. Livestock Research for Rural Development, 22 (2) 2010 cited in <http://www.lrrd.org/lrrd22/2/tade22034.htm>.

## تقدير المعايير الوراثية لبعض الصفات الإنتاجية والتناسلية باستخدام نموذج الحيوان فى قطيع فريزيان تجارى فى مصر

حسن غازى العوضى<sup>١</sup>، عبد الخالق السيد عبد الخالق<sup>٢</sup> و محمد عطية أبو الريش<sup>٢</sup>

١- قسم الإنتاج الحيوانى - كلية الزراعة - جامعة كفر الشيخ - مصر.

٢- قسم الإنتاج الحيوانى - كلية الزراعة - جامعة المنصورة - مصر

الدراسة اجريت فى قطيع تجارى تابع لجمعية الثروة الحيوانية بشبرياتانا- طنطا - محافظة الغربية على ٤٣٧٠ سجل لعدد ٨٧٤ بقرة فريزيان أبناء ٤٤ طلوقة خلال الفترة من ١٩٨٥ وحتى ٢٠٠٣م لدراسة واستخدام نموذج الحيوان لتقدير التباين والتباين المشترك لكلا من كمية اللبن فى ٣٠٥ يوم وطول موسم الحليب و طول فترة الجفاف وطول فترة التلقيح وطول الفترة بين ولادتين واشتمل نموذج التحليل الاحصائى على شهر وسنة الولادة ورقم موسم الحليب كعوامل ثابتة. واما التأثيرات العشوائية فاشتملت على الحيوان والتاثير الوراثى المباشر والامى والبيئى الدائم والتاثير المتبقى. كانت المتوسطات العامة لكلا من كمية اللبن فى ٣٠٥ يوم و طول موسم الحليب و طول فترة الجفاف و طول فترة التلقيح وطول الفترة بين ولادتين هى ٤٨٩٧ كجم و ٣٢٧ يوم و ١٢٠ يوم و ٦٦.٣ يوم و ٤٠١.١٦ يوم على التوالي. كان تاثير شهر وسنة الولادة على المعنوية (١%) على جميع الصفات المدروسة. كانت قيم المكافئ الوراثى المباشر للصفات سالفة الذكر هى ٠.٣٨ و ٠.١٣ و ٠.١٤ و ٠.١١ و ٠.١٦ على التوالي. بينما كتنت تقديرات المكافئ الوراثى الاموى هى ٠.١٣ و ٠.١٨ و ٠.٢٣ و ٠.١٤ و ٠.٢١ لنفس الصفات المدروسة على التوالي. كان الارتباط الوراثى المباشر والامى يتراوح بين -٠.٦٠ الى ٠.٧٤. وكانت الارتباطات الوراثية بين مختلف الصفات المدروسة تتراوح بين (-٠.٨٠ الى ١.٠٠) وبينما كانت الارتباطات المظهرية بين الصفات تحت الدراسة موجبة وتتراوح بين (٠.٠٧ الى ٠.٩٩). كان مدى القيم التربوية المقدر للطلانق هى ٤٦٢ كجم و ١.٠٥ يوم و ٠.٢٤ يوم و ٥.٩٣ يوم و ٢.٢٦ يوم لكمية اللبن فى ٣٠٥ يوم وطول موسم الحليب وطول فترة الجفاف وطول فترة التلقيح و طول الفترة بين ولادتين على التوالي. بالمقارنة مدى تقديرات القيم التربوية للامهات لنفس الصفات السابقة هى ٥١٠ كجم و ١.٤٢ يوم و ٠.٨٠ يوم و ٧.٢٠ يوم و ٣ يوم على التوالي. وكانت دقة التقديرات للقيمة التربوية تتراوح بين (٣٨ الى ٨٨ و ٧٣ الى ٨٨ و ٣٩ الى ٨٩ %) لكلا من الطلائق والابقار والامهات على التوالي وهذا يوضح إمكانية التحسين الوراثى من خلال اى منهما. كانت تقديرات التباين البيئى المؤقت كجزء من التباين المظهرى الكلى لكلا من طول موسم الحليب وطول فترة الجفاف وطول فترة التلقيح وطول الفترة بين ولادتين تتروح بين ٣٥ الى ٧١% وهذا بين إمكانية التحسين من خلال تحسين العوامل البيئية الموقفة.