ASSESSMENT OF LAND DEGRADATION IN WADI EL NATRUN AREA, WESTERN DESERT, EGYPT

Aqrawi, P. M.^{*}; A. A. Abdel Hady^{*}; W.A. Abde IKawy^{*} and A. El-Nahry^{**}

* Soils Science Dept., Faculty of Agriculture, Cairo Uni., Giza, Egypt.

** National Authority for Remote Sensing and Space Sciences

psaqrawi@y ahoo.com

ABSTRACT

Most forms of land degradation are human resource (mismanagement and misuse); some physical and chemical environmental factors are still considered. Quantitative assessment of land degradation and monitoring the changes in land qualities in Wadi El-Natrun are the main objective of this study. Physiographic map of the area was produced by using ETM+, ENVI 5.0 and ArcGIS10. Physiographic map used to determine soil profiles location and soil samples. From the physical and chemical analysis the results compared with the data extracted from Mohamed, (2011). Land degradation rate, relative extent, degree, and severity level in the study area were assessed. The results indicate that the dominant active land degradation features are; water logging, salinity, alkalinity and compaction. Based on the FAO/UNEP, (1979) program of degradation (rate, relative extent, degree, and severity) and the application on that data of water logged, salinity, alkalinity and compaction compared with Mohamed, (2011). The results indicate that the following: there is no effect of compaction so there is no compaction degradation . Waterlogged degradation as water table depth changed from 2011 to 2013 as following (50-100 cm) the degraded area increased from $(234.32 \text{ to } 341.28 \text{ km}^2)$,

(100-150 cm) the soil improved and the degraded area decreased (from 356.31 to 218.42 km²) and (> 150 cm) were improved. Salinity degradation as electrical conductivity(EC dS/m) data changed from (2011 to 2013) that the (8-16 dS/m) the degraded area increased from (0.0 to 16.83 km²), (4-8 dS/m) the soil improved and the degraded area decreased (from 190.86 to 110.43 km²) and (<4 dS/m) the soil improved and the area increased (from 399.76 to 451.87). Alkalinity degradation data changed from (2011 to 2013) that the (>15 %) no change , (10-15%) the soil improved and the degraded area decreased (from 437.98 to 190.82 km²) and (<10 %) the soil improved and the area increased (152.64 to 388.32 km²)

Keywords: Physiographic map, land degradation, compaction, salinity, alkalinity.

INTRODUCT ION

Wadi El-Natrun area lies to the west of Nile Delta and it is considered the natural extension of Nile Delta. The studied area lies between longitudes 30° 06′ 21″.37 to 30° 28′ 50″.02 East and latitudes 30° 18′ 02″.88 to 30° 31′ 06″.66 North. It covers an area of about 142,687 fed. (Fig. 1).

The climatic conditions of Wadi El-Natrun area are those characterizing the desert areas of Egypt. It is characterized by a hot rainless summer.(Fig1) The maximum temperature (34.5 \degree C) was recorded in July and August, while the minimum one (7.5 \degree C) was recorded in January with an average of 13.4 \degree C and 27.9 \degree C for the mean minimum and maximum annual temperature, respectively. The precipitation is rare and recorded only during

November, December, January, February, March and April. The highest value 4.9 mm was recorded in January and the lowest one 0.8 mm was recorded in April. Daily evaporation is high.

Map1

Fig1

The lowest value of evaporation (1.8 mm/day) was recorded during January, while the highest value (7.9 mm /day) was recorded in June. After Meteorological Authority, 2006 Shata and El Fayoumy, (1970) classified the geological features of the area located between Nile Delta and Wadi El-Natrun as follows:

1. The sediments belong to the Pliocene are distributed on a large scale beneath Wadi El-Natrun area and developed into marine and fresh water faces. In the vicinity of Wadi El Natrun, the Pliocene section is thick and distinct into two main portions mostly showing slight variations among themselves, such as a lower portion composed of green sandy clays and the upper one built up by calcareous grits. Brown "Oolite" calcareous silt and sand clayey lime soil with iron oxides are associated with the Pliocene formation in Wadi El-Natrun Depression.

2. The Pleistocene and Holocene deposits are located east of Wadi El-Natrun and to the west of the Nile Delta. These sediments have distribution in the studied area and essentially developed into gravel and sand faces.

Said, (1990) stated that the western desert of Egypt stretches westwards from the Nile Valley to the borders of Libya and occupies an area (exclusive of EI-Fayoum, and some natural depressions) of about 681000 km². It is essentially a plateau with vast expanses of rocky ground and numerous extensive and deep closed depressions. It attains its greatest altitude in the extreme southwestern corner of Egypt. It is disturbed by the great mountain mass of Gabel Oweinat lying just outside Egypt; with only the northern flames of the mountain are within the borders of Egypt. Northeast ward from Gabel Oweinat a broad tract of high ground extends for more than 200 km. This is the extensive sandstone palteau of Gilf El-Kebir, nearly 1000 m above sea level. On the other side of this tract and to its north the ground slopes gradually to the depressions in which the Oases of Abu Mungar, EL-Dakhla and El-Kharga are situated. Immediately north these Oases a high and much embayed escarpment, which is the southern edge of a great plateau of Eocene limestone. This plateau rises in places over 500 m above sea level. In this limestone plateau the great hollows containing the Oases of EI-Farafra and EI-Baharia are situated.

Soil texture of the investigated area is sandy with very low percentage of clay and silt, where sand percentage ranged between 90.71 to 98.52%. The gravel percentage of the surface soil is relatively high as it gravel percentage reached 34%, Abdel-Hamid ,(2008). Soils salinity are slightly in cultivated soils and moderately to strongly in virgin area ranged between 0.64 to 244 dS/m (Ashmawy, 2003) and (Yehia, 2004). Soil reaction (pH values) varied between 8.2 and 8.9 (Erian, (2000), Bahnassy et al,. (2001), and Ashmawy, 2003).Organic matter content ranges between 0.3 -0.8 % (Erian, 2000 and Ashmawy, 2003). calcium carbonate content ranged between 9 - 17% (Ashmawy, 2003).

Hefny (1993) stated that the aquifer of Wadi El-Natrun area, Wadi El-Natrun aquifer and Moghra aquifer. The upper one is Wadi El-Natrun Aquifer, which is local and of low production. It is multi-layered, with alternation of sands and clay, which belong to pliocene. The lower aquifer is the Moghra aquifer which consists of sands and gravels of the Moghra formation. Oligocene basalt or oligocene shale underly the aquifer.

Attia, (1975) The most natural vegetation, which exist in the studied area is annual weeds that flourish during the rainy season. The soil surface is barren, with regard to vegetation, which only small patches (depressions) are covered with natural vegetation, typical of the Western Desert of Egypt as: Artemisia monosperma, Pityranthus tostuesus, Aristida pliniosa, and other common perennial species.

Geographical Information Systems (GIS) have proved to be immensely helpful in the organization of the huge database generated through space technology (Trotter, 1991). The utility of GIS in the analysis and modeling of integrated information is well established (Burrough, 1986).

GIS has been used in the development of digital databases, assessment of status and trends of resources utilization of the areas and to support and assess various resource management alternatives (Clark, 1991). Spectacular developments in the field of GIS to synthesize various thematic information with collateral data have not only made this technology effective and economical but also a tool to arrive at development strategies for sustainable land and water resources management.

Farmers can order spectral imagery of their fields to determine the status of their land and whatever is growing on it. For example, spectral imagery can indicate the amount of fertilization required in specific locations that are designated with GPS coordinates. Agricultural machinery on the market today has the capability to load this information into computers built into the machinery and automatically adjust the amount of fertilizer deposited based on the information contained in the spectral imagery. The type of vegetation can also be determined from spectral remote sensing. Clark (1991) shows the types of plants growing in a field. Because stressed vegetation looks different from healthy vegetation, mapped remote sensing information can be an indication of plant disease or drought.

Soil degradation is defined as the process which lowers (quantitatively or qualitatively) the current and/or the potential capability of s oil to produce goods or services. Soil degradation implies a regression in capability from a higher to lower state; a deterioration in s oil productivity and land capability, Mashali, (1991), Ayoub, (1991), UNEP Staff, (1992), Wim, and El Hadji., (2002). The food gap due to increasing population puts more pressure on the us e of land, resulting in serious forms of land degradation. These are considered irreversible processes particularly with the severe and continued misuse and poor management. The intensification of agriculture coupled with poor management accelerates the rate of land degradation. Food supply situation will be worse in the future if the current trend of land degradation does not change drastically. The livelihoods of more than 900 million people in some 100 countries are now directly and adversely affected by land degradation United Nations, (1994) Unless the current rate of land degradation is s lowed and reversed, food security of humanity will be threatened and the ability of poor nations to increase their wealth through improved productivity will be impeded. Land degradation can be observed in all agro climatic regions on all continents . Although climatic conditions , such as drought and floods, contribute to degradation, the main causes are human activities. Land degradation is a local problem in vas t number of locations, but it has cumulative effects at regional and global scales. The countries of the developing world, and particularly those in the arid and semi- arid zones, are the most seriously affected UNEP Staff, (1986). The status of s oil degradation is an expression of the severity of the process. The severity of the processes is characterized by the degree in which the s oil is degraded and by the relative extent of the degraded area within a delineated physiographic unit UNEP Staff, (1991).

The main objectives of this investigation was to study the quantitative

assessment of land degradation and monitoring the changes in land qualities in wadi El-Natrun .

MATERIALS AND METHODS

The Physiographic Map extracted by using ETM+ taken during the year 2005 (Fig. 2).



(Fig. 2): ETM+ image of the studied area ENVI 5.0 and ArcGIS 10.1 software has been us e d to produce the physiographic map of the studied area (Map.2 and table 1)

Map2



physiograp units	land form	Area Km2	Area%
Torroio	High aeolian	11.01	1.25
Terrais	Low aeolian	16.83	1.92
Depentation basin	High decantation	242.69	27.67
Decantation basin	Low decantation	36.69	4.18
Overflow basin	High overflow	19.79	2.26
	Low overflow	35.25	4.02
Sand choot	High	305.12	34.79
Sand Sheet	Low	196.37	22.39
Sapkhas		6.15	0.70
Water body		7.19	0.82
		877.08	100.00

Table	(1)	ph	ysio	graphi	c legen	d of	studied	area
-------	-----	----	------	--------	---------	------	---------	------

Soil analysis

The studied area representative by 8 different mapping units as shown in legend. 16 soil profiles were represented the mapping units and 60 minpits taken to corrected the boundaries of mapping units , the soil profiles were dug and descripted according to FAO guidelines **FAO**, **(2010)**. The soil samples collected to preparing for laboratory analyses . physical analyses(texture class, bulk density gm/cm³) and chemical analyses as (pH reaction , O.M % , calcium carbonate % , EC dS/m , CEC meq/L , ESP%) using the s oil survey laboratory methods manual USDA, (2004) . The American Soil taxonomy USDA ,(2010) was used to classify the s oil of the studied area.

Soil degradation assessment :

Based on the comparing between the data extracted from Mohamed, 2011 and the data resulting from this study. The FAO/UNEP,(1979) methodology for assessing soil degradation was used and the results were evaluated and confirmed with the physiographic units, the ratings used are presented in Tables (2 and 3).

Chemical degradation	Salinization (Cs) increase in (EC) per dS/m/year	Alkalinization (Ca) increase in ESP/Year
Non to slight	<0.5	<0.5
Moderate	0.5 – 3	0.5 – 3
High	3 – 5	3 – 7
Very high	>5	>7
	Compaction/increase in bulk	Water
Physical degradation	density per g/cm³/year	logging/increase in water table incm/year
Physical degradation Non to slight	density per g/cm³/year <0.1	logging/increase in water table incm/year <1
Physical degradation Non to slight Moderat e	density per g/cm ³ /year <0.1 0.1 - 0.2	logging/increase in water table incm/year <1 1 - 3
Physical degradation Non to slight Moderat e High	density per g/cm ³ /year <0.1 0.1 - 0.2 0.2 - 0.3	logging/increase in water table incm/year <1 1 - 3 3 - 5

Table(2) Soil degradation types, classes and rates

				Haza	ard class	
Hazard type	Indicator	Unit	Low	Moderate	High	Very high
Salinization	EC	dS/m	4	4 – 8	8 – 16	>16
Alkalinization	ESP	value	10	10 – 15	15 – 30	>30
Compaction	Bulk density	g/Cm ³	1.2	1.2 – 1.4	1.4 – 1.6	>1.6
Water Logging	Water Table level	cm	150	150 – 100	100-50	<50

Table(3) Criteria used to determine the degree of the different degradation types

Land degradation degree, relative extent, severity level and causative factors were defined and described using the UNEP, (UNEP Staff, 1991) approach. The relative extent of each type of soil degradation within the mapped unit is recognized as :The s oil degradation severity level is indicated by the combination of the degree and the relative extent as shown in (Table 4&5).

unit

Table (4): The severity level of soil degradation

Category	% of the mapping
1. Infrequent	up to 5%
2. Common	6-10%
3. Frequent	11-25%
4. Very frequent	26-50%
5. Dominant	over 50%

Table(5): Relative extant (%) of degradation soil

Degree of soil	Relative extent (%)				
degradati on	0-5	6-11	11-25	26-50	50-100
S light	1.1	1.2	1.3	1.4	1.5
Moderate	2.1	2.2	2.3	2.4	2.5
Strong	3.1	3.2	3.3	3.4	3.5
Extreme	4.1	4.2	4.3	4.4	4.5

Low Moderat e High Very high

RESULTS AND DISCUSSION

Morphological features of the studied area :-

16 soil profiles representative the main physiographic units of the studied area were descripted as show in Table (6) and samples collected for laboratory analyses as the following :-

The physical and chemical analysis of the studied area were shown in Table (7)

The variation in electrical conductivity between the studied (2010 and 2013)shown in (Map.3)



(Map.3) electrical conductivity 2010 to 2013

The variation in the Alkalinity between the studied (2010 and 2013) (Map.4)



(Map.4) Alkalinity 2010 to 2013 The variation in the waterlogging between the studied (2010 and 2013) (Map.5)



Τ6

t7



(Map.5) Waterlogging 2010 to 2013 Land degradation of the soil :-

Based on UNEP Staff, (1991) the soil degradation calculated as the following :-

Table (8) B:the fina	I Land degradation	on rates from	compared	between t	he
studied ((2010-2013):				

Profile No.	e Mapping unit	Water logging	Compaction	Salinity	Alkalinity
1	LSS	1	1	1	1
2	LDB	1	1	1	1
3	HAT	1	1	1	1
4	HDB	1	1	1	1
5	HSS	1	1	1	1
6	LOB	1	1	1	1
7	LAT	1	1	1	1
8	HOB	1	1	1	1
1-1.0%	2- Modorato	2_ ∐iah			

1= Low 2= Moderate 3= High

From Table (8) clear that the compared between studied of the area 2010 and 2013 there is no differentiation in the data of water logged , compaction , salinity and alkalinity and the rates of the degradation is low

Table	(9)	The	main	causative	factors	of	human	induced	land
degradation types in the studied area.									

Profile No.	Mapping unit	Water logging	Compaction	Salinity	Alkalinity
1	LSS	i/o	0	0	0
2	LDB	i/o	0	0	0
3	HAT	i/o	0	Mi	0
4	HDB	i/d/o	0	0	0
5	HSS	i/o	0	0	0
6	LOB	i/o	0	Mi	0
7	LAT	i/o	Ő	Ō	Ó
8	HOB	i/o	0	Mi	0

i: over irrigation, mi: Poor management of irrigation scheme,

m: improperly timed used of heavy machinery. **d**: human intervention in natural drainage **o**: other activities which include (shorting of the follow periods and the absence of conservation measurements from the Table (9) clear that the over irrigation and other activities were the first factors caused the degradation , poor management of irrigation scheme were the second factors of soil degradation and human intervention in natural drainage were the third factors of causative land degradation .

Table (10): Land degradation status in the different mapping unit of the studied area:

Mapping unit	Land degradation status
LSS	(Pw i/ o 1,4) & (Pc m/o 1,5) & (Cs mi 1,5) (Ca mi 1,5)
LDB	(Pw i/ o 1,3) & (Pc m/o 1,5) & (Cs mi 1,5) (Ca mi 1,2)
HAT	(Pw i/ o 1,4) & (Pc m/o 1,5) & (Cs mi 2,5) (Ca mi 1,5)
HDB	(Pw i/ o 2,3) & (Pc m/o 1,5) & (Cs mi 1,2) (Ca mi 1,2)
HSS	(Pw i/ o 1,4) & (Pc m/o 1,5) & (Cs mi/o 1,4) (Ca mi 1,5)
LOB	(Pw i/ o 1,3) & (Pc m/o 1,5) & (Cs mi 1,4) (Ca mi 1,2)
LAT	(Pw i/ o 1,4) & (Pc m/o 1,5) & (Cs mi 1,5) (Ca mi 1,5)
HOB	(Pw i/ o 2,3) & (Pc m/o 1,5) & (Cs mi 1,2) (Ca mi 1,2)

two letters= causative factors as,

The first two letters = degradation types as, $Pw \rightarrow physical degradation/ water logging.$ intervention in natural drainage. $Pc \rightarrow physical degradation/ soil compaction.$ use of heavy machinery. $Cs \rightarrow chemical degradation/ Salinity$

management of irrigation scheme.

Ca \rightarrow chemical degradation/ alkalinity

The following one or

 $I \rightarrow \text{over irrigation}$ $d \rightarrow \text{human}$

 $m \rightarrow$ improperly time

mi→ poor

 $o \rightarrow$ other activities

From the Table (10) the dominant land degradation states in whole area caused by physical (water logging and compaction) and chemical (salinity and alkalinity) with some causative factors (over irrigation, heavy machinery, poor management of irrigation scheme, over activities), but the area units different in severity level

From degraded studied area clear that there is no effect of compaction so there is no compaction. Waterlogged degradation as water table depth changed from 2011 to 2013 as following (50-100 cm) the degraded area increased from (234.32 to 341.28 km²), (100-150 cm) the soil improved and the degraded area decreased ($356.31 \text{ to } 218.42 \text{ km}^2$) and (> 150 cm) were improved. Saliniity degradation as electrical conductivity(EC dS/m) data changed from (2011 to 2013) that the (8-16 dS/m) the degraded area increased from ($0.0 \text{ to } 16.83 \text{ km}^2$), (4-8 dS/m) the soil improved and the degraded area decreased (399.76 to 451.87). Alkalinity degradation data changed from (2011 to 2013) that the (>150 cm) the soil improved and the area increased (399.76 to 451.87). Alkalinity degradation data changed from (2011 to 2013) that the (>15 %) no change, (10 - 15%) the soil improved and the degraded area decreased ($437.98 \text{ to } 190.82 \text{ km}^2$) and (<10 %) the soil improved and the area increased ($152.64 \text{ to } 388.32 \text{ km}^2$)

CONCLUSION

Wadi El-Natrun area lies to the west of Nile Delta and it is considered the natural extension of Nile Delta. The studied area lies between longitudes 30° 06' 21".37 to 30° 28' 50".02 East and latitudes 30° 18' 02".88 to 30° 31' 06".66 North. T he total area of Wadi El Natrun is about 281.7 K m 2 (i.e. 67608 feddans), extended in a NW-SE direction and 23 m below sea level. This area has always been confine d as a possible area for reclamation and utilization due to its location and the presence of ground water in a suitable quality for irrigation. T he origin of the underground water in W adi El-Natroun is seepage from the Nile stream, due to its proximity and low level (E I M aghraby, 199 0). Most forms of land degradation are human resource (mismanagement and misuse); some physical and chemical environmental factors are still considered. Quantitative assessment of land degradation and monitoring the changes in land qualities in Wadi El-Natrun are the main objective of this study . physiographic map of the area produced by using ETM+, ENVI 5.0 and ArcGIS10 . physiographic map used to determined soil profiles location and soil samples. From the physical and chemical analysis the results compared with the data extracted from Mohamed (2011). Land degradation rate, relative extent, degree, and severity level in the study area were assessed. The results indicate that the dominant active land degradation features are; water logging salinity, alkalinity and compaction. Based on the program FAO/UNEP(1979). of degradation (rate, relative extent, degree, and severity) and the application on that data of water logged salinity alkalinity and compaction compared with Mohamed (2011). clear that the following :- there is no effect of compaction so there is no compaction. Waterlogged degradation as water table depth changed from 2011 to 2013 as following (50-100 cm) the degraded area increased from (234.32 to 341.28 km²), (100-150 cm) the soil improved and the degraded area decreased (356.31 to 218.42 km²) and (> 150 cm) were improved. Salinity degradation as electrical conductivity(EC dS/m) data changed from (2011 to 2013) that the ($8\mathchar`-16\mbox{ dS/m}$) the degraded area increased from (0.0 to 16.83 km²), (4-8 dS/m) the soil improved and the degraded area decreased (190.86 to 110.43 km²) and (<4 dS/m) the soil improved and the area increased (399.76 to 451.87). Alkalinity degradation data changed from (2011 to 2013) that the (>15%) no change, (10-15%) the soil improved and the degraded area decreased (437.98 to 190.82 km²) and (<10 %) the soil improved and the area increased (152.64 to 388.32 km²)

REFERENCES

- Abdel-Hamid, A. A. (2008). Integration of New Technology Systems for Soil Reclamation and Evaluation at El-Gahar Area, Wadi El-Natrun, Egypt. M.Sc. Thesis, Fac. Agric., Alexandria. Univ., Egypt, 108 p.
- Ashmawy, S. A. (2003). Pedological Studies Bearing on Gensis, Morphological and Classification of Soils of Wadi El-Natrun Depression. Ph.D. Thesis, Fac. Agric., Moshtohor, Benha branch, Zagazig Univ., Egypt, 176 p.
- Attia, S.H. (1975). Petrology and Soil Genesis of the Quaternary Deposits in the Region West of the Nile Delta (North and East of Wadi El Natrun). Ph.D. Thesis, Fac. Sci., Ain Shams Univ., Cario, Egypt, 358 p. Cited by Abu Sleem (2010).
- Ayoub, A. T., 1991. An assessment of human induced s oil degradation in Africa. U.N. environmental program, Second Soil Sci. conf. Cairo Egypt.
- Bahnassy, M.; Ramadan, H.M.; Abdel-Kader, F. and Yehia, H.M. (2001). Coupling GIS with modeling tools to support land use planning and management of Sugar Beet area, west Nubaria, Egypt. Alex. J. Agric. Res., 46: 169-180.
- Burrough, P. A. 1986. Principles of Geographical Information Systems for Land Resources Assess-ment, Monographs on Soil and Resources Survey No. 12, Oxford Science Publica-tions, Oxford, England: Clarendon Press.
 - Clarck M.D.(1991) Global database and their implications for GIS. In: Maguire DJ , Goodchild MF, Rhind DW (eds). Geographical information systems : principles and applications. Longman, London , pp. 217-31 Vol 2.
- El- Maghraby, M.M., 1990. "Geographical and hydorological studies of Sadat City, Egypt" M.Sc.Thesis, Fac. Sci., Alexandria University.
- Erian, W. F.; Gomaa, F. A. and Ismail, S. A. (2000). Towards sustainable agriculture on central and marginal slopes of Wadi El-Natrun depression, Egypt. Bull. Fac. Agric. Alexandria. Univ., Egypt, 2: 71-82.
- FAO/UNEP, 1979. A Provis ional methodology for degradation as s es s ment. Bul. No. 48. ,FAO, Rome, Italy
- FAO, 2010. Guidelines of Land Evaluation for Rainfed Agriculture. FAO Soils Bull., No. 52, Rome, Italy, 66 p
- Hefny, K. (1993). The role of groundwater in the sustainable development planning of the region of Wadi El-Natrun. The Research Institute for groundwater, Bull. Fac. Agric. Alexandria. Univ., Egypt, 7: 79-91
- Mas hali A. M., 1991. Land degradation and desertification in Africa 2nd African Soil Sci. Soc. Conf.
- Meteorological Authority (2006). The normals for Egypt up to 2007. Ministry of Civil Aviation, Cairo, Egypt. Meteorological Res. Bull., 22:86-87

- Mohamed, Y. K. 2011. The use of remote sensing and GIS Techniques For assessment of soils of Wadi El-Natrun, Egypt. Ph. D. Thesis, Fac. Agric., Cairo Univ., Egypt .
- Said, R. (1990). The Geology of Egypt. Elservier Publishing Co., Amsterdam, New York, USA, 734 p.
- Shata, A.A. and El Fayoumy, I.F. (1970). Remarks on the regional geological structure of the Nile Delta. Proceedings of the Bucharest Symposium on Delta, 6-14 May 1969. Geol Soc. J., 41: 99-127.
- Trotter CM (1991), Remotely sensed data as an information source for geographical information systems in natural resource management: a review. Int. J. of GIS, Vol. 5 No. 2 pp 225-39
- Wim, G. and El Hadji, M., 2002. Causes, general extent and physical consequence of land degradation in arid, s emiarid and dry sub humid areas. Forest conservation and natural resources, forest dept. FAO, Rome, Italy.
- UNEP Staff, 1986. Sands of change: Why land becomes des ert and what can be done about it. UNEP Environmental Brief No 2, United Nations Environment Program, Nairobi, Kenya.
- UNEP, 1991, Status of Desertification and Implementation of the United Nations Plan of Action to Combat Desertification. UNEP, Nairobi, Kenya, 77 pp.
- UNEP Staff,1992. World atlas of decertification. Publ. E. Arnold, London, 69 pp.USDA, 2004. Soil Survey Laboratory Methods Manual. Soil Survey Investigation Report No. 42 Vers ion 4.0 November 2004
- United Nations, 1994. Earth Summit Convention on Des ertification. Proceedings of the United Nations. Conference on Environment and Development (UNCED), Rio De Janeiro, Brazil, 3-14 June 1992. Department of Public Information, United Nations, New York, USA.
- USDA, 2010. Soil Survey Laboratory Methods Manual. Soil Survey Investigation Report No. 42 Vers ion 4.0November 2004.
- Yehia, H.A.M. (2004). Land Resource Assessment for Sustainable Agriculture Development at Multi Spatial Scale; A case study for El-Behira and Wadi El-Natrun district, Egypt. Ph. D. Thesis, Fac. Agric., Alexandria Univ., Egypt, 250 p.

تقييم تدهور الأراضي بمنطقه وادي النطرون , الصحراء الغربيه , مصر عقراوى بيرس مجيد* ، عبد الحميد على* ، عبد القوى وائل* و النهرى علاء ** * قسم الاراضي / كلية الزراعة / جامعة القاهرة

** الهيئة العامة للاستشعار عن بعد وعلوم الفضاء

معظم أشكال تدهور الأراضي هي الموارد البشرية (سوء الإدارة و سوء الاستخدام) ؛ تعتبر بعض العوامل البيئية الفيزيائية والكيميائية هي الاساس في هذا التدهور . التقييم الكمي لتدهور الأراضي ورصد التغيرات في صفات الأراضي في وادي النطرون هي الهدف الرئيسي من هذه الدر اسة .

تم إنتاج خريطة جغرافية للمنطقة باستخدام صور القمر الصناعي (+ ETM)،وانتاجها باستخدام برامج ENVI 5.0 وArcGIS10

خريطة جغرافية طبيعية استخدمت لتحديد ملامح التربة الموقع و عينات من التربة . من التحاليل الفيزيائية والكيميائيوة مقارنة النتائج مع البيانات المستخرجة من محمد ، (٢٠١١) . تم تقييم معدل تدهور الأراضي ، مدى النسبية ، ودرجة التدهور ، و مستوى خطورة التدهور في منطقة الدراسة . وتشير النتائج إلى أن تدهور الأراضي بالموقع والتي تم الاعتماد عليها لاستخراج النتائج هي

(التشَّبع والملوحة و القلوية و التضاغط)، على أساس برنامج منظمة الأغذية والزراعة / برنامج الأمم المتحدة ، (١٩٧٩) للتدهور (معدل التدهور ومدى النسبية ، ودرجة الدهور ، وشدة التدهور) و تطبيقها على تلك البيانات (التشبع ، والملوحة ، والقلوية و التضاغط) مقارنة مع محمد ، . (7 • 1 1)

وتشير النتائج إلى أن ما يلي : - ليس هناك أي تأثير ملموس في تضاغط التربة، اما التدهور بالنسبة للتشبع بالمياه من حيث عمق منسوب المياه الجوفية فقد تغيرت من ٢٠١١ الي ٢٠١٣ على النحو التالي (٥٠-١٠٠ سم) زادت المساحة ا

لمتدهورة من (٢٣٤.٣٢ الى ٣٤١.٢٨ كم) ، (100-150) (سم) تحسنت و انخفضت المنطقة المتدهورة (٣١٦.٣٥الي٢١٨.٤٢ كم) وكذلك تحسنت المناطق ذات الاعماق (> ١٥٠ سم). بالنسبة للتدهورُ بالملوحة و الموصلية الكهربائية / EC DS) م) فان البيانات التي تغييرت من (۲۰۱۲-۲۰۱۱) أن (dS/m ا6-8) فان المساحة المتدهورة ارتفعت من (0.0الي 16.83 كم)،(4-8 dS/m) فان التربة تحسنت وانخفاض المساحة المتدهورة (190.86 الى 110.43 كم٢) وكذلك (dS/m >>) قد تحسنت التربة و مساحة التحسن (399.76 الى 451.87 كم ً) . بالنسبة لبيانات التدهور على اساس القلوية فقد طرأ تغيير على المنطقة خلال الفترة (٢٠١١-٢٠١٣) أن (> ١٠٪) لا تغيير ، (١٠-١٠ ٪) تحسين التربة وانخفاض منطقة المتدهورة (437.98 الى 190.82 كم) و (<١٠ ٪) قد تحسنت التربة و زادت المساحة (152.64 الى 388.32 كم ً) .

قام بتحكيم البحث

كلية الزراعة – جامعة المنصورة	<u>اً د</u> / ایمن محمد الغمری
مركز البحوث الزراعية	<u>اً د</u> / رأفت رمضان على