

## Serological Diagnosis Of Camel Brucellosis At Sharkia Governorate, Egypt

Eisa MI, Monazie AM, \*\*Khoudair RM and El-Shymaa AA

\* Dept. Animal Med., Fac. Vet. Med., Zagazig Univ., Egypt

\*\* Brucellosis Research Dept., Animal Health Research Institute, Dokki, Giza, Egypt

\*\*\*Vet. Clinic., Fac. Vet. Med., Zagazig Univ., Egypt

### ABSTRACT

The present study was carried out on 210 camels serum samples collected from a various abattoirs at Sharkia Governorate to clarify the prevalence of the brucellosis among camels during the period from June 2012 to August 2013 in addition to find the most reliable diagnostic method. In this investigation, serological tests on 210 camel sera using Buffered Acidified Plate Antigen Test (BAPAT), Rose Bengal Plate Test (RBPT), Rivanol Test (Riv.T), Complement Fixation Test (CFT) and Immunochromatographic Assay (ICA) to evaluate traditional and recent tests which used for the diagnosis.

The obtained results were 7.6%, 6.7%, 6.2%, 6.2% and 6.7% using BAPAT, RBPT, Riv.T, CFT and ICA respectively. The sensitivity percent of different serological tests were 95.45%, 87.75%, 90.90%, 88.85% and 92.15% for BAPAT, RBPT, Riv. T., CFT and ICA respectively. While their specificity percent were 99.27%, 99.61%, 99.80%, 99.80% and 99.75% for the same test respectively.

ICA could be recommended as a rapid screening test, easily performed, sensitive and high specific and as confirmatory test for diagnosis of brucellosis in camels. So, ICA the method of choice when testing animals in remote areas, nomadic and other migratory population. Thus study throw a spot light to include camels in the national program for control and eradication of brucellosis in Egypt on the base that where brucellosis exists in stock animals.

### INTRODUCTION

Animal brucellosis has been recorded in Egypt since 1939 and the prevalence of serological reactors on limited surveys has varied from one survey to another (1). The prevalence of the disease is related to the management practices of the farm and the ability of a country to finance prevention or control. Early detection, control and elimination of reactors are important consideration for the control of brucellosis (2). The prevalence of serological reactors among camels was 9%, 9% and 8.4% by RBPT, BAPAT and CFT respectively (3) while, it was 8.74%, 9.53% and 9.26%, by using RBPT, BAPAT and Riv.T respectively (4) but, it was 5% by using RBPT (5).

Classical serological tests are routinely used for the diagnosis of brucellosis. These tests (RBPT and BAPAT tests) deactivate IgM, which is responsible for non specific reactions. These tests are highly sensitive, but have low specificity and are ineffective in discriminating vaccinal antibodies from those by infection, but these tests still now are a rapid easily applied field tests (1).

As complicated assays may not available in many places, more recently, the convenience and speed of the test have been achieved by a novel concept of immunochromatographic assay (ICA) which is a simplified version of ELISA (6).

ICA assay does not require specific expertise, equipment or electricity, and test kits may be kept in stock without the need for refrigeration, thus, making the assay a very useful one for poor resource countries including most African Countries and migratory herds/flock (7). ICA test used for detection of antibodies against *Br. abortus* in sera samples, the ICA had 94.44% sensitivity and 100% specificity versus RBPT as a gold standard (8).

This study was formulated to study the prevalence of brucellosis in camels and diagnosis of brucellosis by using traditional and recent serological tests and evaluate these tests which used for the diagnosis.

## MATERIAL AND METHODS

**Animals:** A total of 210 adult mature and apparently healthy camels in different localities at Sharkia governorate during the period from June 2012 to August 2013. These camels were admitted for slaughter in different abattoirs.

**Serum samples:** Blood serum samples were collected from 210 investigated camels.

The serum stored at  $-20^{\circ}\text{C}$  in the deep freezer till used for different serological tests (9).

**Antigens for serological tests:** *Brucella abortus* antigen for BAPAT, RBPT and Riv.T were obtained kindly from the Veterinary Serum and Vaccine Research Institute (VSVRI), Abbassia, Cairo, Egypt.

*Brucella abortus* concentrate for CFT was kindly offered by the National Veterinary Service Laboratories (NVSL), Ames, USA.

**Immunochromatographic assay (ICA):** Brucella ICA manufactured according to (10). Product Feature (Kit component) obtained kindly from Brucellosis Research Department, Animal Health Research Institute, Dokki, Giza, Egypt.

**Serological tests:** Serum samples were tested by different serological tests. Buffered acidified plate antigen test (BAPAT), Rivanol test (Riv.T.) and Complement fixation tests were carried out according to (9), while, Rose Bengal plate test (RBPT) according to (11) and ICA procedures according to (12).

**Interpretation of Immunochromatographic Assay**

**Negative:** The presence of only one purple color band within the result window indicates a negative result .



**Fig. 1. Negative ICA**

**Positive:** The presence of two color bands ("T" band and "C" band) within the result window, no matter which band appears first, indicates a positive result.



**Fig. 2. Positive ICA**

**Invalid:** If the purple color band is not visible within the result window after performing the test, the result is considered invalid. The directions may not have been followed correctly or the test may have deteriorated. It is recommended that the specimen be re-tested.



**Fig. 3. Invalid ICA**


Determination of sensitivity% and specificity% of serological tests: Sensitivity and specificity% of different serological tests were determined according to (9).

**RESULTS**

Results of serological tests: The prevalence of brucellosis among camels was 7.6%, 6.7%, 6.2%, 6.2% and 6.7% by BAPAT, RBPT, Riv. T, CFT and ICA respectively as shown in Table (1) and Figure (4)

**Table 1. Prevalence of brucella reactors camels by different serological tests**

Total No.of examined camels	Test	BAPAT		RBPT		Riv. T		CFT		ICA	
		Positive		Positive		Positive		Positive		Positive	
		No.	%	No.	%	No.	%	No.	%	No.	%
210		16	7.6	14	6.7	13	6.2	13	6.2	14	6.7



**Fig. 4. Prevalence of brucella reactors among camels by different serological tests**

Results of sensitivity and specificity of different serological tests

The sensitivity of different serological tests was 95.45%, 87.75%, 90.90%, 88.85% and 92.15% for BAPAT, RBPT, Riv. T, CFT and ICA respectively. While the specificity of the same tests was 99.27%, 99.61%, 99.80%, 99.80% and 99.75% . It was found as shown in Table (2).

**Table 2. Results of sensitivity %and specificity % of different serological tests**

Test	Sensitivity %	Specificity %
BAPAT	95.45	99.27
RBPT	87.75	99.61
Riv. T	90.90	99.80
CFT	88.85	99.80
ICA	92.15	99.75

## DISCUSSION

In order to control and eradicate brucellosis from livestock animals it is very important to establish an appropriate serological method for diagnosis of brucellosis in the enzootic areas (13). In the present study, the prevalence of brucella reactors among camels by different serological tests as shown in Tables (1) and Figures (4) revealed that camels showed the highest percentage of brucella infection (7.6%) using BAPAT, while 6.7% by RBPT. In camels, the prevalence of *Brucella* recorded cases reached 7.9% (14), 11.37% (15), 10.7% (16) and 9.53% (4).

The results of CFT (6.2 %) revealed that the majority of positive CFT titers were 1/16 or higher (up to 1/256) indicating very active infection status where the rate of transmission of the disease among animals was high. This reflected some shortage in the disease control process as well as poor hygienic measures. Rivanol test in this study detected (6.2%) positive reactors. The results of the test showed detection of large number of positive reactors this due to precipitation of IgM and detection of IgG only this was in harmony to that obtained by (17,18).

Immunochromatographic assay (ICA) in this study detected (6.7%) positive reactors. The assay is based on the binding of specific antibodies to antigen immobilised on a test strip (cellulose membrane matrix). It allows the detection of specific IgM as well as specific IgG antibodies and that a high sensitivity is assured for all stages of the disease as reported by (19). The results also agree with (6,20) who recorded that the speed of ICA make it available for the rapid presumptive test which can replace RBPT in brucellosis control programs.

In this study, the sensitivity % and specificity % of different serological tests have been calculated. Table (2) illustrated that the sensitivity was 95.45%, 87.75%, 90.90%, 88.85% and 92.25% for BAPAT, RBPT, Riv. T., ICA and CFT respectively while the specificity was 99.27%, 99.61%, 99.80%, 99.80% and 99.75% for the same tests

respectively. The highest sensitivity in this study was that of BAPAT (95.45%). Complement fixation test showed the highest specificity (99.80%) followed by RBPT (99.61%). Such finding indicate that the suitability of BAPAT and RBPT for screening purposes. The obtained results agreed with those reported by (12) who reported that conventional agglutination tests have good sensitivity but their lack of specificity and occurrence of false positive serological results make a specific test necessary. The same results were obtained by (22) who stated that infected animals with brucellosis could be screened with a test such as RBPT or BAPAT and confirmed with ICA as a more specific test.

The sensitivity % of ICA (92.15%) and specificity% (99.75%) were higher than RBPT. Such findings indicate the suitability of the test to be used to replace RBPT in brucellosis control program advantages that make it the method of choice when testing animals in remote areas, nomadic and other migratory population. Practical advantages include that the use of the ICA does either requires specific training, expertise, electricity nor expensive equipment, that assay devices may be stored without the need for refrigeration and that test results are obtained almost instantaneously and by visual inspection with naked eye. Immunochromatographic assays may have important advantages when testing in remote areas where access to laboratory facilities is problematic and when testing animals from rural, nomadic and other mobile migratory farmers. Moreover ICA is a sensitive and highly specific test.

Comparing the results obtained by the different serological tests employed in this study, it is clear that no single test is cable to identify all positive cases of brucella infected animals due to variation in sensitivity and specificity of different serological tests as a result of difference in sensitivity of these tests to different antibody classes and variation in antibody classes in the examined sera. The obtained results assured that different



serological tests could detect infected animals with brucella with a different sensitivity and specificity. CFT is still superior one among the employed tests as it gave the higher balance of sensitivity and specificity than other tests used. Whereas BAPAT was found to be an effective test for initial screening of brucellosis in farm animals as it was more sensitive for detection of brucella infected animals than RBPT beside that, it is simple, easy and inexpensive test.

The prevalence of brucella infection in camels in this study may be attributed to lacking of a national program for camel brucellosis eradication including periodical testing and slaughtering of reactors, absence of vaccination program for camels according to Egyptian Field Strain and which proved with imported camels and several exposure of this species to brucella infection than other animals.

### CONCLUSION

From the results of the present study, it is concluded that:

1. The buffered acidified plate antigen test is an effective test for initial screening of brucellosis in farm animals
2. No single serological test could be identify all brucellosis infected camels.
3. Complement Fixation test is still the superior one among the employed tests as it gave the highest balance of sensitivity and specificity.
4. Immunochromatographic assay (ICA) could be recommended as a rapid screening or presumptive test in brucellosis control programs as it easily performed ,high sensitive and specific. Also as confirmatory test so it can replaces the RBPT .
5. Thus study throw a spot light to include camels in the national program for control and eradication of brucellosis in Egypt as the base that where brucellosis exists in stock animals.

### REFERENCES

1. **Refai M (2002):** Incidence and control of brucellosis in the near East region. J. Egypt. Vet. Med. Asso., 90 (1-4): 81-110.
2. **OIE (2008):** Manual of the diagnostic tests and vaccines for terrestrial animals, Vol.1, 5 Edition, pp.328-345 (Office International Des Epizooties, Paris, France).
3. **El-Hady MM (2002):** Some studies on brucellosis in farm animals with special reference to camels. M.V.Sc., Thesis, (Infectious Diseases), Faculty of Veterinary Medicine, Zagazig University.
4. **Abdel Moghney AF (2004):** A preliminary study on brucellosis on camels at Behira province. Ass. Univ. Bull. Environ. Res., 7 (1): 39-43.
5. **Lobna, MAS (2006):** Epidemiological studies on brucellosis in animal and man with special reference to prevention and control. Ph.D., Thesis, (Zoonosis) Faculty of Veterinary Medicine Moshtohor , Benha University.
6. **Abdoel T, Dias IT, Cardoso R and Smits HL (2008):** Simple and rapid field tests for brucellosis in livestock. Vet. Microbiol., 130: 312-319.
7. **Baddour MM (2012):** Diagnosis of Brucellosis in Humans. J. Vet. Adv. 2 (4): 149-156.
8. **Elshemey TM and Abd-Elrahman AH (2014):** Evaluation of a rapid immunochromatographic test for detection of *Brucella abortus* antibodies in Egyptian cattle sera and milk. Alexandria Journal of Veterinary Sciences, 40:24-28.
9. **Alton G, Jones L, Angus R and Veger J (1988):** Techniques for the Brucellosis Laboratory. Institute national de la recherche Agronomique Paris, France.
10. **Bronsvoot BM, Koterwas B, Land F, Handle IG and Tucker J (2009) :** Comparison of a flow assay for brucellosis antibodies with the reference cELISA test

- in west African *Bos indicus*. PLoS ONE 4 (4):5221.
11. **Morgan WJB., MacKinnon DJ, Gill KPW, Gower SGM and Norris PIW (1978):** Standard laboratory techniques for the diagnosis of brucellosis. Report Series No. 1, Weybridge cent. Vet. Lab., England.
12. **Montaser AM, Khoudair RM, Hazem SS and Khairy EA (2012):** Evaluation of immunochromatographic assay for serodiagnosis of brucella among cattle, sheep and goats in Egypt. *Global Veterinaria*, 8 (5): 512-518.
13. **Junaidu AU, Oboegbulem SI and Salihu MD (2008):** Seroprevalence of brucellosis in prison farm in Sokoto, Nigeria. *Asian J. Epidemiol.*, 1: 24-28.
14. **Nada AR (1990):** Further studies on brucellosis in camels. Ph. D., Thesis (Infectious Diseases) Faculty of Veterinary Medicine, Cairo University.
15. **Abou Zaid AA (1998):** Some studies on camel brucellosis. 8<sup>th</sup> sci. Conf., Fac. Vet. Med., Assiut University, Egypt, pp 690-705.
16. **Barsoum SA, El-Sayed MM and El-Fayoumy (1995):** Sero-epidemiological study on camel brucellosis. *Beni Suif. Vet. Med. Res.*, V. (2): 119-126.
17. **Ali MM, Makar NH and Scddck SR (2005):** Serological study of brucellosis on camels in Assiut and the New Valley governorates. *Assiut Vet. Med. J.*, 51 (105): 158-14.
18. **Khoudair RM, Ibrahim EM, Saker GG and Hafez MA (2009):** Clinicodiagnostic and pathological studies on cattle and buffaloes suffering from brucellosis and tuberculosis in Kafr El Sheikh Governorate. *Egypt. J. Comp. Path. and Clinic. Path.*, 22 (1): 148-174.
19. **Nielsen K and Yu WL (2010):** Serological diagnosis of brucellosis. *Contributions, Sec. Biol. Med. Sci.*, XXXI (1): 65-89.
20. **Nielsen K, Gall DP, Smith S, Balsevicius and Garrido F (2004):** Response of serological tests for the detection of ovine and caprine antibody to *Brucella melitensis*. *Rev. Sci. Tech.*, 23: 979-987.
21. **Khoudair RM, Hazem SS and Khairy EA (2013):** Seroprevalence of *Brucella* infection among buffaloes in Gharbyia Governorate. *Global Veterinaria*, 11 (2): 206-213.

## المخلص العربي

## التشخيص السيرولوجي لمرض البروسيللا في الجمال بمحافظة الشرقية ، مصر

محمد إبراهيم عيسى\*، عفاف محمد منازع\*، رمضان محمد خضير\*\*، الشيماء عبدالفتاح عبدالله\*\*\*  
 \* قسم طب الحيوان- كلية الطب البيطرى – جامعة الزقازيق – مصر  
 \*\* معهد بحوث صحة الحيوان – الدقى – الجيزة – مصر  
 \*\*\* المستشفى البيطرى التعليمى – كلية الطب البيطرى – جامعة الزقازيق – مصر

تم تجميع ٢١٠ عينة سيرم من الجمال من عدة مجازر مختلفة بمحافظة الشرقية وذلك للتعرف على معدل الإصابة بمرض البروسيللا فى الجمال. وتم فحصها سيرولوجيا بواسطة اختبار الانتيجين الحمضى المتوازن والروزبنجال الأحمر الحمضى السريع وتحليل المناعة الكروماتوجرافى بالإضافة إلى إجراء الإختبارات التأكيدية الأخرى مثل اختبار التثبيت المكمل واختبار الريفانول ، ومقارنتهم باختبار التحليل المناعى الكروماتوجرافى وذلك الوصول إلى أنسب طريقة لتشخيص المرض ودراسة التحليل الإحصائى لحساسية وخصوصية هذه الإختبارات التشخيص.

وقد أظهرت النتائج بالاختبارات السيرولوجية المختلفة نسبة إصابة بمعدل ٧,٦% ، ٦,٧% ، ٦,٢% ، ٦,٢% و ٦,٧% وذلك باستخدام إختبار الأنتيجين الحمضى المتوازن وإختبار الروز بنجال وإختبار الريفانول وإختبار التثبيت المكمل وتحليل المناعة الكروماتوجرافى على التوالى. بينما كانت نسبة الحساسية لكل من إختبار الأنتيجين الحمضى المتوازن وإختبار الروزبنجال وإختبار الريفانول وإختبار التثبيت المكمل وتحليل المناعة الكروماتوجرافى ٩٥,٤٥% ، ٨٧,٧٥% ، ٩٠,٩٠% ، ٨٨,٨٥% و ٩٢,١٥% على التوالى أما بالنسبة للخصوصية فكانت ٩٩,٢٧% ، ٩٩,٦١% ، ٩٩,٨٠% ، ٩٩,٨٠% و ٩٩,٧٥% لنفس الإختبارات السابقة على التوالى.

ومن نتائج هذه الدراسة نوصى بالآتى:

- ١- استعمال إختبار الانتيجين الشريحي الحمضى كإختبار مسح أولى لتشخيص المرض لشدة حساسيته فى الكشف عن الحيوانات المصابة بالإضافة إلى أنه إختبار بسيط وسريع وغير مكلف. مع تأكيد نتائجه بنتائج إختبار تثبيت المكمل التى أكدت الدراسة كفاءته العالية فى معرفة الحيوانات المصابة.
- ٢- استعمال التحليل الكروماتوجرافى كإختبار مسح لتشخيص المرض وأيضاً كإختبار تأكيدى فمن الممكن استخدامه ليحل محل إختبار الروزبنجال الأقل فى الحساسية والخصوصية كما أنه يمكن استخدامه كبديل لكل من الإختبارات المعقدة عالية التكلفة وعالية الجهد وذلك لشدة حساسيته وسرعته وسهولة إجرائه وكفاءته وانخفاض تكلفته كما أنه من الممكن استخدامه فى المناطق الريفية والنائية دون الاحتياج إلى الامكانيات الخاصة بالمعامل لإجرائه.