

**Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.**



The Journal of Medical Entomology and Parasitology is one of the series issued quarterly by the Egyptian Academic Journal of Biological Sciences. It is an important specialist journal covering the latest advances in that subject.

It publishes original research and review papers on all aspects of basic and applied medical entomology, parasitology and host-parasite relationships, including the latest discoveries in parasite biochemistry, molecular biology, genetics, ecology and epidemiology in the content of the biological, medical entomology and veterinary sciences.

In addition to that, the journal promotes research on the impact of living organisms on their environment with emphasis on subjects such a resource, depletion, pollution, biodiversity, ecosystem.....etc.

www.eajbs.eg.net



Insecticidal Properties of Three Plant Extracts Against *Culex quinquefasciatus* Say and *Aedes aegypti* Linnaeus

Abhay J Khandagle

Department of Zoology Prof.Ramkrishna More ACS College, Akurdi,Pune- 411035

E-mail: ajkhandagle@gmail.com

ARTICLE INFO

Article History

Received:30/10/2019

Accepted:25/11/2019

Keywords:

Lantena camera;
Mentha piperita
L.; *Eucalyptis*
grandis; *Aedes*
aegypti; *Culex*
quinquefasciatus;
larvicide;
repellency.

ABSTRACT

Aedes aegypti is one of the mosquito species responsible for the transmission of Dengue fever, chikungunya, yellow fever and the worst, dengue hemorrhagic fever while *Culex quinquefasciatus* vectors Japanese encephalitis (JE) *Wuchereria bancrofti* and is responsible for several human and animal diseases. World Health Organization stated that about 2/5 of the global human population are currently threaten of dengue and the best way to control the transmission of dengue virus is fighting the mosquitoes that cause the disease. Indiscriminate use of several conventional mosquitocidal agents though effective cause several problems to non-target organism including human and affect the ecological balance as well. Thus there is a need to develop an alternative strategy to manage mosquito populations. Biological products like plant extracts are one of the ways to deal with mosquito control. The secondary metabolites of several plants due to their co-evolution with insects are known to have novel mosquitocidal molecules. The objective of the present study is to evaluate the bioactive potential of *Lantena camera*, *Mentha piperita* L. and *Eucalyptis grandis*. against *Culex quinquefasciatus* and *Aedes aegypti*. The leaf extracts were assessed for its larvicidal and repellency activity by standard methods. The highest larvicidal activity was obtained at LC50=96 ppm and 103 ppm against *C.quinquifasciatus* and *Aedes aegypti* respectively with *M.piperita* extract while 100 % repellency was exhibited by *Eucalyptis grandis* extract up to 240 minutes for both the mosquito species. These results reveal that the selected plants have the potential to be considered in mosquito control programs.

INTRODUCTION

Mosquitoes are well-known vectors that spread several pathogens causing human diseases such as malaria, dengue, filariasis, and several types of encephalitis including West Nile fever (Service, 1993). They are markers of poor sanitation and unhygienic conditions. Mosquito transmitted diseases are the major cause of human death worldwide. No part of the world is free from vector borne diseases (Fradin and Day 2002). About 700 million people suffer from these diseases every year (Taubes 1997). Mosquitoes are recognized vectors of various human diseases in tropical and subtropical countries (Kovendan 2011). In addition to their role as vectors they cause lot of nuisance and conditions like allergic reaction that includes local skin and systemic sensitivity. Global warming, periodic flooding, and deforestation have opened new habitats to mosquitoes which show high plasticity in their breeding behavior and readily spread their distribution. In fact, sporadic malaria outbreaks

have been reported in non-endemic countries and transmission of the disease was caused by the bite of a locally infected *Anopheles* species (Zucker, 1996).

More than two billion people, mostly in tropical countries, are at risk of mosquito-borne diseases, such as malaria, dengue, haemorrhagic fever and filariasis (Snow RW, 2005). The widely and commonly used chemical method though effective has some major demerits making insect control practically difficult. The major drawbacks of these synthetic insecticides are that they are generally non-biodegradable, toxic to non-targets, and vectors develop resistance against them (Evans and Raj, 1988). In view of the above, it is unavoidable to search for new molecules, which are eco-friendly, cheaper, and safer (Khandagle *et al.*, 2011). Recently, the environmentally friendly and biodegradable natural insecticides of plants origin have been receiving attention as an alternative green measure of control of arthropods of public health importance (Nathan SS *et al.*, 2005).

Compared to other synthetic compounds, natural products are presumed to be safer for human use, justifying, therefore, a broad search for eco-friendly biological materials to be used for the control of vectors of medical importance. The chemical contents extracted from plant materials can be useful as repellents, larvicides, oviposition attractants, insect growth hormone regulators and deterrent agents (Kilonzo BS *et al.*, 2001). Plant products have been used in many parts of the world for killing or repelling mosquitoes either as extracts or oils or as the whole plant. Many plant extracts have been studied for their efficacy in controlling larvae of different mosquito species (Kumar and Dutta, 1987; Evans and Raj, 1988; Markouk *et al.*, 2000). Plant products can be used, either as insecticides for killing larvae or adult mosquitoes or as repellents for protection against mosquito bites, depending on the type of activity they possess. A large number of plant extracts have been reported

to have mosquitocidal or repellent activity against mosquito vectors, but very few plant products have shown practical utility for mosquito control. Some indigenous plant-based products are very promising against mosquitoes and can be used as insecticides and/or repellents. In the present study *Mentha piperita* L., *Lantana camara* Linn. and *Eucalyptis grandis* were selected to evaluate their potential as mosquitocide against *Culex quinquefasciatus* and *Aedes aegypti*.

Lantana camara Linn. is a flowering ornamental plant belonging to family Verbenaceae. *L. camara* is also known as Lantana, Wild Sage, Surinam Tea Plant, Spanish flag and West Indian lantana. It is a well-known medicinal plant in the traditional medicinal system.

L. camara is a low erect or subscaudent vigorous shrub with tetragonal stem, stout recurved pickles and a strong odour of black currents. The plant grows up to 1 to 3 meters and it can spread to 2.5 meters in width. Leaves are ovate or ovate-oblong, acute or subacute, crenate serrate, rugose above, scabrid on both sides. The leaves are 3-8 cm long by 3-6 cm wide and green in colour. Leaves and stems are covered with rough hair. Small flower held in clusters. Colour usually orange, sometime varying from white to red in various shades and the flower usually change colours as they age (Khare, 2007; Kirtikar and Basu 2006; Chopra *et al.*, 1956). Different parts of *L. camara* are reported to possess essential oils, phenolic compounds, flavonoids, carbohydrates, proteins, alkaloids, glycosides, iridoid glycosides, phenyl ethanoid, oligosaccharides, quinine, saponins, steroids, triterpens, sesquiterpenoides and tannin as major phytochemical groups. (Ganjewala *et al.*, 2009; Kensa, 2011)

M. piperita is commonly called as peppermint. It is widely used in food, cosmetics, and medicines. It has been proven helpful in symptomatic relief of the common cold. It may also decrease symptoms of

irritable bowel syndrome and decrease digestive symptoms, such as dyspepsia and nausea. It is used topically as an analgesic and to treat headaches. Peppermint is on FDA's generally recognized as a safe list. Its essential oil contains 44 % menthol, 15–30 % Methone, and 5 % esters, in addition to various terpenoids. Other compounds found in the peppermint are flavonoids (12 %), polymerized polyphenols (19 %), carotenes, tocopherols, betaine, and choline (Murray 1995).

Eucalyptus grandis attains a height of 45-55 m, usually with an excellent trunk and a wide-spreading, rather thin crown; most of the bark and branches are smooth, white or silvery, sometimes greenish, rough on the lower stem, smooth above, debark easily. Juvenile leaves are petiolate, opposite for several pairs then alternate, ovate up to 16 x 8.5 cm, green to dark green and slightly wavy; adult leaves are petiolate, alternate, stalked, lanceolate to broad lanceolate, up to 15 x 3 cm, green on topside and pale green on underside, slightly wavy, with a long point. Inflorescence axillary and simple, 7 flowered; peduncles flattened, to 1.8 cm long; buds have a bluish bloom. Fruit or seed capsules several, short-stalked, pear-shaped

or conical, slightly narrowed at the rim, thin, 8 x 6 mm, with whitish waxy coating, narrow sunken disc, and 4-6 pointed, thin teeth, slightly projecting and curved inward, persisting on twigs. (Orwa *et al*, 2009) The main constituents of the oil of the *E. grandis* are α -Pinene (29.69%), p-Cymene (19.89%), 1,8-cineole (12.80%), α -Terpineol (6.48%), Borneol (3.48%) and 3.14% D-Limonene (Oluwagbemiga *et al.*, 2013).

The methanolic extracts of the selected plant leaves were assessed for their larvicidal and repellency potential against the mosquito species.

MATERIALS AND METHODS

Preparation of the Plant Samples and Extraction of Plants:

Leaves of the selected plants were dried for four weeks and the 50-gram powder was used for extraction through Soxhlet apparatus in 400 ml methanol for about 36 hours and a concentrated solution was obtained. After evaporation of solvent, the extracted compound in dried form was obtained. The extracted compound was stored in air-tight desiccator and further used for experiments.



Fig:1 Photographs of the selected plants:

Mosquito Rearing:

The mosquitoes, *A. Aegypti* and *Culex quinquefasciatus* were reared (as per WHO guidelines) in the Department of Zoology, Prof. Ramkrishna More College, Akurdi Pune -44. Mosquitoes were kept at (28± 2) °C, 75%-85% relative humidity (RH), with a

photoperiod of 12 h light, 12 h dark. The adults were reared in separate metal cages 24"x24"x12" with cotton sleeve at one end to have easy access to the culture. The larvae were fed on dog biscuits and yeast powder. Adults were provided with 10% sucrose solution and blood meal.

Larvicidal Bioassay:

The larvicidal activity was performed according to the guidelines for laboratory and field testing of mosquito larvicides published by WHO (who/cds/whopes/gcdpp/2005.13), with minor modifications.

Initially, the mosquito larvae were exposed to a wide range of test concentrations and a control to find out the activity range of the extracts under test. Batches of 25 third-instar larvae of the mosquitoes were placed in a small plastic container with 200 ml dechlorinated water. Small, unhealthy or damaged larvae were removed and replaced. The depth of the water in the containers was between 8 cm and 10 cm. Larval bioassays were carried out using the desired concentration of the extract. Five replicates per concentration and 6 concentrations in the activity range of the extract were used. Larval mortality was recorded after 24 h exposure and was calculated using the following formula :

Percentage mortality = Number of dead larvae / Number of larvae introduced x 100

Results were subjected to statistical analysis.

Repellent activity:

The repellent activity was evaluated using the human bait technique (WHO 1996; Schreck and McGovern 1989). An evaluation was carried out in net cage 30×30×30 cm containing 30 blood-starved female mosquitoes. The volunteer had no contact with lotions, perfumes, oils, or perfume soap on the day of assay. Dose used was 1.0 mg/cm². The surgical glove with a window of 2 cm×2 cm was used. A 2 cm×2 cm muslin cloth uniformly treated with the test extracts were fixed on window of the glove while for control only the solvent was used to treat the muslin cloth. After every 30 minutes, the hand with the glove was offered to the mosquitoes separately to check number of bites for 5 min. This was continued till the bites were received. The experiment was repeated five times. Method used for this assay was same for both mosquito species.

RESULTS AND DISCUSSION

The disadvantages of the conventional chemicals used to control mosquitoes have made it imperative to search for new safer and cheaper methods. Plant products due to their metabolites have always been preferred to be considered as an eco-friendly alternative to the conventional chemical method of insect control. The results of the present study reveal that all three selected plant extracts have considerable larvicidal activity. The most larvicidal potential was observed in the extract of *Mentha piperita* against both the mosquito species viz. *Culex quinquefasciatus* and *Aedes aegypti* with LC₅₀ value of 96 ppm and 103 ppm respectively, while LC₅₀ of 116 ppm and 121 ppm was exhibited by the extract of *Eucalyptis grandis* against *Culex quinquefasciatus* and *Aedes aegypti* respectively. As compared to the other two extracts the extract of *Lantana camera* showed weak larvicidal potential viz, LC₅₀ = 159 ppm against *Culex quinquefasciatus* and LC₅₀ = 167 ppm against *Aedes aegypti*. These results are more significant or at par with the several reported findings, Rahuman and Venkatesan (2008), reported the larvicidal activity of methanolic leaf extracts of *Coccinia indica*, *Momordica charantia*, *Trichosanthes anguina*, and *Cucumis sativus* against *A. aegypti*, i.e., LC₅₀=309.46, 199.14, 554.20, and 492.73 ppm, respectively. They have also reported larvicidal activity of the same plant extracts against *C. quinquefasciatus*, i.e., LC₅₀=377.69, 207.61, 842.34, and 623.80 ppm, respectively. Ethanol fractionate of *E. crassipes* showed the highest larvicidal and pupicidal activity against *C. quinquefasciatus* with LC₅₀ =71.43, 94.68,120.42, 152.15 and 173.35 ppm for I, II, III, IV instars and pupae, respectively (Jayanthi et al 2012). The first- to fourth-instar larvae and pupae of *A. stephensi* had values of LC₅₀ = 272.50, 311.40, 361.51, 442.51, and 477.23 ppm, and the LC₉₀ = 590.07, 688.81, 789.34, 901.59, and 959.30 ppm; the *A. aegypti* had values of LC₅₀ = 300.84, 338.79, 394.69, 470.74, and 542.11 ppm, and the LC₉₀ = 646.67, 726.07,

805.49, 892.01, and 991.29 ppm, respectively. (Panneerselvam *et al.*, 2012) Leaf extract of *Vitex negundo* showed an LC₅₀ of 212.57 ppm against the fourth instar larvae of *C. quinquefasciatus* (Kannathasan *et al.*, 2007). Mullai and Jebanesan 2007 have reported larvicidal activity against the third instar larvae of *C. quinquefasciatus* (LC₅₀ = 118.74 ppm). Methanol extract of *Jatropha curcas* and *Bacillus thuringiensis israelensis* against I, II, III and IV instar larvae of filarial vector showed promising larvicidal activity (Kovendan *et al.* 2011). LC₅₀=456.29ppm was reported by Gunabalan Madhumitha *et al.*, 2012 against *C. quinquefasciatus*. Sharma *et al.* (2005) reported that the acetone extract of *Nerium indicum* and *Thuja orientalis* has been studied with LC₅₀ values of 200.87, 127.53, 209.00 and 155.97 ppm against III instar larvae of *A. stephensi* and *C. quinquefasciatus*, respectively. The LC₅₀ values of aqueous extract from leaves of *Ricinus communis* were 1,091.44; 1,364.58; and 1,445.44 ppm against 2nd, 3rd, and 4th larval instars of *C. quinquefasciatus* (Elimam *et al.* 2009). Murugan *et al.* (2012) described the LC₅₀ for *C. sinensis*, was determined for the larvicidal and pupicidal activities against mosquito vector species from first to fourth larval instars and pupae the values for *A. stephensi* were 182.24, 227.93, 291.69, 398.00 and 490.84 ppm; *A. aegypti* values were 92.27, 106.60, 204.87, 264.26, 342.45, 436.93 and 497.41 ppm; and *C. quinquefasciatus* values were 244.70, 324.04, 385.32, 452.78 and 530.97 ppm, respectively. Shahi *et al.*, 2010, reported the alcoholic extract of *C. procera* showed to be less toxic than latex in both mosquito species. The LC₅₀ values were 109.71 and 387.93 mg/l for *An. stephensi* and *Cx. quinquefasciatus*, respectively. These figures were 13.06 and 86.47 mg/l respectively for latex of the plant. The 512 ppm concentration of plant extract didn't show a mortality rate >78% in *Cx. quinquefasciatus* after 24 h. But in the case of *An. stephensi* we observed >95% mortality after 24 h from 256 ppm. Tests with latex showed 99%

mortality at 64 ppm for *An. stephensi*, only 44% mortality against *Cx. quinquefasciatus* and a maximum of 67% in 256 ppm.

Gandhi *et al.*, 2016, have reported the bioactive potential of both the extract and the isolated compound and upon screening one of the fraction from the methanol extract of *R. cordifolia* showed good mosquitocidal activity against *C. quinquefasciatus* and *A. aegypti*. LC₅₀ and LC₉₀ values of fraction 2 were 3.53 and 7.26 ppm for *C. quinquefasciatus* and 3.86 and 8.28 ppm for *A. aegypti* larvae, and 3.76 and 7.50 ppm for *C. quinquefasciatus* and 3.92 and 8.05 ppm for *A. aegypti* pupae, respectively. Further, the isolated compound alizarin presented good larvicidal and pupicidal activities. LC₅₀ and LC₉₀ values of alizarin for larvae were 0.81 and 3.86 ppm against *C. quinquefasciatus* and 1.31 and 6.04 ppm for *A. aegypti* larvae, respectively. Similarly, the LC₅₀ and LC₉₀ values of alizarin for pupae were 1.97 and 4.79 ppm for *C. quinquefasciatus* and 2.05 and 5.59 ppm for *A. aegypti* pupae, respectively.

The highest larval mortality was found in the hexane extract of *Z. zerumbet*, ethyl acetate extract of *D. biflorus*, and methanol extracts of *A. indica* against *C. gelidus* (LC₅₀ = 26.48, 33.02, and 12.47 ppm; LC₉₀ = 127.73, 128.79, and 62.33 ppm) and against *C. quinquefasciatus* (LC₅₀ = 69.18, 34.76, and 25.60 ppm; LC₉₀ = 324.40, 172.78, and 105.52 ppm), respectively, after 24 h (Kamaraj *et al.*, 2010). The maximum repellent activity was observed at 450 ppm in ethanol extracts of *C. sinensis* and the mean complete protection time ranged from 150 to 180 min. The ethanol extract of *C. sinensis* showed 100 % repellency up to 150 min and showed complete protection up to 90 min at 350 ppm against *A. stephensi*, *A. aegypti* and *C. quinquefasciatus*, respectively (Kadarkarai Murugan *et al.*, 2012). Amer and Mehlhorn, 2006, have reported that the five most effective oils were those of *Litsea cubeba*, Cajeput (*Melaleuca leucadendron*), Niaouli (*Melaleuca quinquenervia*), Violet (*Viola odorata*), and Catnip (*Nepeta cataria*), which induced a

protection time of 8 h at the maximum and a 100% repellency against *Ae. aegypti*, *An. stephensi*, and *Cx. quinquefasciatus*. The essential oil of *Zingiber officinalis* showed repellent activity at 4.0 mg/cm², which provided 100% protection up to 120 min against *Cx. Quinquefasciatus* (Pushpanathan *et al.*, 2008)

Effiom *et al.*, 2012, reported the mosquito repellent activity of diethyl ether extracts from Peels of five citrus fruit species, *Citrus sinensis*, *Citrus limonum*, *Citrus aurantifolia*, *Citrus reticulata*, and *Citrus vitis*, at five different concentrations, 5%, 10%, 15%, 20%, and 25%. Topical application revealed that 20% and 25% repelled mosquitoes 2 hours and 5 hours, respectively. At a dose of 0.1 mg/cm², potent repellency against mosquito adults was obtained with the extracts of *Cinnamomum cassia* Blume bark (91%), *Nardostachys chinensis* Batalin rhizome (81%), *Paeonia suffruticosa* Andrews root bark (80 %), and *C. camphora* steam distillate (94 %). *Eugenia caryophyllata* Thunb. extract provided 75 % repellency (Young *et al* 2004)

Kamaraj *et al.*, 2011 reported for maximum repellent activity at 500 ppm in methanol extracts of *N. nucifera*, ethyl acetate and methanol extract of *P. nigrum* and methanol extract of *T. ammi* the mean complete protection time ranged from 30 to 150 min with the different extracts tested. Table 2 depicts the repellency potential of the extracts of the selected plants against both the mosquito species and the obtained results show that *Eucalyptis grandis* extract has 100 % repellency up to 240 minutes against both mosquito species.

These plants certainly have bioactive potential that can be further explored with different field trials. Toxicological tests should also be done before including them in the mosquito control strategy. These extracts may also be used in a combinatorial way to step up the larvicidal potential.

Though the knockdown effect of synthetic chemicals is remarkable they bring irreversible environmental hazards, severe side effects and pernicious toxicity to human beings and beneficial organisms. In the light of the recognized demerits of the chemical control method, emphasis on controlling mosquito vectors has shifted steadily from the use of conventional chemicals toward alternative insecticides that are target-specific, biodegradable, and environmentally safe. Recently, the use of environment-friendly and biodegradable natural insecticides of plant origin has established renewed consideration as agents for vector control as they are rich in bioactive chemicals, active against a limited number of species including specific target insects. Among the biopesticides in practice plant extracts and essential oils are of choice and great help to control mosquitoes. These results indicate the potential of the plant extracts and their possibility to be included in the mosquito management program. It can also be a part of IPM practice to control mosquitoes. Further exploration of these plants for their active principle and insecticidal potential even against other vectors can be thought of. With further field trials and toxicity tests these extracts may be a helpful addition to the mosquito control strategy.

Table 1: Larvicidal activity of some plant extracts against *A.aegypti* and *C. quinquefasciatus*

Extract	Mosquito	LC ₅₀ ±SE (ppm)	95% Confidential limit		Regression equation	LC ₉₀ (ppm)
			LCL	UCL		
<i>Lantana camera</i> ,	<i>A.aegypti</i>	167±0.77	124	291	Y=3.82X-4.06	297
	<i>C.quinquefasciatus</i>	159±1.05	118	284	Y=4.32X-5.24	269
<i>Mentha piperita</i>	<i>A.aegypti</i>	103±0.45	88	143	Y=3.98X-6.76	161
	<i>C.quinquefasciatus</i>	96± 0.67	74	132	Y=6.24X-4.87	130
<i>Eucalyptis grandis.</i>	<i>A.aegypti</i>	121± 0.91	91	162	Y=3.29X-5.22	171
	<i>C.quinquefasciatus</i>	116±0.88	82	152	Y=2.84X-5.77	161

Table 2: Repellent activity of some plant extracts against *A.aegypti* and *C. quinquefasciatus*

Extract	Mosquito	Percent Repellency after								
		30 min	60 min	90 min	120 min	150 min	180 min	210 min	240 min	270 min
<i>Lantana camera</i> ,	<i>A.aegypti</i>	100	100	95	87	82	75	70	64	56
	<i>C.quinquefasciatus</i>	100	100	94	89	84	79	73	68	62
<i>Mentha piperita L.</i>	<i>A.aegypti</i>	100	100	100	97	91	85	79	73	69
	<i>C.quinquefasciatus</i>	100	100	100	96	90	84	80	76	70
<i>Eucalyptis grandis.</i>	<i>A.aegypti</i>	100	100	100	100	100	100	100	100	94
	<i>C.quinquefasciatus</i>	100	100	100	100	100	100	100	100	93

REFERENCES

Amer A, Mehlhorn H. (2006) Repellency effect of forty-one essential oils against *Aedes*, *Anopheles*, and *Culex mosquitoes*. Parasitol Res, 99: 478-490.

Chopra RN, Nayar SL and Chopra IC. (1956) Glossary of Indian medicinal plants. CSIR New Delhi, India.

Effiom, O. E. Avoaja D. A, & Ohaeri, C. C (2012) Mosquito Repellent Activity of Phytochemical Extracts from Peels of Citrus Fruit Species, Global Journal of Science Frontier Research Interdisciplinary, Volume XII Issue I version I

Elimam AM, Elmalik KH, Ali FS (2009) Larvicidal, adult emergence inhibition and oviposition deterrent effects of foliage extract from *Ricinus communis* L. against *Anopheles arabiensis* and *Culex quinquefasciatus* in Sudan. Trop Biomed 26(2):130–139.

Evans DA, Raj PK. (1988); Extract of Indian plants as mosquito larvicides. *Indian J Med Res* 88: 38–41.

Fradin MS, Day JF (2002) Comparative efficiency of insect repellents against mosquito bites. N Engl J Med 347:13–18.

Gandhi, M.R., Reegan, A.D., Ganesan, P. et al. (2016) Larvicidal and Pupicidal Activities of Alizarin Isolated from Roots of *Rubia cordifolia* Against *Culex quinquefasciatus* Say and *Aedes aegypti* (L.) (Diptera: Culicidae). Neotrop Entomol 45: 441.

Ganjewala D, Sam S and Khan KH. (2009) Biochemical compositions and antibacterial activities of *Lantana camara* plants with yellow, lavender, red and white flowers. EurAsian Journal of BioSciences. 3: 69-77.

Gunabalan Madhumitha, Govindasamy Rajakumar, Selvaraj Mohana Roopan, Abdul Abdul Rahuman, Kanagaraj Mohana Priya , Antoneyraj Mary Saral , Fazlur Rahman Nawaz Khan ,

- Venkatesh Gopiesh Khanna , Kannaiyaram Velayutham , Chidambaram Jayaseelan , Chinnaperumal Kamaraj and Gandhi Elango (2012) Acaricidal, insecticidal, and larvicidal efficacy of fruit peel aqueous extract of *Annona squamosa* and its compounds against blood-feeding parasites, *Parasitol Res* 111:2189–2199.
- Jayanthi P. & P. Lalitha & N. Aarthi (2012) Larvicidal and pupicidal activity of extracts and fractionates of *Eichhornia crassipes* (Mart.) Solms against the filarial vector *Culex quinquefasciatus* Say, *Parasitol Res* 111:2129–2135
- Kadarkarai Murugan , Palanisamy Mahesh Kumar, Kalimuthu Kovendan, Duraisamy Amerasan, Jayapal Subrmaniam and Jiang-Shiou Hwang (2012) Larvicidal, pupicidal, repellent and adulticidal activity of *Citrus sinensis* orange peel extract against *Anopheles stephensi* *Aedes aegypti* and *Culex quinquefasciatus* (Diptera: Parasitology Research 111(4). DOI: 10.1007/s00436-012-3021-8
- Culicidae), *Parasitol Res* 111:1757–1769
- Kamaraj, C., Rahuman, A.A., Mahapatra, A. et al. (2010) Insecticidal and larvicidal activities of medicinal plant extracts against mosquitoes *Parasitol Res* 107: 1337.
- Kamaraj C, Rahuman AA, Bagavan A, Elango G, Zahir AA, Santhoshkumar T. (2011) Larvicidal and repellent activity of medicinal plant extracts from Eastern Ghats of South India against malaria and filariasis vectors. *Asian Pac J Trop Med.*4(9):698-705.
- Kannathasan K, Senthilkumar A, Chandrasekaran M, Venkatesalu V(2007) Differential larvicidal efficacy of four species of *Vitex* against *Culex quinquefasciatus* larvae. *Parasitol Res* 101:1721–1723
- Kensa VM.(2011) Studies on phytochemical screening and antibacterial activities of *Lantana camara* Linn. *Plant Sciences Feed.* 1 (5): 74-79. 17.
- Khandagle Abhay J. & Vrushali S. Tare & Kishor D. Raut & Rashmi A. Morey, Bioactivity of essential oils of *Zingiber officinalis* and *Achyranthes aspera* against mosquitoes, *Parasitol Res.* (2011) 109:339–343
- Khare CP. (2007) *Indian Medicinal Plants - An Illustrated Dictionary.* Berlin, Springer.
- Kilonzo BS, Ngomuo AJ, Sabuni CA, Mgoe GF (2001): Effect of *Azadirachta indica* (NEEM) extract on livestock fleas in Morogoro district, Tanzania. *Insect Science Applic*, 21:89-92.
- Kirtikar KR, Basu BD. (2006) *Indian medicinal plants.* New Delhi, India.
- Kovendan K, Murugan K, Vincent S, Kamalakannan S (2011) Larvicidal efficacy of *Jatropha curcas* and *bacterial insecticide, Bacillus thuringiensis*, against lymphatic filarial vector, *Culex quinquefasciatus* Say (Diptera: Culicidae). *Parasitol Res* 109:1251–1257
- Kovendan K. and Murugan K. (2011), Effective of Medicinal Plants on the Mosquito Vectors from the Different Agroclimatic Regions of Tamil Nadu, India, *Advances in Environmental Biology*, 5(2), 335-344.
- Kumar A, Dutta GP (1987). Indigenous plant oils as larvicidal agents against *Anopheles stephensi*. *Curr Sci.*;56: 959-60.
- Markouk M, Bekkouche K, Larhsini M, Bousaid M, Lazrek H B, Jana M. (2000) Evaluation of some Moroccan medicinal plant extracts for larvicidal activity. *J Ethnopharmacol*; 73: 293-7.
- Mullai K, Jebanesan A (2007) Larvicidal, ovicidal and repellent activities of the leaf extract of two cucurbitaceous plants against filarial vector *Culex quinquefasciatus* (Say) (Diptera: Culicidae). *Tropical Biomed* 24:1–6
- Murray MT (1995), *The healing power of herbs: the enlightened person's guide to the wonders of medicinal plants*, 2nd edn. Prima, Roseville, 410

- Murugan, K., Mahesh Kumar, P., Kovendan, K. et al. (2012) Larvicidal, pupicidal, repellent and adulticidal activity of *Citrus sinensis* orange peel extract against *Anopheles stephensi*, *Aedes aegypti* and *Culex quinquefasciatus* (Diptera: Culicidae) *Parasitol Res* 111: 1757.
- Nathan SS, Kandaswamy K, Kadarkarai M (2005): Effect of neem limonoids on the malaria vector *Anopheles stephensi* Liston (Diptera: Culicidae). *Acta Trop*, 96:47-55.
- Oluwagbemiga Sewanu Soyingbe , Adebola Oyedeji , Albert Kortze Basson , Andy Rowland Opoku (2013) The Essential Oil of *Eucalyptus grandis* W. Hill ex Maiden Inhibits Microbial Growth by Inducing Membrane Damage, *Chinese Medicine*, 4, 7-14
- Orwa C., Mutua A., Kindt R., Jamnadass R., Anthony S., (2009), *Agroforestry Database 4.0*. World Agroforestry Center, Nairobi, Kenya
- Panneerselvam, C., Murugan, K., Kovendan, K. et al. (2012) Mosquito larvicidal, pupicidal, adulticidal, and repellent activity of *Artemisia nilagirica* (Family: Compositae) against *Anopheles stephensi* and *Aedes aegypti* *Parasitol Res* 111: 2241.
- Pushpanathan T, Jebanesan A, Govindarajan M. (2008) The essential oil of *Zingiber officinalis* Linn (Zingiberaceae) as a mosquito larvicidal and repellent agent against the filarial vector *Culex quinquefasciatus* Say (Diptera: Culicidae). *Parasitol Res*; 102:1289-1291.
- Rahuman AA, Venkatesan P (2008) Larvicidal efficacy of five cucurbitaceous plant leaf extracts against mosquito species. *Parasitol Res* 103:133-139.
- Schreck, C. E., and McGovern, T. P. (1989) Repellents and other personal protection strategies against *Aedes albopictus*. *Mosq. Control. Assoc. Journal of the American Mosquito Control Association* 5(2):247-50.
- Service M.W. (1993) Mosquitoes (Culicidae): medical importance, *in: Medical Insects and Arachnids*. R.P. & Crosskey R.W.(eds), Chapman and Hall, London, 5, 196-208
- Shahi M, Hanafi-Bojd AA, Iranshahi M, Vatandoost H, Hanafi-Bojd MY,(2010), Larvicidal efficacy of latex and extract of *Calotropis procera* (Gentianales: Asclepiadaceae) against *Culex quinquefasciatus* and *Anopheles stephensi* (Diptera: Culicidae), *J Vector Borne Dis.*47(3):185-8
- Sharma P, Mohan L, Srivastava CN (2005) Larvicidal potential of *Nerium indicum* and *Thuja orientalis* extracts against malaria and Japanese encephalitis vector. *J Environ Biol* 26(4):657-660
- Snow RW, Guerra CA, Noor AM, Myint HY, Hay SI (2005): The global distribution of clinical episodes of *Plasmodium falciparum* Malaria *Nature*, 434:214-217.
- Taubes, G. (1997). A mosquito bites back. *New York Times Magazine*, 40-46.
- WHO (1996) Report of the WHO informal consultation on the evaluation on the testing of insecticides. *CTD/WHO PES/IC/96.1:69*
- Young-Cheo, Yang, T Eun-Hae Lee, Hoi-Seon Lee, Dong-Kyu Lee and Young-Joon Ahn (2004) Repellency of Aromatic Medicinal Plant extracts and a Steam Distillate to *Aedes aegypti*, *Journal of the American Mosquito Control Association*, 20(2):146-149,
- Zucker J.R. (1996) Changing patterns of autochthonous malaria transmission in the United States: a review of recent outbreaks. *Emerging Infectious Diseases*, 2, 37-43