

## **HETEROSIS AND YIELD COMPONENTS OF SOME F1 HYBRIDS OF SWEET POTATO.**

**Gendy,A. S. and A. G. Mohamed**

**Department of Vegetable Breeding Research, Horticulture Research Institute Egypt Agriculture, Research Center, Egypt.**

### **ABSTRACT**

True seeds of twenty sweet potato lines were collected and used for croup production. The six hybrids were evaluated in a randomized complete blocks design with four replicates with best two check cvs (Mabrouka and Abees) for some morphological and root yield characters at Motopous conditions Kaferel Sheikh, Governorate . Differences were found among the F1 hybrids in all studied traits. All the F1 hybrids significantly exceeded the two check cvs. The percentage of increase in the hybrids with Mabrouka cv. was ranged from (168%) in the hybrid No (1) to (51.5%) in the No (3). While it ranged from (173%) to (56%) compared with Abees cv. Average tuber root was highly significant in hybrids No (4) compared with the two check cvs. Two degrees of skin colour, i.e., purple and dark purple, were observed in roots of the hybrids, while six degrees of flesh colour from white to purple were noticed in this study. Most of F1 hybrids significantly exceeded the to check cultivar in most studied traits. Hybrid vigor attributed to over dominance was detected in total tuber root yield and the other studied traits. Dominance in complete and partial was also observed.

### **INTRODUCTION**

Sweet potato tubers are a good source of vitamin C, carotenes, proteins, carbohydrates and some minerals. Sweet potato is a cross pollinated and hexaploid ( $2n=6x$ ) crop with 90 chromosomes (Jones, 1965).The genetics of sweet potato is little understood and the inheritance pattern is quite complex one. Genetic information on many traits of direct economic importance in Sweet potato are not available and most published information are from the clones of similar genetic back ground Jones, 1966 and Jones *et al.*, 1969.

Studies on the entire spectrum of the variability are therefore necessary to acquire knowledge on the inheritance pattern (Vimalan, 1993).On the other hand foliage in terms of leaves and shoots contain high amount of vitamins A and B as well as proteins, so it also used for animal feeding. Sweet potato flour could be mixed with wheat flour in bread manufacturing and it also a good raw material for sweet industries cakes and biscuits. It also used as a staple food-vegetable (both fresh roots and tender leaves and petioles) - snack food animal feed for, industrial starch extraction and fermentation and for various processed products (Chalfant *et al.*, 1990). Sweet potato has long been very important food crops the movgh tropical and sub tropical regions of the world especially in the eastern countries of Asia it is considered the six most important food crops in the world (Marrison *et al* 1993). It produce agreed yield under a wide range of environments, but most countries of the world cultivated it in summer seasons. Mwanga *et al.* (2007) reported that

high dry matter content is an important characteristic of a good sweet potato variety. Recently, the orange fleshed sweet potato varieties with high  $\beta$ -carotene are being promoted to improve vitamin A nutrition of the poor. Therefore, these varieties should carry increased  $\beta$ -carotene and dry matter to promote their adoption and large-scale production (Cervantes-Flores *et al.*, 2010 and Mwanga *et al.*, 2010). The gene actions need a particular analysis in each case study involving various genotypes and environments through appropriate genetic designs (Acquaah, 2007). Various gene actions have been identified and described in different studies on sweet potato. Miller, (1939) found that white skin, green stem and white flesh are dominant to red skin, red stem and yellow flesh of sweetpotato. Ma. *et al.* (2009) observed that the inheritance of  $\beta$ - carotene is controlled by additive gene effects. Vimalan and Hariprakash (2011) reported that most sweetpotato varieties used in their study were self- and cross-incompatible and therefore these varieties were not able to produce viable seeds when hybridized through both self- and cross-pollinations. The additive gene effects were identified in the inheritance of dry matter and  $\beta$ -carotene content (Chiona, 2009). Heterosis was observed for the size and number of roots per plant, indicating dominance gene action or intra-allelic interaction between alleles of the same gene (Gasura *et al.*, 2010).

As soon as in Egypt. This study aims to produce sweet potato under winter season. To give the highest prices of yields and also high yield from some promising genotypes (F1) of sweet potato with high productivity and aurally in winter season. In addition average degree of heterosis and standered heterosis for some characters were study.

## **MATERIALS AND METHODS**

The present study was carried out at Motopous Kafr El sheikh governorate during two seasons (2012 and 2013) in El Zhraa village. The genetic material used in the study was 20 hybrids which were obtained from the PhD study from A.S.Gendy 2002 Egypt. In July 2012 the seeds were soaked in concentrated sulfuric acid for 30 minutes and washed in water. (Jones *et al.*, 1969) before sowing because sweet potato seeds germination is restricted by a hard seeds coat which delays inhibition and overcame inbreeding programs by mechanical or chemical scarification and sown in nursery. In October (2012), plants were randomly chosen and vegetative propagated by vine cutting to produce the required plants for evaluation traits. The cutting stems were transplanted in an evaluation experimental trial. A randomized complete block design with four replications was used. Each replicate contained six genotypes as well as two sweet potato cultivars Mabrouka and Abees which widely used in Egyptian cultivars as control data were recorded on vegetative traits and tuber root characteristics. Usual fertilization, irrigations, disease and insects control programmers were practical as used with commercial production of sweet potato. The harvest was done at full maturity about 150 days after transplanting. Data were recorded on the following characters:-

#### **A. Plant characters**

These characters were determined after 100 days from transplanting.

- 1- Number of branches/plant.
- 2- Stem length of the main stem (cm), from ground level to the terminal bud of longest vegetative branch.
- 3- Number of leaves /plant.
- 4- Fifth leaf area. This character was determined for the fifth leaf area from the top by using fresh weight method.
- 5- Dry matter percentage 100g of leaves and stems were determined by drying in an oven at 70°C for 3 days (until constant weight). These characters were determined for four plants in each genotype in the two seasons.
- 6-Skin colour.
- 7- Flesh colour.
- 8-Tuber root shape.

#### **B. Yield components**

- 1- Total yield of tuber roots /plant(kg).
- 2- Average number of tuber roots/plant.
- 3- Average tuber root weight (g). It was determined by dividing the weight of tuber roots per plant on the total tuber root number

#### **C- Tuber root characters.**

- 1- Average tuber root shape. It was determined by dividing the root length on the diameter (L/D ratio) and the shape was described according to Human (1991), where 1= round (circular), 2= round elliptic (slightly circular) and 3= elliptic (oblong).
- 2- Skin and flesh colour. They were visual determined the skin colour was rated subjectively from 1 (creamy –white) to 9 (dark-purple) where 1 white-cream, 2 yellow, 3 orange, 4 brownish, 5 pink, 6 red, 7 purplish-red, 8-purple and 9 dark purple for skin colour. The flesh colour was rated subjectively 1-9 from white to deeply orange (dark-orange), where 1 white, 2 cream, 3 pale yellow, 4 yellow, 5 deep yellow, 6 red, 7 violet, 8 purple and 9 other for (flesh colour), according to Human *et al.* (1977).

#### **D. Statistical analysis**

All data obtained from both experimental seasons were subjected to statistical analysis according to Sendecor and Cochran (1972). Mean values represented the various investigated genotypes were compared by least significant difference value (L.S.D) at both 5% and 1% level of significant. Estimates of genetic parameters were calculated according to Fehr (1987) as follows:

a-Average degree of heterosis (ADH%), was expressed as percent increase or decrease of F1 crosses performance above the mid parents (MP) values and the high parent (HP) values as follows :

$$ADH \% (\text{ based on SC}) = \frac{F1 - SC}{SC} \times 100$$

Where: F1 are means of F1 Hybrid and SC are means of standard cultivars respectively.

## RESULTS AND DISCUSSION

Mean performances of resulted six F1 hybrid for some characters during two experiments of seasons are presented in Tables (1 to 5).

### A. Plant characters

#### Number of branches/ plant

As shown in Table (1) number of branches/ plant showed high significant differences among the F1 and check cvs during the two seasons. Number of branches/plant ranged from (26 to 36) with a mean of (31.1) branches in the first season while in the second these values were (24 to 35) with a mean of (29.6) branches. The genotype No( 6) showed the highest branch number per plant (36 branches) in the first season while the genotype No ( 3) and (4) gave the lowest number of branches with values of (24 and 25) in the second season respectively. Regarding check cvs. they showed that the check cultural Mabrouka gave the lowest number of branches /plant in the two season with values of (20 to 21) respectively. Our results are supported by many works so Shalaby *et al.* (1993 a and b). El- Shini (1996) and Salem (1999).

**Table(1). Mean performances of the evaluated sweet potato F1 crosses and check cvs for number of branches/plant and stem length(cm).**

Genotypes	No of branches/ plant		Stem length (cm)	
	2010	2011	2010	2011
H .No1	34.0	32.0	125.0	119.0
H .No2	30.0	28.0	132.0	136.0
H .No3	26.0	24.0	105.0	109.0
H .No4	28.0	25.0	109.0	111.0
H .No5	35.0	34.0	139.0	141.0
H .No6	36.0	35.0	121.0	126.0
Mabrouka	20.0	21.0	181.0	179.0
Abees	27.0	25.0	193.0	187.0

#### Stem length

As shown in Table (1) the studied genotypes significantly differed in this trait during the two seasons, the average stem length ranged from 105 to 139cm with a mean of 121.3 cm in the first season, while ranged from (109 to 141cm ) with a mean of (123.6cm ) in the second season. The line No (5) gave the tallest stem length during the two seasons as showed 139 cm in the first season and to 141cm in the second season. On the other hand the genotype No (3) gave the shortest stem length in the two seasons, with values of (105cm) and 109 respectively. The check cultivars Mabrouka and Abees gave the highest values of stem length Mabrouka recorded 181 to 179 cm while Abees recorded 193 to 187cm in the first and second season respectively. The results are confirmed with those by El- Denary (1998) and Salem (1999) who found significant differences among lines in stem length. Data in Table (1) regarding check cvs, Mabrouka and Abees had high values

of stem length with an average of 180.0 and 190.0 cm respectively. All the six hybrids were found significantly shorter than the check cvs. Mabrouka and Abees in their stem length during the two seasons. These results agree with those by Sozki (1990) Shalaby *et al.* (1993a) and El-Shini (1996) and Gendy (2002).

**Number of leaves/plant**

Significant differences were observed among the studies 6 hybrids concerning this trait during the two seasons were observed as shown in Table (2). The number of leaves per plant ranged from 145 to 295 in first season and ranged from 139 to 305 in second season respectively. The check cv Mabrouka recorded the highest number of leaves/plant in the second season with means of 305.0, the hybrid No (5) gave the better value of 226 in the second season. Data in Table (2) refer to that the 6 hybrids recorded approximately similar behavior in the two seasons. While Mabrouka cv recorded the highest values in the two seasons with values of 295 and 305 respectively. Similar results were obtained from El-Shini 1996, Salem (1999) and Gendy 2002 home found significant differences for this trait among sweet potato hybrids and cultivars.

**The Fifth leaf area**

The Fifth leaf area ranged from 18.9 to 26.5cm<sup>2</sup> in first season and from 19.4 to 27.0 in the second season with means values 22.95cm<sup>2</sup> and the 6 hybrids gave higher value than the two check cultivars Mabrouka and Abees cvs, Mabrouka gave (24.6 and 23.9cm<sup>2</sup>) in the first and second respectively with means of 24.25cm<sup>2</sup> also Abees gave values of 20.1 and 20.2cm<sup>2</sup> with means of 20.15cm<sup>2</sup>.

**Dry matter %**

Data in Table (2) show dry matter % among the hybrids during the two seasons, the hybrids No (6) gave the highest values of 49.2 and 48.3% in the first and second season respectively with means of 48.7% and the 6 hybrids gave higher values than the two check cultivars Mabrouka and Abees cvs, Mabrouka gave (30.1 and 29.7%) in the first season with means 29.9 % also Abees gave values of 31.8 and 32.0% with mean of 32.0%. However all evaluated hybrids exceeded the two cvs cultivars in the two seasons. These results agree with that recorded by Singh *et al.* (1993) and Gendy (2002).

**Table (2). Mean performances of the evaluated sweet potato F1 crosses and check cvs for number of leaves /plant and fifth leaf area/plant .**

Genotypes	No of leaves/plant		Fifth leaf area cm <sup>2</sup>		Dry matter%	
	2010	2011	2010	2011	2010	2011
H .No1	170.0	172.0	26.5	27.0	43.1	42.7
H .No2	195.0	199.0	22.6	23.5	46.5	47.1
H .No3	205.0	222.0	25.1	23.7	44.3	45.1
H .No4	145.0	139.0	25.3	26.1	42.9	43.7
H .No5	218.0	226.0	18.9	19.4	45.2	44.9
H .No6	195.0	190.0	23.6	24.1	49.2	48.3
Mabrouka	295.0	305.0	24.6	23.9	30.1	29.7
Abees	200.0	207.0	20.1	20.2	31.8	32.0

### Skin colour

Data in Table (3) showed that, the studied genotypes did not varied in root skin colour, dark-purple colour was observed in all genotypes as well as the two check cultivars Mabrouka and Abees . These results agree with Gendy (2002) and Vimalan and Hariprakash (2011).

**Table (3). Performances of the evaluated sweet potato F1 crosses and check cvs for skin, flesh colour and root shape.**

Genotypes	Skin colour		Flesh colour		Root shape	
	2010	2011	2010	2011	2010	2011
H .No1	(9)Dark - purple	(9)Dark - purple	8 violet	8 violet	Round	Round
H .No2	(9)Dark - purple	(9)Dark - purple	6 orange	6 orange	Round - elliptical	Round - elliptical
H .No3	(9)Dark - purple	Dark -purple	5 deep- yellow	5 deep- purple	Round - elliptical	Round - elliptical
H .No4	(9)Dark - purple	(9)Dark - purple	5 deep- purple	5 deep- purple	Round - elliptical	Round - elliptical
H .No5	(9)Dark - purple	(9)Dark - purple	1 white	1 white	Round- elliptical	Round- elliptical
H .No6	(9)Dark - purple	(9)Dark - purple	Deep- yellow	Deep- yellow	Round	Round
Mabrouka	(9)Dark - purple	(9)Dark - purple	1 white	1 white	elliptical	elliptical
Abees	(9)Dark - purple	(9)Dark - purple	6 orange	6 orange	elliptical	elliptical

### Flesh colour

Data concerning tuber flesh colour in the evaluated genotypes are shown in Table (3). The hybrids varied greatly in root flesh colour, 6 degrees of colour from white to purple were observed as follow 1 white colour such as the hybrids No (5) and Mabrouka cvs. While the hybrid No 2 and the Abees cvs were 6 orange. The hybrid No 3 and 6 were deep yellow –while the hybrid No (1) was 8 violet. Similar results were reported by some investigators the yellow groups of flesh colour was reported by Wanas (1979)and Sozki (1990). The orange flesh colour in sweet potato tubers was also observed in studies conducted by Jones *et al.* (1977). Our results also agree with those of Salem (1999), Gendy 2002 and Vimalan and Hariprakash (2011), who selected 52 new bred lines reported that the lines varied from white to purple for Flesh colour.

### Tuber root shape

According to data listed in Table (3) three types of shape were observed in the tuber roots of the evaluated 6 hybrids and two cheek cultivars.

- 1- Round such as the hybrids No. 1 and 6.
- 2- Round elliptic which observed in hybrids No. 2, 3 and 5.
- 3- Elliptic such as the two check cultivars Mabrouka and Abees they produce tuber roots with elliptic shape. This results agree with these of Gendy 2002 and Vimalan and Hariprakash (2011).

**Yield and its components:**

**Total yield/plant**

The total yield in the evaluated hybrids and check cultivars are listed in Table (4). The evaluated genotypes significantly differed in total yield as weight of roots/plant during the two seasons. The highest total yield values were obtained by the hybrids No (1) and No (4) they produced more than 2 kg/ plant. On the other hand the lowest yield less than 0.5 kg /plant was produced by the two check cultivars Mabrouka and Abees with mean of 475 g for Mabrouka while Abees gave 415 and 425g with of 420g. The percentage of increase in the best hybrid compared with the best check cultivare Mabrouka was 462.3%. Accordingly, it is very good result of hybrid regarding the 6 genotypes since they gives yield over than the two check cultivars Mabrouka and Abees. The lowest genotypes No (3) gave (980 g), compared with Mabrouka cv.(the best check cvs.) gave (465.0 g), with 103% increase. Accordingly, it is very good result when genotypes outyielded the commercial cultivars Mabrouka and Abees cvs. These results are confirmed with those of El-denary (1998) and Salem (1999) who found significant differences among sweetpotato lines and cultivars studied. Gasura *et al.* (2008) and Vimalan and Hariprakash (2011), demonstrated the possibility to improve sweet potato for yield and quality using the available germplasm.

**Table (4). Mean performances of the evaluated sweet potato F1 crosses and check cvs for yield per plant, number of roots, plant and average root weight .**

Genotypes	Yield/plant (Kg)		Root number/plant		Average root weight (g)	
	2010	2011	2010	2011	2010	2011
H .No1	2.150	1.900	4.0	3.9	537.5	487.2
H .No2	1.950	1.780	2.8	2.6	870	684.6
H .No3	0.980	1.025	2.7	2.9	362.9	353.5
H .No4	2.100	1.985	2.9	2.8	724.1	685.0
H .No5	1.350	1.300	3.1	2.9	435.4	448.2
H .No6	1.650	1.750	3.2	3.3	515.6	530.3
Mabrouka	465.0	485.0	3.6	3.7	129.2	131.0
Abees	415.0	425.0	3.8	3.6	110.0	118.0

**Root number plant**

Significant differences among the evaluated breeding lines were observed in total number of tuber roots per plant during the two seasons as shown in table (4). The hybrid No (1) gave the highest number of roots/ plant (4) while the hybrid No (2) gave the lowest number of roots/plant (2) in the first season in the second season they gave 3.9 and 2.6 number of

roots/plant respectively in contrast Mabrouka gave 3.6 and 3.7 number of roots/plant while Abees gave 3.8 and 3.6. These results were in agreement with those of Sheng and Wang (1992), Reddy *et al* (1996), Salem (1999) and Gendy (2002) who found significant differences among the sweet potato lines and cultivars.

**Average tuber root weight (g)**

Data concerning average tuber root weight are shown in Table (4). Significant differences for this trait were observed among the studied lines. Average tuber root weight showed the highest values of 870.0 and 724.1g in the hybrid No (2) and (4) in the first season while they were 684.6 and 685.0g in the second season . The hybrid No (3) gave the lowest values in the first and second season with means of 362.0 and 352.5 g respectively. These hybrids exceeded the best check cultivar Mabrouka which gave 131.0g in the second season. The percentage of increase in these lines compared with Mabrouka cvs was 169% . However, the tuber roots in this study were significantly heavier than the check cvs. It is very good result when some hybrids outyielded the commercial cultivars Mabrouka and Abees cvs. These similar results were reported by other researchers among of them El-Gazzar (1988), El-Shini (1996), Gendy (2002) and Vimalan and Hariprakash (2011), who found significant differences among the studied sweet potato lines and cultivars for average tuber root weight.

**Heterosis**

**Average degree of heterosis :**

Average degree of heterosis (ADH %) was expressed as percent increase or decrease of each F1 population above the standered cultivars. The average degree of heterosis in relation to the standered cultivars was estimated for all traits.

**Number of branches /plant**

Data presented in Table (5) show significant positive ADH% heterosis values based on the stander cultivars in the all studied crosses suggesting dominance towards the large number of branches /plant. These results are similar to those obtained by Salem (1999) who found over dominance for the large number of branches in sweet potato.

**Table(5).Average degree of stander heterosis (ADHS) based on cultivars Mabrouka and Abees for number of branches /plant, stem length (cm) and Dry matter% of six sweetpotato hybrids.**

Genotypes	Mabrouka						Abees					
	Number of branches/plant		Stem length (cm.)		Dry matter (%)		Number of branches/plant		Stem length (cm.)		Dry matter%	
	2010	2011	2010	2011	2010	2011	2010	2011	2010	2011	2010	2011
H .No1	70.0**	52.3**	-30.90**	-30.90**	43.19**	43.77**	25.0**	28.0**	-35.2**	-36.3**	35.53**	33.44**
H .No2	50.0**	33.3**	-27.01**	-27.01**	54.49**	58.59**	11.0*	12.0*	-31.6**	-27.2**	46.22**	47.19**
H .No3	30.0**	14.2*	-41.90**	-41.90**	47.18**	51.85**	3.7 <sup>ns</sup>	4.0 <sup>ns</sup>	-45.5**	-41.7**	39.31**	40.94**
H .No4	40.0**	20.0**	-39.70**	-39.70**	42.52**	47.14**	3.7	36.0**	-43.5**	-40.6**	34.91**	36.56**
H .No5	75.0**	61.0**	-23.20**	-23.20**	50.17**	51.18**	29.0**	0.0	-27.9**	-24.5**	42.14**	40.31**
H .No6	80.0**	66.0**	-33.10**	-33.10**	63.46**	62.63**	33.0**	40.0**	-37.3**	-32.6**	54.72**	50.94**



**Stem length**

Average degree of standered hetrosis (ADHs) based on standered cultivars Mabrouka and Abees for stem length cm /plant of six sweet potato genotypes. This characters recorded that all the values of this trait were negative heterosis values means that stem length( cm) in the two varieties Mabrouka and Abees had longer stems than the hybrids in the two seasons 2010 and 2011 respectively, this results referres to that theres relation between yield/plant and stem length in sweet potato. Data present in Table (5) show significant negative heterosis values in relation to the standered cultivars. These ADH values suggested partial dominance for the short stem length in these crosses. The obtained results are similar to those reported by Gendy (2002) who found that partial dominances was detected for stem length in sweet potato plants.

**Dry matter%**

Data present in Table (5) show significant positive ADH% heterosis values based on the standered cultivars in the all studied crosses suggesting dominance towards the dry matter%.These results are similar to those obtained by Cervantes-Flores *et al.* (2010) and Placide *et al.* (2013).

**Total yield:**

**Total yield/ plant**

Data presented in Table (6) show that the heterosis percentages relative to standered cultivars ranged from 110.0 to362.0% in the first season and ranged from 111 to 309% in the second season. The hybrid No (1) gave the highest yield than the two check cultivars Mabrouka and Abees suggesting dominance towards the high cultivars. Generally, it could be concluded that the F1 crosses in sweet potato had hybrid vigour regarding total tuber root yield. Then the utilization of F1 hybrids in sweet potato cultivations will be lead to increase the yield production these results were agree with those of Salem (1999) and Vimalan and Hariprakash (2011), who found dominance for the high total tuber yield in their studies.

**Table (6).Average degree of standered hetrosis for yield/plant and root weight from some hybrids of sweet potato in 2010 and 2011 seasons.**

Genotypes	Mabrouka				Abees			
	Yield/ plant		Root weight		Yield/ plant		Root weight	
	2010	2011	2010	2011	2010	2011	2010	2011
H .No1	362.0**	292.0**	316.0**	271.0**	418.0**	347.0**	388.0**	312.0**
H .No2	319.0**	267.0**	574.0**	422.0**	369.0**	318.0**	690.0**	480.0**
H .No3	110.0**	111.0**	180.0**	169.2**	136.0**	141.0**	229.0**	480.0**
H .No4	352.0**	309.0**	461.0**	422.0**	406.0**	367.0**	558.0**	480.0**
H .No5	190.0**	168.0**	237.0**	241.0**	225.0**	205.0**	286.0**	279.0**
H .No6	254.0**	268.0**	299.0**	304.0**	297.0**	311.0**	368.0**	349.1**

**Root weight**

Estimates of ADH% in relation to standered cultivars show that all the hybrids gave significant positive hetrosis values suggesting domanince towards the high average degree tuber root weight. Obtained hetrosis values in the 6 crosses ranged from 180.0 to 574% in the first season and ranged from 169.2 to 422.0% in second season compared with Mabrouka cv. While

they ranged from 229.0 to 558.0% in the first season and from 279 to 480% in the second season compared with Abees cultivar. These significant positive heterosis values suggesting dominance toward the high average tuber root weight and the hybrid vigours verified by the obtained significant positive heterosis values. The obtained results are similar to those reported by Chaurasiya *et al* (2013) found that the gene action studies indicated that there was preponderance of non-additive (dominance) gene action for all the traits under study.

## REFERENCES

- Acquaah, G (2007). Principles of plant genetics and breeding Wiley-Blackwell, Oxford .
- Bourke, R.M (1989). Influence of soil moisture extremes on sweet potato yield in the Papua New Guinea highlands. Mountain Res. Dev., 9(3):322-328.
- Cervantes-Flores, J.; B. Sosinski; K. Pecota; R .Mwanga; G .Catignani; V. Truong; R. Watkins; M. Ulmer and G .Yencho (2010). Identification of quantitative trait loci for dry matter, starch and  $\beta$ -carotene content in sweet potato. Mol Breeding, 5: 1-16.
- Chalfant.R.B; R.K. Gaasson and R.F. Dokshina (1990). Ecology and management of sweet potato insects. Ann. Ref. Entomol., 5:157-180.
- Chaurasiya P. C.; J. Singh; S. N. Dikshit and N. Mehta (2013). Heterosis, Combining Ability and Gene Action Studies in Sweet potato (*Ipomoea batatas* L.) Lam. International J. of Plant Res., 26: 264-270.
- Chiona, M (2009). Towards enhancement of  $\beta$ -carotene content of high dry mass sweet potato genotypes in Zambia, African Centre for Crop Improvement (ACCI), School of Agricultural Sciences and Agribusiness Faculty of Science and Agriculture, KwaZulu-Natal, Pietermaritzburg
- EL-Denary, M.E.M. (1998). The performances of sweet potato (*Ipomoea batata* L.) plant in response to some cultural treatments .M.Sc. of Hort. Sci, vegetable crops. Foc. of Agric, Monoofeiya univ. Egypt.
- EL-Gazzar .T.M. (1988). The isolation of high yield clone of sweet potato (*Ipomoea batata* L.) Balady through spontaneous mutation J.of Agric. Sci. Mansoura. Univ.Egypt, 13 (1):461-470.
- El-Shini, A.A.M. (1996). Effect of some factors on the production and quality of sweet potato and its storage ability Ph.D. Thesis . Agric cairo Univ .Egypt.
- Fehr, W.R. (1987). IN: Principles of cultivar development. Vol. MacMillan Publishing Company, USA.
- Gasura E, A. Mashingaidze and S. Mukasa (2010). Genetic variability for tuber yield, quality, and virus disease complex traits in Uganda sweet potato germplasm. Afr Crop Sci J., 16: 147-160.
- Gasura E., A.B. Mashingaidze and S.B. mukasa (2008). Genetic variability for tuber yield, quality, and virus disease complex traits in Uganda sweet potato germplasm. African Crop Science J., 16 (2):147 – 160.
- Gendy, A.S.S. (2002). Genetical studies on sweet potato Ph.D. Thesis of Hort. Sci. Fac. Agric, Menoufiya Univ. Egypt.
- Huaman.Z (1991). Descriptors for sweet potato. Cip, Avrdc, Ibrgr, pp. 134.

- Huaman.Z. T.T. Williams; W.Salhuana and L. Vincent (1977). Descriptors for the cultivated potato and for the maintenance and distribution of germplasm collection. Int. Board for Plant Genetic Resources, Rome, Italy, p.47. (C.A. Computer Search ).
- Jones A (1965). A proposed breeding procedure for sweet potato. *Crop. Sci.*, 5: 191-192.
- Jones A (1966). Morphological variability in early generations of randomly intermating populaton of Sweet potato (*Ipomoea batatas* (L) Lam.) Uni. Ga. Agr. Expt. Sta. Tech.Bull.N.S., 56:31.
- Jones A, Steinbauer CE, Pope DT (1969). Quantitative inheritance of ten root traits in Sweet potato.*J. Amer. Soc. Hort. Sci.*, 94: 271-275.
- Jones,A. (1977). Heitabilities of seven sweet potato root traits. *J. Amer. Soc.Hort.Sci.*, 102 (4):440-442.
- Jones, A., P.D. Dukes and J.M. Schalk ( 1987). Sweetpotato breeding. Breeding vegetable crops. AVI Publishing Co, Washington, D.C. p. 1-35.
- Ma, D-f.;Q. Li; X-y. Li; H-m. Li; Z-h. Tang; L. Hu; Q-h. Cao; Y-p. Xie and X. Wang (2009). Selection of parents for breeding edible varieties of sweet potato with high carotene content. *Agr Sci China*, 8: 1166-1173.
- Miller, Jc (1939) Further studies and technic used in sweetpotato breeding in Louisiana. *J Heredity*. 30: 485-492
- Morrison.T.A.R. Pressey and S.J.Kaks (1993). Change sin Alpha and Beta amylase during storage of sweet potato lines with varying starch hydrolysis potential. *J. Amer. Soc. Hort. Sci.*, 118 (2) :236-242.
- Mwanga Z.; M. Mataa and M. Msabaha (2007). Quality and yield stability of orange fleshed sweetpotato (*Ipomoea batatas*) varieties grown in different agro-ecologies. *Afr Crop Sci. J.*, 8: 339-345.
- Mwanga. R.; B. Odongo; C. Niringiye;R. Kapinga; S .Tumwegamire; P. Abidin; E. Carey; B. Lemaga; J. Nsumba and D. Zhang (2010). Sweet potato selection releases: Lessons learnt from Uganda. *Afr Crop Sci J.*, 15: 11-23.
- Placide, R .; S.Hussein ; L.Mark and G.Daphrose (2013). Storage root formation, dry matter synthesis, accumulation and genetics in sweet potato *AJCS*, 7(13):2054-2061 .
- Reddy,B.S;A.V.Desai;M.G.Patil;S.A.Hussain;P.N.Reddy; N.G.Hulaman and P.M.Gangadharappa (1996). Evaluation of sweet potato genotypes under semiarid condition of Raichur. *Advances in Agric.Res. in India*.5:33-36.
- Salem, A.A.M. (1999).Genetical and physiological studies on potato. PhD. Thesis of Hort. Sci. Fac. Agric., Menufiyal Univ.
- Shalaby,G.I.; S.A. Abdel AAL; A.M Damarany and A.M.Abd Ei-Salam (1993a). Evaluation of some cultivars and breeding lines of sweet potato under Assiut conditions.I. Vine and Storage Root Yield Data. *Assiut J. Agric. Sci.*, 24(1): 315-328.
- Shalaby,G.I.; S.A. Abdel AAL; A.M Damarany and A.M.Abd Ei-Salam (1993b). Evaluation of some cultivars and breeding lines of sweet potato under Assiut conditions.2. Chemical composition and storage ability of storage roots. *Assiut.J. Agric.Sci.*, 24(1):329-344.

- Sheng, J. and Q. Wang (1992). New sweet potatoes of multipurpose type. Crop Genetic Resources No. 346. (C.A.Plant Breed.Abstr., 8881).
- Singh, P. and S.S. Narayanan (1993). Biometrical Techniques in Plant Breeding. Kalyani Publishers, New Delhi, India, pp: 74-84.
- Snedecor, G. W and W. G. Cochran (1972). Statistical methods applied to experiments in agriculture and biology Iowa State College USA
- Suzuki, M. (1990). New summer crop varieties registered by the Ministry of Agriculture, Forestry and Fisheries. Paddy Rice; sweet potato, Soybean and sugar-cane. Japanese. J. of Breeding, 40(4).537-547.
- Vimalan B and Hariprakash B (2011). Variability of morphological characters and dry matter content in the hybrid progenies of sweetpotato (*Ipomoea batatas* L.) Lam.]. Geneconserve 10: 65-86
- Vimalan, B. (1993). Genetic studies of Sweet potato (*Ipomoea batatas* L.) Lam.)- A Review. J. Root crops. 19 (1) : 40.46.
- Wanas, N.M. (1979). Studies on some factors affecting the quantity and quality of sweet potato.M.Sc. Thesis, Fac. Agric. Menoufeiya Univ.Egypt.

### قوة الهجين ومكونات المحصول لبعض هجن F1 من البطاطا عبد المنعم سيد احمد الجندى و أحمد جمعة محمد سليمان قسم بحوث الخضر - معهد بحوث البساتين - مركز البحوث الزراعية - الجيزة

أجريت هذه الدراسة بمركز مطوبس كفر الشيخ خلال الفترة من عام ٢٠١٠ حتى ٢٠١١ حيث تهدف الدراسة إلى إنتاج هجن متفوقة في المحصول ومكوناته عن أهم وأشهر صنفين تجاريين من البطاطا لدى المزارعين مبروكة و ابيس وذلك تحت ظروف العروة الشتوية وتم الزراعة الست هجن والصنفين التجاريين المحليين في موسمين في أربعة مكررات وتم تصميم التجربة في قطاعات كاملة العشوائية . ويمكن تلخيص أهم النتائج فيما يلي :-

- ١- أظهرت النتائج وجود اختلافات معنوية بين الهجن المستخدمة والأصناف التجارية وكذلك مع بعضها البعض .
- ٢- تفوقت الهجن رقم ١ ورقم ٣ عن صنف مبروكة بزيادة ١٦٨% زيادة محصولية و ٥١.٥% على التوالي.
- ٣- وحقق الهجن زيادة محصولية ١٧٣% و ٥٦% عن الصنف التجاري ابيس.
- ٤- الهجين الرابع حقق زيادة معنوية عالية فى وزن الجذر الواحد مقارنة بصنفي المقارنة.
- ٥- سجلت التراكيب الوراثية درجتان للون اللحم هم القرمزي والقرمزي الداكن.
- ٦- سجل لون اللحم الداخلي ست درجات مختلفة من الأبيض حتى القرمزي .
- ٧- كذلك سجلت الدراسة زيادة معنوية في معظم الصفات المدروسة مقارنة بصنفي المقارنة
- ٨- أثبتت دراسة قوة الهجين وجود سيادة تامة وجزئية تجاه معظم الصفات.