



EFFECT OF GARLIC AS FEED ADDITIVE ON PERFORMANCE, CARCASS CHARACTERISTICS, AND MEAT QUALITY OF MUSCOVY DUCKS REARED IN DIFFERENT STOCKING DENSITIES

Basma M. Bawish , Fayed, R.H*., and Abeer H. Abdel Razek

Department of Veterinary Hygiene and Management, Fac. Vet. Med., Cairo University, Egypt

*Corresponding author: Fayed, R.H., email: rhfayed@cu.edu.eg

ABSTRACT

This study was carried out to evaluate the effect of supplementing diet with different garlic powder concentrations (300 and 600gm/Ton) on performance, carcass characteristics and meat quality (fat %) of Muscovy ducks exposed to different stocking densities; standard stocking density (4 birds/m²) and high stocking density (8 birds/m²). A total number of 180 one -day -old Muscovy ducklings were used in this experiment. Birds were randomly allocated into 18 symmetrical pens classified into 6 treatment groups (3 replicate / treatment). Groups (C, SS+G₃₀₀, SS+G₆₀₀) raised on standard stocking density and fed on basal diet supplemented with (0 , 300 and 600 g. garlic powder,/ton feed) respectively while groups (HS, HS+G₃₀₀, HS+G₆₀₀) raised on high stocking density and fed on basal diet supplemented with (0 , 300 and 600 g. garlic powder/ton feed) respectively. The data of growth performance indicated that supplementation of garlic (600 gm / ton) significantly ($p > 0.05$) improved performance (body weight gain and FCR) of ducks in standard stocking density (SS+G₆₀₀) group and high stocking density in (HS+G₆₀₀) group. Concerning carcass characteristic and meat quality, results indicated that, garlic supplementation reduced fat % in thigh and breast muscles. It can be concluded that, adding garlic powder to duck ration can effectively compensate the negative effects of the high stocking density in Muscovy ducks.

Keywords: Carcass characteristics, garlic, Muscovy ducks, performance, stocking density.

Original Article:

DOI:<https://dx.doi.org/10.21608/javs.2018.67162>

Received: 02 june, 2018.

Accepted: 28 June, 2018.

Published: 12 July, 2018.

This is an open access article under the term of the Creative Commons Attribution 4.0 (CC-BY) International License . To view a copy of this license, visit:

<http://creativecommons.org/licenses/by/4.0/>

INTRODUCTION

World duck production has increased considerably since 1993, and meat production has risen from 1.72 to 4.34 million tons (Demian, 2014). Ducks are capable of increasing the much needed animal protein on account of their reputation for fast growth and efficient feed conversion (Ahaotu and Aghasu (2015). The performance and welfare of ducks are affected by different factors like environment, nutrition, management, and genetic makeup (Estevez, 2007). Stocking density is an important factor affecting duck production and welfare, but economic profit may come at the cost of reduced bird performance, health, and welfare if densities are excessive (Xie. et al .,2014), as high stocking density led to physical restriction of birds movement as locomotors behaviour so effect the access of birds to feed and water (Estevez, et al., 1997 and SCAHAW, 2000), decrease growth

performance (Simitzis et al., 2012; Tong et al., 2012), carcass traits (Dozier et al., 2005 and Dozier et al., 2006) , meat quality (Osman, 1993) . But a higher profitability per kilogram of poultry can be obtained by increasing the stocking density if the performance of birds remains constant (Feddes et al., 2002).

The utilization of plants of natural origin (phytobiotics) as growth promoters considered a practical and achievable way in recent years (Grashorn, 2010; Yang et al., 2015). A large variety of plant compounds (phytobiotics) such as garlic were used as growth promoters (Farhad et al., 2011 and Ismoyowati et al., 2015) .In poultry, garlic has been shown to improves growth and Feed Conversion Ratio (FCR), and decreased mortality rate (Lewis et al., 2003 and Tollba and Hassan ., 2003) and Bampidis, et al., 2005) also induce hypocholesterolemia (Yeh and Liin 2001 and Yoo et al., 2011) which important in

improving the fat composition and overall appearance of duck meat (**Dwiloka., et al 2015**) which can provide more benefits for human health.

Researches about the effect of garlic on productive performance and carcass characteristics of Muscovy ducks on different stocking density was scarce. Therefore, the current study was conducted to assess the effects of different concentration of garlic powder supplementation on productive performance, carcass characteristics, meat quality (fat %) of Muscovy ducks in different stocking density.

MATERIALS AND METHODS

Birds and housing

The present study was carried out at the poultry research unit at the Department of Veterinary Hygiene and Management, Faculty of Veterinary Medicine, Cairo University. A total number of 180 one day old Muscovy ducks (*Carina moschata*) of both sexes were divided into 6 groups in 3 replicates of 10 ducks in standard stocking density groups and 20 duck in high stocking density group. All ducks were weighed before the start of experiment to take the initial weight then brooded at about 35 °C using gas heater during the 1st week of brooding period then gradually reduced 3 °C / week till reaching 25 °C at 4th week of age till the end of the study, Continuous lighting program of 23 h light and 1 h dark was maintained by the natural day light and continued by artificial light throughout the experimental period and good ventilation was maintained through the experiment. (**Coates et al., 2000**).

The experimental birds were reared in floor litter system with wood shaving (10 cm depth) at separate pens of identical size (3 m length x 2.50 m width x 3 m height) which was thoroughly cleaned, washed and disinfected before arrival of duckling; Fresh clean water and feed was supplied ad libitum throughout the experimental period. The ducks were fed A starter diet of (23.34 CP and 3020.80 kcal/kg ME) that have been used from 0–4 weeks while grower diet of (19.26 CP and 3250.37 kcal/kg ME) were fed from 5 - 10 weeks (end of trials). The basal starter and grower diets were formulated to meet the nutrient requirements of Muscovy ducks according to the recommendation of the national research council (**NRC, 1994**).

Experimental design

Birds were randomly allocated into 18 symmetrical pens classified into 6 groups and each group was represented by three replicates / group. The 1st group fed on basal diet with standard stocking density control (C) , the 2nd group fed on basal diet

with high stocking density (HS), the 3rd group fed on basal diet with 300 g. garlic powder / ton feed with standard stocking density (SS+G₃₀₀), the 4th group fed on basal diet with 300 g. garlic powder / ton feed with high stocking density (HS+G₃₀₀), the 5th group fed on basal diet with 600 g. garlic powder / ton feed with standard stocking density (SS+G₆₀₀), the 6th group fed on basal diet with 600 g. garlic powder / ton feed with high stocking density (HS+G₆₀₀).

Parameters measured

Performance Parameters

A random sample of 5 birds from each replicate were weighted individually every week using large- pan beam balance to obtain the average body weight then average weekly weight gain was calculated through the following equation:

$$\text{Average weight gain} = (W_2 - W_1)$$

Where:

W₂= average body weight in a particular week

W₁ = a average body weight in a previous week

The feed consumption was calculated weekly for each group throughout the experimental period. Feed conversion was calculated weekly by dividing the average feed consumption / ducks / week on the average body weight gain / ducks / week. (**Sheng-Qiu et al., 2013**). The mortality rate was recorded daily throughout the experimental period.

Carcass and organ characteristics

At the end of the experiment, Feed was withdrawn 12 h before slaughter and 5 birds from each replicate were taken randomly and slaughtered for their dressing percentage and organ weight (heart, liver, and gizzard). (**Ahaotu and Agbasu, 2015**).

Meat quality measurements (fat %)

At the end of the experiment three birds from each replicate were randomly chosen and were slaughtered by severing of jugular vein then thigh (Biceps femoris) and breast (Pectoralis major) muscle samples were taken from each carcass and then analyzed individually to determine the fat percentage of the breast and thigh muscles (**Galal., et al ., 2011**) by Soxhlet method. (**AOAC, 1995**).

Statistical analysis

Parametrical statistical test were applied (ANOVA) after exploration of data. Descriptive statistics are represented by means ± standard error (SE). The least significance difference (LSD) was used for significance among groups at significance level (p < 0.05) **IBM CORP (2013)**.

RESULTS

Table 1: The effect of different garlic powder concentrations on body weight of, Muscovy ducks exposed to different stocking density

Gps.	(C)	HS	SS+G ₃₀₀	HS+G ₃₀₀	SS+G ₆₀₀	HS+G ₆₀₀
Day 1	51.67±0.29	52.19±0.35	52.13±0.30	51.16±0.15	52.17±0.39	52.44±0.37
W1	141.67±1.65 ^a	134.28±0.94 ^b	144.96±1.24 ^a	137.50±0.83 ^b	150.33±1.70 ^c	137.16±1.95 ^b
W2	321.58±4.01 ^{ad}	306.00±2.17 ^b	329.75±4.17 ^a	311.00±1.85 ^{bd}	359.50±4.74 ^c	314.69±1.35 ^d
W3	591.25±9.66 ^{ad}	553.13±2.20 ^b	603.08±6.29 ^a	562.13±9.31 ^{bd}	654.42±4.75 ^c	580.94±8.37 ^d
W4	934.58±11.11 ^a	854.06±8.54 ^b	948.33±13.32 ^a	872.81±11.96 ^b	1019.1±12.71 ^c	915.94±10.78 ^a
W5	1395.0±6.71 ^a	1274.0±13.25 ^b	1416.2±10.22 ^a	1307.1±8.54 ^c	1495.0±13.05 ^d	1371.8±9.24 ^a
W6	1864.1±25.48 ^a	1716.2±13.85 ^b	1889.1±10.66 ^a	1755.4±9.55 ^b	1990.8±9.54 ^c	1835.0±7.50 ^a
W7	2311.6±7.82 ^a	2125.9±7.89 ^b	2345.4±14.35 ^c	2170.0±6.43 ^d	2461.2±7.41 ^e	2278.13±4.82 ^f
W8	2681.6±14.27 ^a	2476.8±13.58 ^b	2720.8±9.48 ^c	2534.6±9.32 ^d	2860.4±12.74 ^e	2645.9±12.89 ^f
W9	3020.0±18.74 ^a	2759.3±14.62 ^b	3068.7±13.98 ^c	2827.1±12.32 ^d	3225.0±7.04 ^e	2980.6±12.06 ^f
W10	3348.1±24.37 ^a	3026.8±27.88 ^b	3400.4±17.95 ^a	3100.9±25.37 ^c	3570.8±7.49 ^d	3306.8±20.48 ^a

(C): group fed on basal diet with standard stocking density. (HS): fed on basal diet with high stocking density.

(SS+G₃₀₀): fed on basal diet with 300 g. garlic powder / ton with standard stocking density.

(HS+G₃₀₀): fed on basal diet with 300 g. garlic powder / ton with high stocking density.

(SS+G₆₀₀): fed on basal diet with 600 g. garlic powder / ton feed with standard stocking density .

(HS+G₆₀₀): fed on basal diet with 600 g. garlic powder / ton with high stocking density.

^{Abc}: Different small alphabet within the same row denotes statistically difference at 0.05. Similar letters denote there was no statistical significant difference

Table 2: The effect of different garlic powder concentrations on body weight gain of Muscovy ducks exposed to different stocking density

Gps.	(C)	(HS)	(SS+G ₃₀₀)	(HS+G ₃₀₀)	(SS+G ₆₀₀)	(HS+G ₆₀₀)
W1	90.00±0.76 ^a	82.09±0.47 ^b	92.83±0.97 ^c	86.34±0.70 ^d	98.17±0.78 ^e	84.75±0.46 ^d
W2	179.92±0.55 ^a	171.72±0.45 ^b	184.79±1.18 ^c	173.50±0.86 ^d	209.17±1.11 ^e	177.53±0.57 ^a
W3	270.08±2.52 ^a	247.13±1.28 ^b	273.33±2.41 ^{ac}	256.75±1.88 ^c	294.92±1.21 ^d	266.25±2.21 ^e
W4	345.42±2.08 ^a	300.94±3.46 ^b	355.58±6.01 ^{ac}	310.69±2.24 ^b	364.75±3.26 ^c	335.00±2.82 ^d
W5	460.42±3.56 ^a	420.00±1.44 ^b	476.67±3.00 ^c	434.38±1.99 ^d	475.83±3.00 ^c	455.94±6.63 ^a
W6	469.17±1.53 ^{ac}	442.19±2.48 ^b	472.92±2.82 ^a	448.75±3.93 ^b	495.83±2.63 ^c	463.13±2.82 ^e
W7	447.50±2.27 ^a	409.69±2.13 ^b	456.25±1.72 ^c	414.06±3.24 ^b	470.42±2.45 ^d	443.13±2.54 ^a
W8	370.00±1.90 ^a	350.94±1.85 ^b	371.44±3.78 ^a	364.69±2.96 ^a	399.08±4.12 ^c	367.82±1.58 ^a
W9	338.34±2.27 ^a	282.50±1.73 ^b	351.53±6.71 ^c	289.83±3.93 ^b	364.48±2.45 ^d	334.69±3.51 ^a
W10	328.08±3.49 ^a	267.50±1.99 ^b	331.67±3.00 ^a	272.82±5.18 ^b	345.38±2.27 ^c	326.25±5.32 ^a

Different small alphabet within the same row denotes statistically difference at 0.05. Similar letters denote there was no statistical significant difference

Effect Of Garlic As Feed Additive On Performance

Table 3: The effect of different garlic powder concentrations on weekly feed intake of Muscovy ducks exposed to different stocking density

Gps.	(C)	(HS)	(SS+G ₃₀₀)	(HS+G ₃₀₀)	(SS+G ₆₀₀)	(HS-G ₆₀₀)
W1	105.00±2.56 ^a	98.04±2.65 ^{ab}	103.11±2.61 ^{ab}	96.81±2.85 ^{bc}	95.31±2.83 ^{bc}	89.43±2.88 ^c
W2	269.40±2.56 ^a	256.97±2.60 ^{bc}	261.31±2.58 ^c	253.40±2.63 ^b	259.25±2.73 ^{bc}	242.24±2.84 ^d
W3	474.73±2.83 ^{ac}	465.15±4.04 ^b	466.31±3.20 ^{ab}	464.49±4.35 ^b	482.23±2.90 ^c	457.63±4.18 ^b
W4	793.29±4.37 ^a	750.61±4.04 ^b	782.58±3.01 ^c	746.31±3.20 ^b	773.47±3.44 ^c	733.04±2.98 ^d
W5	1107.58±4.05 ^a	1060.85±2.64 ^b	1094.70±2.60 ^c	1051.86±2.56 ^d	1078.00±2.50 ^e	1048.43±2.93 ^d
W6	1145.97±3.00 ^a	1121.61±4.04 ^b	1140.65±3.40 ^{ac}	1092.81±2.58 ^d	1132.11±3.00 ^c	1087.91±4.50 ^d
W7	1181.11±3.00 ^a	1128.30±3.89 ^b	1166.41±2.81 ^c	1121.54±2.56 ^{bc}	1149.63±2.55 ^d	1114.86±3.91 ^e
W8	1212.54±4.29 ^a	1161.27±5.20 ^b	1209.92±2.99 ^a	1145.97±4.15 ^c	1183.81±2.58 ^d	1135.61±4.04 ^c
W9	1221.48±4.21 ^a	1177.23±2.61 ^b	1221.09±3.58 ^a	1169.81±2.58 ^b	1211.81±2.58 ^c	1168.58±2.85 ^b
W10	1246.81±2.58 ^a	1206.31±2.56 ^b	1237.39±3.40 ^c	1198.54±4.07 ^b	1229.69±3.31 ^c	1188.18±3.01 ^d

Table 4: The effect of different garlic powder concentrations on FCR of Muscovy ducks exposed to different stocking density

Gps.	(C)	(HS)	(SS+G ₃₀₀)	(HS+G ₃₀₀)	(SS+G ₆₀₀)	(HS+G ₆₀₀)
W1	1.17±0.03 ^a	1.19±0.03 ^a	1.11±0.02 ^{ac}	1.18±0.07 ^a	0.97±0.03 ^b	1.05±0.03 ^{bc}
W2	1.50±0.01 ^a	1.50±0.01 ^a	1.41±0.02 ^b	1.51±0.02 ^a	1.24±0.01 ^c	1.36±0.01 ^d
W3	1.76±0.01 ^a	1.88±0.01 ^b	1.71±0.02 ^c	1.82±0.03 ^d	1.64±0.01 ^e	1.72±0.01 ^{ac}
W4	2.30±0.02 ^a	2.50±0.04 ^b	2.20±0.04 ^c	2.40±0.01 ^d	2.12±0.03 ^c	2.19±0.02 ^c
W5	2.41±0.01 ^a	2.53±0.01 ^b	2.30±0.01 ^c	2.42±0.01 ^a	2.27±0.02 ^c	2.30±0.03 ^c
W6	2.44±0.01 ^a	2.54±0.01 ^b	2.41±0.01 ^a	2.44±0.02 ^a	2.28±0.01 ^c	2.35±0.02 ^d
W7	2.64±0.02 ^a	2.75±0.01 ^b	2.56±0.01 ^c	2.71±0.02 ^b	2.44±0.01 ^d	2.52±0.02 ^c
W8	3.28±0.01 ^a	3.31±0.03 ^a	3.26±0.03 ^a	3.14±0.02 ^b	2.97±0.03 ^c	3.09±0.01 ^b
W9	3.61±0.03 ^a	4.17±0.03 ^b	3.48±0.07 ^c	4.05±0.05 ^d	3.32±0.02 ^e	3.49±0.03 ^{ac}
W10	3.80±0.04 ^a	4.51±0.03 ^b	3.73±0.03 ^{ad}	4.40±0.09 ^b	3.56±0.02 ^c	3.65±0.06 ^{cd}

Table 5: The effect of different garlic powder concentrations on carcass characteristics of Muscovy ducks exposed to different stocking density

Groups	(C)	(HS)	(SS+G ₃₀₀)	(HS-G ₃₀₀)	(SS+G ₆₀₀)	(HS+G ₆₀₀)
Dressing %	77.50±0.39 ^a	76.27±0.16 ^b	78.03±0.38 ^a	76.44±0.25 ^b	78.23±0.62 ^a	77.37±0.34 ^a
Gizzard weight	81.32±5.80	71.19±2.65 ^a	83.62±6.58 ^a	74.87±2.56 ^a	87.25±4.22 ^a	80.10±2.66 ^a
Liver weight	72.95±4.69 ^a	62.85±2.64 ^a	73.39±4.43 ^a	64.86±3.24 ^a	77.07±4.73 ^a	69.23±3.11 ^a
Heart weight	17.75±1.36 ^a	15.23±0.49 ^a	18.24±1.72 ^a	15.54±0.60 ^a	19.06±1.62 ^a	16.71±0.69 ^a

Table 6: The effect of different garlic powder concentrations on meat quality (fat %) of Muscovy ducks exposed to different stocking density

Groups	(C)	(HS)	(SS+G ₃₀₀)	(HS+G ₃₀₀)	(SS+G ₆₀₀)	(HS+G ₆₀₀)
Breast muscle fat %	1.56±0.04 ^a	1.52±0.12 ^{ab}	1.44±0.07 ^{ab}	1.26±0.12 ^b	0.85±0.11 ^c	0.71±0.07 ^c
Thigh muscle fat %	2.44±0.48 ^a	2.37±0.17 ^a	2.22±0.10 ^a	2.17±0.13 ^{ab}	1.46±0.09 ^{bc}	1.49±0.18 ^b

DISCUSSION

Body weight and body weight gain

Results in Table (1) showed that the body weight of muscovy ducks significantly increased ($p < 0.05$) gradually from 1st week till the end of the experiment at 10th week of age where SS+G₆₀₀ group showed significantly ($p < 0.05$) higher final body weight however the final body weight of (HS+G₆₀₀) group not significantly ($p < 0.05$) different from control group and (SS+G₃₀₀) group, on the other hand (HS+G₃₀₀) group and (HS) group showed the lower final body weight .The significant decrease in live body weight throughout the experiment due to increasing of stocking density as in (HS) group and (HS+G₃₀₀) group may be attributed to the decrease in feed intake and nutrient utilization at all growing age which not allowing them to grow to their potential (Taboosha ,2014) our findings are in agreement with some previous evaluations which showed a linear decrease in BW with increasing population density (Dozier *et al.*, 2005, Xie *et al.*, 2014 and Ahaotu and Agbasu (2015) but not in agreement with Feddes *et al.*, (2002) who reported similar body weight of broiler reared at densities of 12, 18, and 24 birds/m² and Buijs *et al.*, (2009) who found no difference in final BW at 39 d of age as stocking density increased.

However, on the other hand supplementation of garlic (600 gm / ton) increase the body weight of ducks in standard stocking density as in (SS+G₆₀₀) and could effectively compensate the decrease in BW under high stocking density as in (HS+G₆₀₀) due to the growth promoting effect of garlic which be attributed to allicin (an organosulfur compound) contained in garlic that promotes the performances of the intestinal flora thereby enhance digestion which increase body weight of ducks (Pourali *et al.*, 2010) , these results in line with El-Ghamry *et al.*, (2004) who reported that addition of fresh garlic 2% could enhance the productive performance and improved body weight of Muscovy ducks and also with (Brzóska *et al.*, (2015) , El-katcha *et al.*, (2016) but in contrary with studies of Ghazalah and Ibrahim (1996) who mentioned that supplementation of garlic oil (1.25 mg/100g body weight) reduce body weight in Muscovy ducks and

with Ismoyowati *et al.*, (2015) who found that there was no significant effects of inclusion of 1 % of garlic on B.W of ducks during the growing period.

Concerning with body weight gain as in Table (2), it was obvious that BWG significantly increased ($p \leq 0.05$) till reach maximum value at 6th week of age and then decreased gradually till the end of the experiment, the highest BWG was recorded in (SS+G₆₀₀) group followed by (SS+G₃₀₀) group, (C) group, (HS+G₆₀₀) group while the lowest BWG was recorded in (HS) group then (HS+G₃₀₀) group respectively. These results, showed that ducks housed at high stocking density as in (HS group) and (HS+G₃₀₀) had lower body weight gain which may be explained through decreasing of time spent resting and increase disturbance in the groups of high stocking density (Febrer *et al.*, 2006) which enable the birds to gain less body weight than those of lower stocking density, it agreed with (Taboosha ,2014 , Xie *et al.*, (2014 and Cengiz *et al.*, 2015) who showed that the final BWG decreased significantly when stocking density increased but in contrast with (Ravindran *et al.*, 2006) who found no differences in average weight gain between the 3 density treatments (16, 20, and 24 birds/m²),

Regarding garlic supplementation (600gm /ton feed), increase body weight gain in high stocking density group (HS+G₆₀₀) group and led the highest body weight gain in standard stocking density group (SS+G₆₀₀) group which may be due to the fact that allicin (active principle of garlic) control of growth and colonization of various pathogenic microorganisms in the gut of birds so when the load of these bacteria in the intestine is low, birds may absorb more nutrients, thus leading to the improvement in weight gain of the birds (Bedford , 2000) , these results agreed with Bidura (1999) who demonstrated an increase in live body weight gain of growing duck aged 0 – 8 weeks compared to control one when diet supplemented with 3% and 6 % garlic leaf and also with (Bampidis, *et al.*, 2005 and Abd El Azim *et al.*, 2009) while disagreed with Ismoyowati *et al.*, (2015) who found that weight gain of ducklings were not affected by inclusion of 1 % of garlic .

Feed intake and feed conversion ratio (FCR)

The results in Table (3) showed that the weekly feed intakes significantly ($p \leq 0.05$) increased gradually from 1st week till the end of the experiment where groups of high stocking density showed lower feed intake with the lowest mean recorded in (HS+G₆₀₀) group followed by (HS+G₃₀₀) group and (HS) group,. On the other hand the higher feed intake observed in groups of standard stocking density but (SS+G₆₀₀) group fed lower amount of feed than (SS+G₃₀₀) group and (C) group , feed intake decreased with increasing stocking density due to reduction of access of birds to feed (**Feddes et al., 2002**) , these agreed with **Osman (1993) and Baeza et al., (2003)** who found reduction of feed intake at high stocking density in Muscovy and Pekin ducks.

However decreasing of feed intake by garlic supplementation especially in groups supplemented with (600 gm/ton feed) may be attributed to the effect of garlic supplementation in fulfill the nutrient requirement of birds by increase absorptive capacity of nutrients ,these results agreed with **Ibrahim et al., (2004)** who found that a decrease in feed consumption in all supplemented groups of muscovy ducks specially in groups received diet supplemented with 3% garlic or 1% onion + 3% garlic (5.2% - 4.6%) respectively but disagreed with **Kirkpinar et al., (2011)** who found that feed intake of broilers not affected by supplementation of garlic oil (300 mg/kg) and **Oladele et al., (2012)** who showed that feed intake was higher in garlic supplemented broilers (0.125% , 0.25% and 0.5%) than in control group .

Feed conversion ratio in Table (4) revealed that there was a significance ($p < 0.05$) difference in the FCR between groups. FCR increased with age to reach the highest value at the end of the experiment. It was noticed that groups supplemented with the higher concentration of garlic showed the best FCR as in (SS+G₆₀₀) group and (HS+G₆₀₀) group while the worst FCR was observed in HS group and HS-G₃₀₀ group respectively. it was obvious that ducks reared at standard stocking density convert their feed more efficiently than those kept at high stocking density which may be attributed to the loss of the energy which used for growth as the group at high density were more disturbed and had less opportunity to rest (**Hall (2001)** , these results in agreement with **Bilgili and Hess (1995)** who concluded that feed conversion was significantly improved when birds were given more space. however, but in variance with **Thaxton et al., (2006)** who reported a higher feed conversion ratio due to a higher stocking density used.

On the other hand supplementation of garlic (600 gm /ton feed) improve the FCR in (HS+G₆₀₀) group and this may be due to the enhancing effect of garlic on villi length, villi width and cryptal depth and goblet cell numbers in the duodenum, jejunum and ileum of birds which activates the entire absorptive process of nutrients and results in better feed efficiency (**Masoud, 2006; Incharoen et al., 2010; Oladele et al., 2012**), these results agreed with (**Ibrahim et al., 2004**) who found that FCR of muscovy ducks was significantly better by supplementation of 35 of garlic at 10 weeks than that at the end of the 12th week (3 versus 3.67) but disagreed with **Ismoyowati et al., (2015)** which found that FCR of ducklings were not affected by inclusion of 1 % of garlic .

Dressing yield and visceral organs weight and percentage

Table (5) showed that, dressing yield % was significance ($p < 0.05$) decreased in (HS+G₃₀₀) group and (HS) group compared with other groups and these may related to the lower final live weight of these groups. These agreed with **Ahaotu and Agbasu (2015)** who showed that increasing of stocking density above 4 birds /m² decrease dressing % of pekin ducks,. However this wasn't occurred in (HS+G₆₀₀) group and this may be due to the effect of garlic supplementation which increased it final body weight and dressing % . Of it These agreed with (**Demir et al., (2003)** who showed that using garlic powder in broiler's diet influenced carcass yield positively. While disagreed with **Onibi et al., (2009)** who stated that garlic supplementation had no significant effects on carcass characteristics

Visceral organ weight in table (5) showed no significance ($p < 0.05$) difference among groups in gizzard ,liver and heart weight , our results agreed with (**Sekerglu et al., 2011**) who showed no significant difference of different stocking density of broiler(9,13,17 birds/m² on organ weigh) and also with (**Onibi et al., 2009, Raeesi et al., 2010 and Fayed et al., 2011**) who showed that garlic supplementation had no significant effects on giblet weight among different treatments however disagreed with **Jayalakshmi et al., (2009)** who studied different stocking density 900 cm, 750 cm, 600 cm and 450 cm per bird and found higher organ weight at 450 cm/ bird .

Meat quality (fat %)

Meat quality (fat %) in table (6) showed that (HS+G₆₀₀) group and (SS+G₆₀₀) group showed the lower breast and thigh fat % compared with other groups and this indicate that garlic supplementation of (600 gm/ton) of ducks decrease the fat % in breast and thigh muscle than the other concentration, these may be due to the greater effect of garlic in reducing fatty acid

synthetase enzyme. These results in agreement with **Kim et al., (2009)** who found that dietary supplementation of different level of garlic 2% and 4% for 5 weeks for broiler lower fat content in chicken thigh muscle, **Choi et al., (2010)** who found that higher concentration of dietary garlic powder supplementation in broiler lower fat contents of thigh muscle significantly ($P < 0.05$) than the control group. However these results in disagreement with **Amouzmehr et al., (2012)** who showed that there were no significant differences of broiler breast and thigh fat in groups fed with different levels (0.3% and 0.6%) of garlic extracts .

CONCLUSION

Garlic supplementation of 600 g / ton feed in Muscovy ducks improve body weight and weight gain and showed better FCR and lower fat % in carcass in standard stocking density and even in high stocking density which may be useful for economical and efficient production of ducks

REFERENCES

- ABD EL AZIM, A, EL-KAMASH, E.M., AND NADIA M. EL-BAHY., HASSAN ,Z. S .2009.** Physiological effect of some feed additives under different feeding regimes of muscovy ducks . Egypt. Poult. Sci. Vol 29, IV: 1047-1060.
- AHAOTU, E.O. AND AGBASU, C.A (2015):** Evaluation of the stocking rate on growth performance, carcass traits and meat quality of male peking ducks. Scientific Journal of Biological Sciences .4 (3) 23-29.
- AOAC “ASSOCIATION OF OFFICIAL ANALYTICAL CHEMISTS. 1995.** Official Methods of Analysis. Association of Official Analytical Chemists.16th Ed., Washington, DC, USA.
- BAEZA, E., CHARTRIN, P. AND ARNOULD, C. 2003.** Effects of stocking density on welfare, growth performance and carcass quality in Muscovy ducks. Sci. Tech. Avi. 45:4–
- BAMPIDIS, V. A ., CHRISTODOULOU ,V., CHRISTAKI ,E., FLOROU-PANERI ,P. AND SPAIS , A. B. 2005.** Effect of dietary garlic bulb and garlic husk supplementation on performance and carcass characteristics of growing lambs. Anim. Feed Sci. Technol. 121: 273 – 283.
- BEDFORD, M. 2000.** Removal of antibiotic growth promoters from poultry diets: Implications and strategies to minimize subsequent problems. World’s Poultry Science Journal. 56: 347-65.
- BILGILI, S, F AND HESS J. B.1995.** placement density influences broiler carcass grade and meat yields. J.Appl .Poultry. Res, 4: 384-389.
- Brzóska F., Śliwiński ,B., Michalik-Rutkowska ,O. and Śliwa,J. 2015.** The effect of garlic (*Allium sativum* L.) on growth performance, mortality rate, meat and blood parameters in broilers. Ann. Anim. Sci., 15 (4): 961–975
- BUIJS, S., L., KEELING, S., RETTENBACHER, E., POUCKE,V AND TUYTTENS,F. A. M. 2009.** Stocking density effects on broiler welfare: Identifying sensitive ranges for different indicators. Poult. Sci. 88:1536–1543.
- CENGIZ ,O ., K’OKSAL ,B. H.,TATLI ,O ., SEVIM ,O ., AHSAN ,U ., UNER, A. G., ULUTA ,P. A., BEYAZ, D ., BUYUKYORUK,S ., YAKAN,A AND ONOL , A.G.2015.** Effect of dietary probiotic and high stocking density on the performance, carcass yield, gut microflora, and stress indicators of broilers. Poultry Science 94:2395–2403.
- DEMIR, E., SARICA, S., OZCAN, M.A., AND SUICMEZ, M. 2003.** The use of natural feed additives as alternatives for an antibiotic growth promoter in broiler diets. Br. J. Poultry Sci. 44:S44 S45.
- DERMAN, C. 2014.** Le point sur le marché du canard en. TeMA .31: 34-39.
- COATES, W., RALPH, A. AND COUNTY, S.2000.** Raising duck in small flocks. Review. Journal of Poultry Science 42.
- DOZIER, W. A., THAXTON, J. P., BRANTON, S. L., MORGAN, G. W., MILES, D. M., ROUSH, W. B., LOTT, B. D., AND VIZZIER-THAXTON, Y.(2005):** Stocking density effects on growth performance and processing yields of heavy broilers. Poult. Sci., 84: 1332–1338.
- DOZIER, W. A., THAXTON, J. P., PURSWELL, J. L., OLANREWAJU, H. A., BRANTON, S. L., AND ROUSH, W. B. 2006 .** Stocking density effects on male broilers grown to 1.8 kilograms of body weight. Poult. Sci., 85: 344–351.
- DWILOKA, B ., SETIADI , A ., SANTOSO , S.I ., SUPRIJATNA ,E., SUSANTI ,S. 2015.** Effects of duck feed supplemented with invasive giant salvinia (*Salvinia molesta*) on duck meat characteristics .Turk J Vet Anim Sci . 39: 668-675.
- EL-EDEL, M.A., ABO GHANEMA, I.I AND RASHED,R.R. 2015.** Impact of BCG Vaccination on Productive Performance and Immune Response To avian Influenza Vaccine in Two Breeds of Ducks. Global Veterinaria 14 (3): 425-431.
- EL-GHAMRY, A.A.; EL-MALLAH, G.M. AND EL YAMANI, A.T.2002.** The effect of incorporating yeast culture, *Nigella sativa* seed and fesh garlic in broiler diets on their performance. Egypt. Poult. Sci. 22 (II): 445-459.
- EL-KATCHA, M ., SOLTAN, M. A ., MOHAMED ,M ., SHARAF, M. M. AND HASEN,A.2016.** Growth Performance, Immune Response, Blood serum parameters, Nutrient Digestibility and Carcass Traits of Broiler Chicken as Affected by Dietary Supplementation of Garlic Extract (Allicin) .Alexandria Journal of Veterinary Sciences 2016 , pr. A 49 (2) : 50 – 64.
- ESTEVEZ, I. 2007.** Density allowances for broilers: Where to set the line. Poult. Sci. 86:1265–1272.
- ESTEVEZ, I., NEWBERRY, R.C., ARIAS DE REYNA, L. 1997.** Broiler chickens: a tolerant social system? Etologia .5, 19–29.
- FARHAD, K., GHORBANALI, S., AHMAD, K., AND ASAAD, V. 2011.** Evaluation of different medicinal plants blends in diets for broiler chickens. J. Med. Plants. Res., 5(10), 1971-1977.

- FAYED, R.H., ABEER.H ABD EL RAZIK. AND JEHAN, M. OUF .2011.** Effect of dietary garlic supplementation on performance, carcass traits and meat quality in broiler chicken. XVISAH Congress, Vienna, Austria.
- FEBRER, K., JONES, T.A., DONNELLY, C.A. & DAWKINS, M.S. 2006.** Forced to crowd or choosing to cluster? Spatial distribution indicates social attraction in broiler chickens. *Animal Behaviour*, 72: 1291–1300.
- FEDDES, J.J., EMMANUEL, E.J., AND ZUIDHOFT, M.J. 2002.** Broiler performance, body weight variance, feed and water intake, and carcass quality at different stocking densities. *Poult. Sci.*, 81: 774-779.
- GALAL, A., ALI, W.A.H., AHMED, A.M.H. AND ALI, K.A.A.2011.** Performance and carcass characteristics of Dumyat, Muscovy and pekin duck breeds. *Egyptian J. Anim. Prod.* 48(2):191-202.
- GHAZALAH, M., AND IBRAHIM, A. 1996.** The possibility of using some edible and aromatic oils in the nutrition of Muscovy ducks. *Egypt. Poult. Sci.* 16 (II): 305-328.
- GRASHORN, M.A. 2010.** Use of phytobiotics in broiler nutrition—an alternative to infeed antibiotics? *Journal of Animal and Feed Sciences*, 19: 338–347.
- IBM CORP (2013):** IBM SPSS statistics for windows, version 22.0, Armonk. NY.
- IBRAHIEM, I.A., ELAM, T.A., MOHAMED, F.F., AWADALLA , S.A .AND YOUSIF,Y.I.2004.** Effect of onion and /or garlic as feed additives on growth performance and immunity in broiler muscovy ducks. 1st Ann. Confr. , FVM., Moshtohor, Sept, 2004 Ibrahim A.I., et al., 2004
- INCHAROEN, T., YAMAUCHI, K., THONGWITTAYA, N. 2010.** Intestinal villus histological alterations in broilers fed dietary dried fermented ginger. *J Anim Physiol Anim Nutr (Burl)*. 94: 130-137.
- ISMOYOWATI. INDRASANTI, D ., MUFTI, M ., AND FARJAM,A.S. 2015.** Photobiotic Properties of Garlic, Red Ginger, Turmeric and Kencur in Growing Ducks . *Animal Production*. 17(1):49-55.
- JAYALAKSHMI, T., KUMARARAJ, R., SIVAKUMAR ,T. AND VANAN ,T T.(2009):** Carcass characteristics of commercial broilers reared under varying stocking density. *Tamilnadu J. Veterinary & Animal Sciences* 5 (4) 132- 135.
- KIRKPINAR , F , BORA ÜNLÜ , H , ÖZDEMİR, G .(2011):**Effects of oregano and garlic essential oils on performance, carcass, organ and blood characteristics and intestinal microflora of broilers *Livestock Science* .137 : 219–225
- LEWIS, M.R., ROSE, S.P., MACKENZIE, A.M AND TUCKER, L.A. 2003.** Effects of dietary inclusion of plant extracts on the growth performance of male broiler chickens, *Br, Poult, Sci*, 44, S43-S44.
- MASOUD, A.2006.** Effect of dietary garlic meal on histological structure of small intestine in broiler chickens. *J. Poult. Sci.*, 43(4): 378-383.
- NATIONAL RESEARCH COUNCIL. 1994.** Nutrient requirements of poultry. 9th rev. ed. Natl Acad Press, Washington, DC.
- OLADELE, O. A., EMIKPE, B. O. AND BAKARE, H. 2012.** Effects of Dietary Garlic (*Allium sativum* Linn.) Supplementation on body weight and gut morphometry of Commercial Broilers *Efect. Int. J. Morphol.*, 30(1):238-240.
- ONIBI, G.E., OLUWATOYIN, E., ADEBISI,A., FAJEMISIN,N., AYODE,V. AND ADETUN, J.I . 2009.** Response of broiler chickens in terms of performance and meat quality to garlic (*Allium sativum*) supplementation. *African Journal of Agricultural Research* 4 (5): 511 - 517.
- OSMAN, A. M. A. 1993.** Effect of stocking rate on growth performance, carcass traits and meat quality of male Peking ducks. *J. Agric. Rural. Dev. Trop.* 94:147–156.
- POURALI, M., MIRGHELENJ, S.A AND KERMANSHASHI, D.2010.** Effect of garlic powder on productive Performance and immune response of broiler chickens challenged with Newcastle disease virus, *Global Veterinaria* 4,616-621
- RAEESI, M., HOSEINI-ALIABAD, S.A ., ROOFCHAE, A., ZARE SHAHNEH,A.Z .AND PIRALI, S. 2010.** Effect of periodically use of Garlic (*Allium sativum*) powder on performance and carcass characteristics in broiler chickens, *World Academy Science*, 4(8):1213-1219.
- RAVINDRAN V., THOMAS D.V., THOMAS D.G., MOREL P.C.H. (2006):** Performance and welfare of broilers as affected by stocking density and zinc bacitracin supplementation. *Anim. Sci. J.*, 77: 110–116.
- SCAHAW (SCIENTIFIC COMMITTEE ON ANIMAL HEALTH AND ANIMAL WELFARE) 2000.** The Welfare of Chickens Kept for Meat Production (Broilers). European Commission, Health and Consumer Protection Directorate- General.
- SHENG-QIU, T., XIAO-YING, D., CHUN-MEI, J., JING-JING, P., SHAN-SHAN, L., JIN-DING, C. 2013.** Effect of *Bacillus subtilis* natto on Growth Performance in Muscovy Ducks. *Brazilian Journal of Poultry Science*. V.15, N.3: 169-286.
- SEKEROGLU, A., M. SARICA, M., GULAY, M.S .AND DUMAN ,M. 2011.** Effect of stocking density on chick performance, I nternal organ weights and blood parameters in broilers. *J. Anim. Vet. Adv.* 10: 246-250
- SIMITZIS, P.E ., KALOGERAKI ,E ., GOLIOMYTIS ,M , CHARISMIADOU ,M.A ., TRIANTAPHYLLOPOULOS ,K ., AYOUTANTI ,A ., NIFOROU , ,K ., HAGER-THEODORIDES ,A.L AND S.G. DELIGEORGIS, S.G.2012.** Impact of stocking density on broiler growth performance, meat characteristics, behavioural components and indicators of physiological and oxidative stress. *British Poultry Science* .V. 53, N. 6, pp. 721-730.
- TABOOSHA, M. F. 2014.** Effect of Stocking Density and Slaughter age on Growth Performance and Carcass Traits of Female Mule Ducks (Crossbred of Muscovy Drake and Pekin Ducks) During Summer Season in Egypt .*Middle East Journal of Applied Sciences*, 4(4): 1023-1033.

- THAXTON, J. P., DOZIER, W. A., BRANTON ,S. L., MORGAN ,G. W., MILES, D. M., ROUSH ,W. B., LOTT, B. D. AND VIZZIER-THAXTON, Y. 2005.** Evaluation of stocking density on physiological adaptive responses of broilers. Southern Poultry Science Society Meeting. Abstract 70. p. 17.
- TOLLBA, A.A. AND HASSAN, M.S. 2003.** Using some natural additives to improve physiological and productive performance of broiler chicks under high temperature conditions. Black cumin (*Nigella sativa*) or garlic (*Allium sativum*). *Poult. Sci.*, 23: 327-340.
- TONG , H. B ., LU ,J ., JZOU ,J.M .,WANG ,Q AND S. R. SHI, S.R. 2012.** Effects of stocking density on growth performance, carcass yield, and immune status of a local chicken breed . *Poultry Science* 91:667–673
- XIE, M., JIANG, Y., TANG, J., WEN, Z.G., HUANG, W AND HOU, S.S. 2014.** Effects of stocking density on growth performance, carcass traits, and foot pad lesions of White Pekin ducks. *Poult. Sci.* 93:1644–1648.
- YANG, C., CHOWDHURY, M., HOU, Y AND GONG, J. 2015.** Phytogetic compounds as alternatives to in-feed antibiotics: potentials and challenges in application. *Pathogens*, 4: 137-156.
- YEH, Y.Y., LIU, L. 2001.** Cholesterol lowering effect of garlic extracts and organosulfur compounds: Human and animal studies. *J Nutr* 131: 989-993.
- YOO, J. S., KIM, H. J., WANG, J. P., AO, X., JUNG, J. H. AND KIM, I. H. 2011.** Effects of fermented garlic powder supplementation on production performance, egg quality and blood characteristics of laying hens. *J. Anim. Sci.* Vol. 88.

How to cite this article:

Basma M. Bawish , Fayed, R.H., and Abeer H. Abdel Razek, 2018. Effect of Garlic as Feed Additive on Performance, Carcass Characteristics, and Meat Quality of Muscovy Ducks Reared in Different Stocking Densities. *Journal of Applied Veterinary Sciences*, 3 (1): 43 – 51.

DOI: <https://dx.doi.org/10.21608/javs.2018.67162>