

## **ESTIMATE OF SOME GENETIC PARAMETERS IN A COTTON CROSS THROUGH NORTH CAROLINA DESIGN III**

**Ali, Samia E. S.**

**Cotton Research Institute, Agricultural Research Center, Giza, Egypt.**

### **ABSTRACT**

North Carolina designs are very effective in breaking undesirable linkage and lead to creating genetic variability in a population. F<sub>1</sub> population was obtained by crossing the two cultivars Giza 88 P<sub>1</sub> and Australian P<sub>2</sub>. Followed crossing between F<sub>2</sub> with the original parents, P<sub>1</sub> and P<sub>2</sub>. The analysis of variance was highly significant for all studied traits. While, females sets were highly significant for all studied traits, which exhibit difference between them. Also, the males were significant for boll weight (BW) g /plant, seed cotton yield (SCY) and lint yield g/plant(LY) g/plant, which showed difference between them. The interactions between them were highly significant for boll weight (BW) and seed cotton yield (SCY). That showed variation background. The additive variance was higher than dominance for most studied traits. The contribution of male or female parents was more pronounced in the genetic variation. Additive components of variation were higher than dominance ones for most studied traits reflecting decreased ( $\sqrt{D/A}$ ) values (less than unity). While the dominance portion of the genetic variation played a role for some traits reflecting higher ( $\sqrt{D/A}$ ) values than unity for boll weight (BW) and fiber length (FL.2.5%).

**Keywords:** Cotton, North Carolina design III.epistasis.

### **INTRODUCTION**

The concept of biparental cross or biparental mating was originally developed by Comstock and Robinson, (1948, 1952). This technique provides information about additive and dominance components of genetic variance, which helps breeders in the choice of breeding procedure for genetic improvement of polygenic characters. Also, this technique based on the second order statistics (Singh and Narayanan, 2000). Also, it is very effective in breaking undesirable linkages as well as leading to certain genetic variability in a population by creating heterozygosity. Assessment of the components of genetic variance controlling yield and its components have been studied by several research workers ( Soliman, 2003 and Abd El-Salam, 2005). El-Mansy *et al.*, (2008) indicated that biparental mating was more effective in breaking undesirable linkage.

The objectives of the present study were 1. to evaluate seed cotton yield and its components and fiber properties of F<sub>2</sub> hybrids, 2. to obtain genetic information about the additive and dominance genetic variance, heritability in both broad and narrow senses and degree of dominance.

### **MATERIALS AND METHODS**

The materials used in this study were the selfed seeds of the Egyptian cotton variety Giza 88 and Australian one (Australly), which



**Table 2. Analysis of variance for yield, its components and fiber properties in the biparental cross.**

S.O.V	Mean Squares								
	d.f	B.W	S .C. Y	L. Y	L%	S. I	Mic	F.S	F.L2.5%
sets	3	0.119**	39407.47**	5763.60**	0.371	0.277**	0.022	72.071	0.106
R .sets	4	0.029	2064.23	691.47	0.195	0.064	0.032	62.180	0.358
F.sets	4	0.240**	684512.11**	146875.16**	13.450**	0.765**	2.315**	72.817	19.751**
M.sets	12	0.042*	6274.34**	1811.64**	0.800	0.058	0.051	57.246	0.423
F M									
.sets	12	0.051**	1978.48**	495.66	0.184	0.021	0.031	54.084	0.466
error	28	0.016	828.34	330.540	0.455	0.033	0.032	56.725	0.330

The interaction between females x males mean squares were highly significant for boll weight (B.W) and (S.C.Y), exhibit the difference between females and males, also the variation between males in their genetic background. On the other hand, insignificant mean squares due to this interaction revealed that the male or female had a degree of similarity for these traits.

**Table 3. Proportional contribution as percentage of males, females and their interactions for all studied traits.**

traits	Sources		
	Females%	Males%	Males x females interaction %
B.W	46.36	24.42	29.22
S. C. Y	96.51	2.65	0.84
L.Y	95.50	3.53	0.97
L%	82.00	14.64	3.36
S. I	76.27	17.38	6.34
Mic	90.46	5.95	3.58
F.S	17.90	42.22	39.88
F.L2.5%	88.10	5.67	6.23

Data in Table 3 showed that the male or female parents appeared to contribute the maximum portion in the genetic variation for most traits. The contribution of female parent was more pronounced in all traits except fiber strength. These results confirm the role of additive genetic effects in controlling these traits. Information about additive and dominance components of genetic variance, heritability and degree of dominance are showed in Table 4. Regarding the relative magnitude of additive (A) and dominance (D) components estimates of additive variance were higher than those of dominance variance for most traits resulting in ( $\sqrt{D/A}$ ) values less than unity and explaining that, the additive component was the predominant type in the inheritance of such traits. This was confirmed by high narrow sense heritability estimates. This agrees with Abd El Bary (2003), El-Mansy (2005) and El-Mansy *et al.* (2008). While, the magnitude of dominance component was larger than corresponding additive ones for B.W and F.L.2.5% reflecting higher ( $\sqrt{D/A}$ ) values than unity. These findings showed the importance of over dominance gene effects in the control of these traits and indicated that non-fixable genes could be exploited efficiently through hybrid breeding method for improving these traits. This agrees with Soliman

(2003) and El-Mansy *et al.* (2008), which was confirmed by low estimates of narrow sense heritability.

**Table 4. Component of genetic variance**

Genetic components	Traits							
	B.W	S.C.Y	L.Y	L%	S.I	Mic	F.S	F.L2.5%
A	0.026	1150.14	1481.095	0.345	0.025	0.019	0.521	0.093
D	0.034	828.3415	165.116	0.000	0.000	0.000	0.000	0.135
E	0.0163	828.3415	330.5402	0.455	0.0332	0.032	56.725	0.330
broad sense	78.75	70.49	83.28	43.17	42.84	36.92	0.91	40.87
narrow sense	33.96	40.98	74.93	43.17	42.84	36.92	0.91	16.67
√D/A	1.149	0.849	0.334	0.000	0.000	0.000	0.000	1.205

## REFERENCES

1. Abd El-Bary, A.M.R. 2003. Triallel analysis of some quantitative inherited traits in *G.barbadense*, Ph.D. Thesis, Fac. Agric. Mansoura Univ., Egypt.
2. Abd El-Salam M.E. 2005. Studies on breeding and maintenance of cotton varieties *G.barbadense*, L. Ph.D.Thesis, Fac. Agric., Kafr El-Sheikh. Tanta Univ. Egypt.
3. Comstock, R.E. and H.F. Robinson.1948. The components of genetic variance in populations of biparental progenies and their use in estimating average degree of dominance. *Biometrics*. 4:254-266.
4. Comstock, R.E. and H.F. Robinson .1952. Estimation average dominance of genes . In heterosis, Iowa State College Press, Ames, 494-516.
5. El-mansy, Y.M. 2005. Using genetic components for predicting new recombination in some cotton crosses *G.barbadense*, L. Ph.D. Thesis, Mansoura Univ., Egypt.
6. El-mansy, Y.M. M.A. Abou El-Yazied and M.E. Abd El - Salam 2008. Determining some genetic parameters in cotton (*Gossypium barbadense*, L.) through North Carolina Design III.
7. Soliman, Y.A.M 2003 Studies on some quantitative traits in some interspecific cotton crosses. Ph.D.Thesis Fac. Agric., Kafr El- Sheikh. Tanta Univ. Egypt.
8. Soliman, Y.A.M., M.A. Abou El- yazied and A.M.R Abd El-Bary . 2007. Application of F<sub>2</sub> biparental mating design to produce new cotton recombination. Proceeding Fifth plant breeding conference. Egypt. *J. Plant Breed*.11(1):11-23.

### تقدير بعض المكونات الوراثية فى القطن من خلال النموذج الثالث

ساميه البدر سيد على

معهد بحوث القطن – مركز البحوث الزراعية - الجيزة

أجرى هذا البحث بغرض تقدير مكونات التباين الوراثى و حساب الارتباط الوراثى المضيف و السيادة باستخدام طريقة التهجين الرجعى لعدد ١٦ من نباتات الجيل الثانى مع الابوين حيث تم الحصول على ٣٢ هجين بالتهجين الرجعى للجيل الثانى مع الابوين للهجين (جيزة ٨٨ × استرالى).

اظهر تحليل التباين لمعظم الصفات وجود قدر كبير من الاختلافات يسمح باجراء الانتخاب كما اظهرت أن المساهمة النسبية لكل من الأباء و الأمهات أكبر من المساهمة النسبية للتفاعل. كانت مساهمة الفعل الجينى المضيف أكبر من مساهمة الفعل الجينى السيادة لمعظم الصفات المدروسة مما انعكس على انخفاض قيم درجة السيادة عن الواحد الصحيح لمعظم الصفات بينما لعب الفعل الجينى السيادة دور اكبر فى دراسة بعض الصفات مثل وزن اللوزة و طول التيلة مما ادى الى زيادة متوسط درجة السيادة عن الواحد الصحيح. كان اتجاه السيادة غير معنوى لكل الصفات المدروسة مما يدل على أن السيادة غير موجة أى فى اتجاهين مؤكدا التوزيع غير المتمائل لجينات السيادة للأباء.

قام بتحكيم البحث

أ.د / محمود سليمان سلطان

أ.د / حسين يحي عوض

كلية الزراعة – جامعة المنصورة

مركز البحوث الزراعية