

POPULATION DENSITY OF SOME INSECT PESTS INFESTING GLOBE ARTICHOKE PLANTATIONS IN RELATION TO CERTAIN ECOLOGICAL FACTORS AT DAKAHLIA GOVERNORATE

Taha, M. A.¹; Horia A. Abd-el Wahab²; Hanaa I. Mahmoud¹ and Ghada EL S. Abd el hamed²

1. Science College, Al Azhar Univ., Cairo, Egypt

2. Plant Protection Research Institute, ARC, Dokki- Giza, Egypt

ABSTRACT

Field experiments were conducted at Dakahlia Governorate during two successive seasons 2007/2008 and 2008/2009 to study the population density of some insect pests infesting globe artichoke plantations namely *Capitophorus horni* Born, *Empoasca discipiens* Poali, *Autographa gamma* Linn. and *Spodoptera littoralis* (Biosd). Also the effect of certain weather factors (daily mean temperatures and daily mean R.H.) and plant age were studied on the population dynamics of the various insect pests. The weather factors and plant age had insignificant effect on the population density of *C. horni*, *E. discipiens*, *A. gamma* and *S. littoralis*. Also, the relative humidity had shown significant effect on population density of the four insect pests during the two seasons 2007/2008 and 2008/2009. The percentage of explained variance was 92.5 & 86.5 during the two successive seasons, respectively.

INTRODUCTION

Globe artichoke (*Cynara scolymus* L.) belongs to family Asteraceae (Compositae) is becoming one of the most important vegetable crops grown for both local consumption and export (Mansour, 1983 and Tawfik, 1994). Throughout the growing season, globe artichoke plants are liable to infestation by phytophagous pests such as *Capitophorus horni* Born, *Empoasca discipiens* Poali, *Spodoptera littoralis* Biosd and *Autographa gamma* Linn., which considered the most common and important insect pests of globe artichoke plants. In heavy infestation, these pests are causing serious damage to plants, leading to great reduction in the yield (Schrameyer, 2002). Aphids play an important role as a vector of plant viruses and produce honeydew (Kasperovich, 2002). Therefore, the purpose of this work was to study the effect of plant age and certain weather factors on the population density of some insect pests infesting globe artichoke.

MATERIALS AND METHODS

The experiments were carried out in the farm of Agriculture Research Center at Baramoon, Dakahlia Governorate during two successive seasons 2007/ 2008 and 2008/ 2009. An Area (about feddan) ,divided into nine replicates, was planted with globe artichoke (Balady variety), on 1st November and 2nd October during the 1st and 2nd seasons, respectively. Every two weeks randomly samples of ten leaves from ten plants from each replicate were collected and put in paper bags and taken to the laboratory,

where carefully examined by the aid of a stereoscopic-microscope for counting the insect pests. The whole area was free from any pesticides treatment. The artichoke aphid, *Capitophorus horni* Born. (nymphs and adults), *Empoasca discipiens* Poali (adults), *Autographa gamma* Linn. (immature stages and adults) and *Spodoptera littoralis* (Biosd) (immature stages and adults). were Counted and recorded at two weeks until the end of the season.

The records of meteorological data, the daily mean of minimum, maximum temperatures and daily mean relative humidity, were obtained from the meteorological records of Central Laboratory for Agriculture climate, Agriculture Research Center in Dakahlia Governorate.

Statistical analysis:

To investigate effects of plant age, weather factors on the population dynamics of the insects, simple correlation and partial regression were carried out using MSTAT program.

RESULTS AND DISCUSSION

Population fluctuations studies on, the *Capitophorus horni* Born., *Empoasca discipiens* Poali, *Autographa gamma* Linn. and *Spodoptera littoralis* (Biosd.) were investigated during the two seasons 2007/2008 and 2008/2009.

Season 2007/2008:

The data illustrated in Fig. (1) showed that the infestation of artichoke aphid, *C. horni* was started at 30 days after sowing on 9th December (334 individuals /10 leaves) then it was increased sharply to reach its maximum (1941 individuals /10 leaves) at 72 days after planting date on 6th January. After that the insect population fluctuated and decreased gradually to reach moderate level (505 individuals /10 leaves) on 16th March after 128 days from planting and fluctuated and increased to reach (1566 individuals/ 10 leaves) at 30th March. Finally the insect population decreased sharply to reach a lower level (5 individuals/ 10 leaves) at 184 days after planting on 11th May then the insect disappeared until the end of cultivation on 6th July 2008. Meanwhile, data illustrated that the population of *C. horni* was obviously higher on young plants than the older one. These results were in agreement with Barbagallo (1974).

Data represented in Fig. (1) indicated that the infestation of the artichoke leaves with *E. discipiens* appeared after 30 days after planting on 19th December 2007 (11 individuals/ 10 leaves) then disappeared absolutely until 11 May 2008 (24 individuals/ 10 leaves) after 184 days from planting. After that increased sharply after 212 days on 8th June and reached (146 individuals/ 10 leaves). Finally the insect population decreased gradually on 6th July after 240 days planting date (38 individuals/ 10 leaves). These results indicated that high number of insects infested the older plants of artichoke than the younger ones.

Data explained in Fig. (1) The infestation of the leaves of globe artichoke with *S. littoralis* appeared at 170 days after cultivation on 27th April (12 individuals/ 10 leaves). Then the population decreased until the end of

cultivation on 6th July 2008 after 240 days from planting date. These results also indicated that the older globe artichoke plants infested with *S. littoralis* than the younger plants. These results were in agreement with Rina and Paolo (1980).

Data showed in Fig. (1) illustrated that the infestation of globe artichoke leaves with *Autographa gamma* initiated after 156 days from planting on 13th April 2008 (10 individuals/ 10 leaves). Then the population increased to become (18 individuals/ 10 leaves) after 170 days on 27th April then the population deceased and then increased to reach (12 individuals/ 10 leaves) in the end of cultivation on 6th July 2008 after 240 days from planting date. This means that older globe artichoke plants were infested by *A. gamma* more than younger ones. These results were in agreement with Rina and Paolo (1980).




Fig (1): Population fluctuation of some insects infesting leaves of globe artichoke plant during 2007/ 2008 season in Dakahlia Governorate.

As shown in Table (1) the simple correlation “r” indicated insignificant correlation between the age of globe artichoke plants and the population of *C. horni* and *S. littoralis* on globe artichoke cultivar ($r = 0.487$ and 0.429), respectively, while there was significant correlation between the age of globe artichoke plants and the population of *E. discipiens*, and *A. gamma* ($r = 0.509$ and 0.548). The real effect of this factor which appears from the partial

regression values on *C. horni*, *E. discipiens*, *S. littoralis* and *A. gamma* population revealed significant effect (b. reg.= 4.614, 1.735, 2.913), respectively, during 2007/2008 season. Also, insignificant correlation was found between the insect populations and daily relative humidity ($r=0.206$) during the season 2007/2008 season. The partial regression analysis for the effect of weather factors on the populations, revealed that means of daily maximum and minimum temperatures had significant effect (b. reg.= 7.686 & 1.795), respectively, during the season of 2007/2008. Also, the means of daily relative humidity had significant effect (b. reg.= 1.185). These results are in agreement with El-Khouly *et al.* (1998).

The obtained results revealed that the combined effect of the tested plant age and weather factors was significant on the insects population where the calculated "F" value was 7.60.

The analysis of the variance revealed that the weather factors and the plant age are responsible for about 92.5% of the variability in the populations of the observed pests on the artichoke cultivar during 2007/2008 season.

Table (1): Effect of plant age and weather factors on the population fluctuations of some insect pests infesting globe artichoke cultivar during 2007/2008 season at Dakahlia Governorate.

Factors		Simple correlation and regression values			Partial regression values			Analysis of variance		E.V. %
		r	b	S.E.	b. reg.	S.E.	p	F	P	
Insect pests	<i>Capiophorus horni</i>	0.487	4.532	2.044	4.614	2.081	0.01	7.60	0.004	92.5
	<i>Empoasca discipiens</i>	0.509	2.637	3.079	1.735	2.025	0.03			
	<i>Spodoptera littoralis</i>	0.429	6.877	4.959	2.913	2.101	0.03			
	<i>Autographa gamma</i>	0.548	5.913	3.471	4.886	2.868	0.01			
Weather factors	Daily mean max. temp.	0.327	6.253	4.609	7.686	5.666	0.01			
	Daily mean min. temp.	0.481	2.173	7.366	1.795	6.084	0.03			
	Daily mean R.H.	0.206	3.196	2.352	1.185	3.798	0.01			

r: Simple correlation value.

b: Simple regression coefficient value.

b.reg.: Partial regression coefficient value.

E.V.: Explained variance.

Season 2008/2009:

The data illustrated in Fig. (2) showed that globe artichoke leaves infested with *C. horni* was very high and started after 47 days of planting on 18th November 2008 (101 individuals/ 10 leaves). Then the population increased to reach sharply (738 individuals/ 10 leaves) after 187 days of planting on 7th April 2009. After that the insect population decreased to reach (1 individual/ 10 leaves) at the end of cultivation after 257 days of planting on 16th June 2009. These results indicated that *C. horni* preferred the younger globe artichoke plants than the older ones. These results are in agreement with Barbagallo (1974) and Maisonneuve *et al.* (2003).

Data in Fig. (2) showed that the infestation of globe artichoke leaves with *E. discipiens* started on 24th March 2009 after 173 days of planting date (5 individuals/ 10 leaves). Then, the population increased rapidly to reach (15 individuals/10 leaves) after 187 days from cultivation on 7th April and finally

the population decreased. These results indicated that *E. discipiens* preferred the older globe artichoke plants than the younger ones.

The results showed in Fig. (2) that infestation of the globe artichoke leaves with *S. littoralis* which started in 21st April 2009 after 201 days and then disappeared but after that appeared again and reached (7 individuals/ 10 leaves) in the end of cultivation after 257 days of planting date on 16th June. The results showed that *S. littoralis* preferred the older globe artichoke plants than the younger ones. These results are in agreement with Rina and Paolo (1980).

Data in Fig. (2) indicated that globe artichoke leaves infested with *A. gamma* after 187 days of planting date on 7th April 2009 (3 individuals/ 10 leaves). Then, the population increased rapidly to reach (46 individuals/ 10 leaves) after 215 days of sowing on 5th May. After that the population fluctuated and decreased and finally the population increased again to reach (50 individuals/ 10 leaves) after 257 days on 16th June. These results revealed that *A. gamma* infest the older globe artichoke plants than the younger plants. These results are in agreement with Rina and Paolo (1980).




Fig. (2): Population fluctuation of some insects infesting leaves of globe artichoke plant during 2008/ 2009 season in Dakahlia Governorate.

As shown in Table (2) the simple correlation “r” indicated significant correlation between the age of globe artichoke plants and the population of *C. horni*, *S. littoralis* and *A. gamma* on globe artichoke cultivar ($r = 0.540$,

0.852 and 0.732). The real effect of this factor which appears from the partial regression values on *C. horni*, *E. discipiens*, *S. littoralis* and *A. gamma* population revealed significant effect (b. reg.= 6.591, 6.982, 2.731 and 2.301), respectively, during the season 2008/2009season.

In Table (2)The results indicated insignificant correlation between the populations of insects and mean daily maximum, minimum temperatures and daily mean of relative humidity ($r= 0.038, 0.198$ and 0.183), respectively, during 2008/2009 season. The partial regression analysis for the effect of weather factors on the populations revealed that means of daily maximum, minimum temperatures and daily relative humidity had significant effect (b. reg.= 1.819& 1.409), respectively, during the season of 2008/2009, as shown in table (2). Also, the means of daily relative humidity had significant effect (b. reg.= 9.867). These results are in agreement with El- Khouly et al. 1998 ; CAB, (2003) and Maisonneuve et al (2003).

The obtained results revealed that the combined effect of the tested plant age and weather factors was 86.5% on the insects population where the calculated "f" value was 3.82.

The analysis of the variance revealed that the weather factors and the plant age are responsible for about 86.5% of the variability in the populations of the observed pests on the artichoke cultivar during 2008/2009 season.

Table 2: Effect of plant age and weather factors on the population fluctuations of some insect pests infesting globe artichoke cultivar during 2008/2009 season at Dakahlia Governorate.

Factors		Simple correlation and regression values			Partial regression values			Analysis of variance		E.V.%
		r	b	S.E.	b. reg.	S.E.	p	F	P	
pests	<i>Capiyophorus horni</i>	0.044	2.649	2.891	6.591	7.191	0.01	3.82	0.03	86.5
	<i>Empoasca discipiens</i>	0.852	1.333	1.261	6.982	6.606	0.03			
	<i>Spodoptera littoralis</i>	0.258	1.093	1.089	2.731	2.721	0.01			
	<i>Autographa gamma</i>	0.332	1.019	1.437	2.301	3.243	0.03			
Weather factors	Daily mean max. temp.	0.038	1.792	9.982	1.819	1.013	0.01			
	Daily mean min. temp.	0.198	2.185	1.309	1.409	8.443	0.01			
	Daily mean R.H.	0.183	4.568	2.274	9.867	4.912	0.01			

r: Simple correlation value.

b: Simple regression coefficient value.

b.reg.: Partial regression coefficient value.

E.V.: Explained variance.

REFERENCES

- Barbagallo, S. (1974): Observations on the aphids (Homoptera, Aphidoidea) of artichoke (*Cynara scolymus* L.). Bollettino del Laboratorio di Entomologia Agraria' Filippo Silvestri' Portici. 31, 197- 252.
- CAB, Crop Protection Compendium (2003): global module. Commonwealth Agricultural Bureau International, Wallingford, UK.
- El-Khouly, A.S.; E.M.E. Khalafalla; M.M. Metwally, H.M. Helal and A.B. Mezaïen (1998). Seasonal abundance and population dynamics of certain sucking insects on Soya bean in Kafr-El-sheikh Governorate, Egypt. J. Agric. Res., 76(1): 141-151.

- Kasperovich, E. (2002): Application of bacitaurine in *Aphis gossypii* and *Tetranychus urticae* number limitation in the protected ground. Scientific Conference dedicated to the 90th anniversary of the birth of the corresponding member of the AS RB AL Samersov Minsk Prilukii, 177- 180.
- Maisonneuve, J. C.; N. Hugon and F. Lolivier (2003): Biological control with *Chrysoperla lucasina* against *Aphis fabae* on artichoke in Brittany (France). Bull. OILB/ SROP. 26:3, 149-153.
- Mansour, A.S. (1983): Effect of some growth regulators on some physiological characters and the yield of globe artichoke. M. Sc. Thesis, Fac. Agric. Cairo Univ. 88pp.
- Rina, I and P. Paolo (1980): Observations on catches of Lepidoptera in a cultivated field of Polignano (Bari). Entomologica. 16, 143- 182.
- Schrameyer, K. (2002): Pests of artichokes. Gemuse Munchen. 38:3, 16-18.
- Tawfik, A.M. (1994): Morphological and physiological studies on globe artichoke. Ph.D. Thesis, Fac. Agric. Cairo Univ. 150 pp.

الكثافة العددية لبعض الآفات الحشرية التي تصيب زراعات الخرشوف وعلاقتها ببعض العوامل البيئية في محافظة الدقهلية
مصطفى أمين طه^١، حورية على عبد الوهاب^٢، هناء إبراهيم محمود^١ و غادة السيد عبد الحميد^٢

١ - كلية العلوم- جامعة الأزهر- القاهرة

٢ - معهد بحوث وقاية النباتات- مركز البحوث الزراعية- الدقى- الجيزة

أجريت التجارب الحقلية في محافظة الدقهلية خلال سنتين متتاليتين ٢٠٠٧ / ٢٠٠٨ ، ٢٠٠٩ / ٢٠٠٨ لدراسة الكثافة العددية لبعض الآفات الحشرية التي تصيب زراعات الخرشوف وهى من الخرشوف *Capitophorus horni* Born ، نطاط أوراق البطاطس *Empoasca discipiens* Poali وورق القطن *Spodoptera littoralis* Biosd والفراشة ذات الحرف *Autographa gamma* Linn. Y وأيضاً دراسة تأثير بعض العوامل الجوية وهى درجة الحرارة الصغرى والكبرى والرطوبة النسبية وعمر النبات على التغير لتعداد الآفات الحشرية المذكورة.

وأظهرت النتائج تأثير بعض العوامل الجوية (الحرارة العظمى والصغرى والرطوبة النسبية) وعمر النبات على الكثافة العددية و وجدت علاقة ارتباط معنوية مختلفة بين درجة الحرارة العظمى والصغرى وتعداد الحشرات محل الدراسة ، بينما بالنسبة للرطوبة النسبية. توجد علاقة ارتباط معنوية مختلفة سالبة على نبات الخرشوف خلال عامي الدراسة وكانت نسبة الاختلاف بين العوامل الجوية وعمر النبات ٩٢,٥ % و ٨٦,٥ % خلال سنتين الدراسة.

قام بتحكيم البحث

كلية الزراعة – جامعة المنصورة
مركز البحوث الزراعية

أ.د / هاله احمد كامل الصيرفى
أ.د / حسن على طه