

## **RESPONSE OF POTATO (*Solanum tuberosum* L.) GROWTH, YIELD AND STORABILITY TO SOME ORGANIC MANURE RATES AND FOLIAR SPRAY OF SOME MACRONUTRIENTS: 1-VEGETATIVE GROWTH, YIELD AND YIELD COMPONENTS**

**EL-Sayed, Hala A.\* ; A. H. A. El-Morsy\*\* and H. M. B. EL-Metwally\*\***

\* Veg. and Flori. Dept., Fac. of Agric., Mans. Univ., Egypt.

\*\* Veg. Res. Dept., Hort. Res. Inst., Agric. Res. Center, Giza, Egypt.

### **ABSTRACT**

Two field experiments were conducted on potato (*Solanum tuberosum* L.) Spunta cv plants during the winter seasons of 2008/2009 and 2009/2010 at Kafre Meet Faris village near El-Mansoura, Dakahlia Governorate, Egypt, to study the effect of organic fertilizer rates (20, 30 and 40 m<sup>3</sup>/fed) and some inorganic fertilizers (NPK) levels (100%, 75% and 50% from recommended dose), either alone or in combination with foliar spray with some macronutrients sources (Hi-Fertil 1 and Ferty More) on the plant growth, yield, and yield components.

The results showed that the plants fertilized with 40 m<sup>3</sup>/fed. farmyard manure (FYM) followed by 30 m<sup>3</sup>/fed. significantly increased plant stem length, number of leaves/plant, leaf area/plant and foliage dry weight/plant as well as average of tubers weight/plant and total yield.

Increasing the supplied NPK level up to 100% from recommended level caused significant increases in the most vegetative growth parameters. Also, tubers weight/plant, marketable yield and total yield were increased significantly.

Also, the results indicated that the plants sprayed with Hi-Fertil 1(19-19-19, NPK), had a significant effect in most vegetative growth parameters while, plants sprayed with Ferty More (5-3-43, NPK) resulted in significant increases in foliage dry weight/plant, tubers weight/plant, marketable yield and total yield.

In general, the best results were obtained by using 40 m<sup>3</sup> FYM, 50% NPK level and foliar spray with Ferty More (5-3-43, NPK) followed by the interaction among 40 m<sup>3</sup> FYM, 50% NPK level and foliar spray with Hi-Fertil 1(19-19-19, NPK). These treatments achieved increases in tubers weight/plant total yield and marketable yield as well as decreasing unmarketable yield. Therefore, these treatments could be recommended for raising productivity, improving quality of potato and will also decrease the pollution of environment under the conditions of El-Dakahlia Governorate.

**Keywords:** Potato, *Solanum tuberosum*, organic farming, farmyard manure, FYM, mineral fertilizers, NPK, foliar application, foliar fertilization, vegetative growth, yield.

### **INTRODUCTION**

Potato (*Solanum tuberosum* L.) is a major world food crop. Potato is exceeded only by wheat, rice, and maize in world production for human consumption. In Egypt, it has been generally cultivated for both local consumption and export.

Therefore, increasing potato yield and improving tuber quality are essential aims for both growers and consumers, but it usually depends on many factors especially that influence the plant growth throughout the growth period. Nitrogen, phosphorus and potassium nutrition are three of major factors affecting growth, yield and quality of potato. However, there are some problems which prevent the farmers to use sufficient amounts of nitrogen phosphorus and potassium, such as nutritional requirements of potato plants are quite high, the continuous increases in the costs of using chemical fertilizers and environment pollution problems.

Therefore, it has become essential using safe substitute or supplements for chemical fertilizers, reduce the costs of produced yield and environmental pollution as well as increase production and improve quality of potato. One of the ways is to use organic manures as a source of essential nutrients, increased nutrient supply and improved the efficiency of macronutrients as well as its ability to meet some micronutrients requirements such as Fe, Zn, Mn and Cu which were reflected on plant uptake and plant growth, in addition to positive effect on the environment and public health (Kolbe *et al.*, 1995; EL-Nagar,1996). Several investigators reported that potato plants growth, yield and its quality as well as N, P, K, Fe, Zn, and Mn content in the plant tubers were affected by organic fertilization. In this respect, Abou-Hussein (1995), Abdel-Ati (1998), Arisha and Bardisi (1999); Abou-Hussein *et al.* (2002a and b), Awad *et al.* (2002) and Abd El-Kader (2002). Abou-Hussein *et al.* (2003) indicated that applying cattle manure combined with chicken manure increased tuber dry matter, total carbohydrates, specific gratify and potato tuber yield. In the same manner, Radwan and Tawfik (2004) reported that organic fertilization improved plant growth characters, yield and its quality and the content of Fe in potato tuber, El-Kassas *et al.* (2005) and El-Morsy *et al.* (2006) found that using chicken manure increased all vegetative growth characters, number of tubers/plant, average tuber weight, total tuber yield and chemical constituents in tubers.

Also, it is notable that, the other way is use the foliar fertilization. The foliar application is more economical than root fertilization due to the efficiency and lower cost. It is usually preferred because very small amounts of fertilizers are applied per feddan. It also reduces the number of passes of the applicant thereby reducing problem of soil compactness as well as less likely to result in ground water pollution. Foliar application of nitrogen (N), phosphorus (P) and potassium (K) maintains leaf nutrition in photosynthesis, enhances N, P and K content, carbon (C) balance and self-destructive mechanisms that have resulted in higher yields in crop plants (Boote *et al.*, 1978), Increasing the yield due to foliar nutrient in different crops has been well documented previously (Singh *et al.*, 1996 and Ali and Mishra, 2001).

Therefore, the main objective of the present study is to evaluate the effect of organic manure (FYM) rates, some inorganic NPK fertilizers levels, either alone or in combination with some sources of macronutrients foliar spray on the plant growth, yield and its components of potato plants under the conditions of El-Dakahlia Governorate.

## MATERIALS AND METHODS

In two field trials, potato (*Solanum tuberosum* L.) Spunta cv plants were grown in clay soil during the two winter seasons of 2008/2009 and 2009/2010 at Kafr Meet Faris village near El-Mansoura, Dakahlia Governorate, Egypt, to study the effect of farmyard manure (FYM) rates and some inorganic fertilizers (NPK) levels, either alone or in combination with some sources of macronutrients foliar spray on the plant growth, yield and its components of potato plants. Tuber seeds were planted on 17<sup>th</sup> and 19<sup>th</sup> of October in the first and the second seasons, respectively.

Randomized samples were obtained from the experiment soil before the application of organic and chemical fertilization in both seasons of this study to determine the physical and chemical contents according to the standard method described by Jackson (1973). The obtained results are presented in Table (1).

**Table (1): Physical and chemical analysis of the experimental soil.**

| Seasons   | Physical properties (%) |       |           |             | Chemical properties |         |             |               |               |                |
|-----------|-------------------------|-------|-----------|-------------|---------------------|---------|-------------|---------------|---------------|----------------|
|           | Clay                    | Silt  | Fine sand | Coarse sand | Texture             | O.M (%) | Total N (%) | Avail P (ppm) | Exch. K (ppm) | pH (1:2.5 w/v) |
| 2008/2009 | 49.71                   | 25.73 | 23.00     | 1.56        | clay                | 1.90    | 0.13        | 7.15          | 215.00        | 7.90           |
| 2009/2010 | 49.93                   | 25.88 | 22.47     | 1.72        | clay                | 2.10    | 0.14        | 7.95          | 229.00        | 8.02           |

The experimental design was split-split plots in a randomized block design with three replicates. Organic fertilizer rates occupied the main plots which were subdivided to 4 sub plots each contained one of the mineral nutrients (NPK) levels, while, the foliar spray treatments with macronutrients were occupied the sub-sub plot. The sub-sub plot area was 17.5 m<sup>2</sup> (1/400 fed.) which contained 5 rows, each 5 m long and 0.7m width. The experiment included 27 treatments which were the combination among 3 rates of farmyard manure fertilizer (FYM), 3 levels of mineral nutrients (NPK) and 3 treatments of foliar spray of macronutrients including control treatment as follows:

**Organic manure rates (main plots):**

- 1- 20 m<sup>3</sup>/fed. FYM (Control treatment).
- 2- 30 m<sup>3</sup>/fed. FYM.
- 3- 40 m<sup>3</sup>/fed. FYM.

Organic manure rates were distributed, spreaded and thoroughly mixed with the surface soil layer (0 - 20 cm) at preparing the soil before planting. The farmyard manure (FYM) used analysis was shown in Table (2) as follows:

**Table (2): Chemical analysis of farmyard manure (FYM).**

| Macroelements (%) |       |       | Microelements (ppm) |     |     |
|-------------------|-------|-------|---------------------|-----|-----|
| N                 | P     | K     | Fe                  | Zn  | Mn  |
| 1.580             | 0.553 | 1.625 | 346                 | 210 | 185 |

According to methods of (Jackson, 1973).

**Mineral Fertilizers (NPK) levels (sub plots):**

- 1- 100% NPK from recommended dose (180 Kg N + 75 Kg P<sub>2</sub>O<sub>5</sub> + 96 Kg K<sub>2</sub>O/fed.).
- 2- 75 % NPK from recommended dose (135 Kg N + 56.25 Kg P<sub>2</sub>O<sub>5</sub> + 72 Kg K<sub>2</sub>O/fed.).
- 3- 50 % NPK from recommended dose (90 Kg N + 37.5 Kg P<sub>2</sub>O<sub>5</sub> – 48 Kg K<sub>2</sub>O/Fed.).

Nitrogen was applied in the form of ordinary ammonium nitrate (33.5%N), phosphorus was applied in the form of calcium superphosphate (15.5% P<sub>2</sub>O<sub>5</sub>), and potassium was applied in the form of potassium sulphate (48% K<sub>2</sub>O).

Ammonium nitrate and superphosphate were divided and applied at two equal doses with the first and second irrigation and potassium sulfate was added at two times before second and third irrigation.

**Macronutrients foliar spray sources (sub-sub plots):**

- 1- Control (untreated with macronutrients)
- 2- Hi-Fertil 1 (19-19-19, NPK).
- 3- Ferty More (5-3-43, NPK).

Hi-Fertil 1 and Ferty More were supplied as a foliar application at 45, 55 and 65 days after planting in the rate of 3 g/L. The control treatment was sprayed with tap water.

Hi-Fertil 1 and Ferty More are a commercial fertilizers locally produced by Egyptian Fertilizer development center, AL DELTA for fertilizer and chemicals industry, Egypt.

The other cultural practices were applied according to the instructions laid down by the Ministry of Agriculture, Egypt.

**Data recorded and statistical analysis:**

**Vegetative growth characteristics:**

A random sample of three potato plants were taken from each plot at 90 days after planting (DAP) to estimate the plant stem length, number of leaves/plant, leaf area/plant (according to the method of Koller, 1972), foliage fresh weight/plant and foliage dry weight/plant.

**Yield and its components:**

At harvest time, 105 days after planting (DAP), all tubers of plants of each sub-sub plots were dig up, weighted in kg and converted to total yield (tons/fed), graded to three classes according to tuber diameter, <30 mm, 30-60 mm, and >60 mm, then each grade was weighted separately. Marketable yield including good shapes healthy tubers which from 30-60 mm and >60 mm, while, unmarketable yield of culls (offshape, blemished, green and diseased) and <30 mm in diameter. Also, average of tubers weight/plant was measured.

All data were subjected to the statistical analysis and means were compared using new L.S.D according to (Gomez and Gomez 1984).

## RESULTS AND DISCUSSION

### Plant growth characteristics:

Data presented in Table (3) show that stem length, number of leaves per plant, Leaf area and foliage fresh and dry weight were increased significantly with increasing FYM rates up to 40 m<sup>3</sup>/fed in both seasons. These increases in all plant growth parameters may be related to the higher contents of macro-elements (NPK) in FYM (Table 2) and this led to an increase of the metabolism activity and consequently increasing of plant growth. These results are in agreement with those of Karadogan (1996), Abou-Hussein *et al.* (2002 b) Singh and Kushwah (2006). El-Sirafy *et al.* (2008) reported that plant vegetative growth parameters expressed as plant height, number of leaves, foliage fresh and dry weight were increased with increasing FYM rate.

**Table (3): Vegetative growth characters as affected by FYM rates, NPK levels and macronutrients foliar spray sources during 2008/2009 (S1) and 2009/2010 (S2) winter seasons.**

| Characters                   | Stem length/plant (cm) |       | Number of leaves / plant |       | Leaf area (m <sup>2</sup> ) |      | Plant fresh weight (g) |        | Plant dry weight (g) |       |
|------------------------------|------------------------|-------|--------------------------|-------|-----------------------------|------|------------------------|--------|----------------------|-------|
|                              | S1                     | S2    | S1                       | S2    | S1                          | S2   | S1                     | S2     | S1                   | S2    |
| <b>FYM rates:</b>            |                        |       |                          |       |                             |      |                        |        |                      |       |
| 20 m <sup>3</sup>            | 55.63                  | 57.22 | 18.13                    | 19.44 | 0.33                        | 0.30 | 227.48                 | 228.07 | 20.81                | 20.33 |
| 30 m <sup>3</sup>            | 60.37                  | 60.22 | 21.44                    | 22.37 | 0.30                        | 0.37 | 203.22                 | 200.74 | 32.02                | 30.48 |
| 40 m <sup>3</sup>            | 63.89                  | 62.63 | 23.78                    | 24.41 | 0.38                        | 0.39 | 280.78                 | 282.07 | 37.07                | 37.22 |
| LSD 5%                       | 1.61                   | 1.80  | 1.37                     | 1.09  | 0.01                        | 0.01 | 3.89                   | 0.49   | 0.23                 | 0.28  |
| <b>NPK levels:</b>           |                        |       |                          |       |                             |      |                        |        |                      |       |
| 100%                         | 62.81                  | 64.41 | 23.26                    | 24.11 | 0.38                        | 0.38 | 276.89                 | 272.19 | 30.10                | 28.74 |
| 75%                          | 59.37                  | 59.80 | 20.70                    | 21.70 | 0.30                        | 0.36 | 249.00                 | 204.11 | 31.89                | 31.10 |
| 50%                          | 57.70                  | 50.81 | 19.89                    | 20.41 | 0.34                        | 0.34 | 230.09                 | 239.09 | 32.80                | 32.10 |
| LSD 5%                       | 1.79                   | 1.37  | 0.86                     | 1.06  | 0.004                       | 0.01 | 2.73                   | 2.80   | 0.22                 | 0.20  |
| <b>Foliar spray sources:</b> |                        |       |                          |       |                             |      |                        |        |                      |       |
| Control                      | 51.80                  | 52.02 | 18.41                    | 18.06 | 0.30                        | 0.36 | 211.96                 | 204.44 | 27.19                | 27.78 |
| Hi-Fertil 1                  | 67.78                  | 68.67 | 24.48                    | 20.26 | 0.41                        | 0.42 | 296.93                 | 309.78 | 31.80                | 31.00 |
| Ferty More                   | 60.26                  | 58.89 | 20.96                    | 22.41 | 0.30                        | 0.32 | 202.09                 | 201.67 | 36.80                | 34.26 |
| LSD at 5%                    | 1.24                   | 1.07  | 0.60                     | 0.62  | 0.01                        | 0.01 | 2.39                   | 2.63   | 0.30                 | 0.23  |

As regard to the effect of NPK levels on vegetative growth, data in Table (3) indicated that increasing NPK level up to level one 100% NPK (180+75+96 unit/fed) recorded the highest values in plant height, number of leaves/plant, leaf area/plant and foliage fresh weight/plant in both seasons. While, the foliage dry weight/plant was reduced at plants received 100% NPK. These results may be related to the vital role of nitrogen, phosphorus and potassium nutrition as the major factors affecting growth such as nitrogen is a main constituent of many organic compounds in plants, phosphorus plays an important role in accumulation and release of energy during cellular metabolism and potassium plays a vital role for a normal cell division of plants (Gardener *et al.*, 1985; Marschner, 1995 and Black, 1973). Several

investigators reported that, plant height, number of main stems, fresh and dry weight of plant foliage were significantly increased with increasing NPK level (Jasiwal, 1995; Rabie, 1996; Arisha and Bardisi, 1999; Sarhan *et al.*, 2004; Al-Moshileh and Moftah, 2005 and Abd El-Aal *et al.*, 2008).

Concerning the effect of foliar spray (NPK) on vegetative growth, data presented in Table (3) indicated that plant height, number of leaves/plant, leaf area/plant and foliage fresh and dry weight/plant significantly increased at the plants sprayed with Hi-Fertil 1 (19-19-19 NPK) in both seasons of study. These results may be related to the fact that these elements which can be readily absorbed by the leaves as a result of foliar spraying application and not lost through fixation, decomposition or leaching under unfavorable soils conditions (Doeing, 1986). Similar results were found by Joudu *et al.* (2001), Ayyub *et al.* (2006) and Al-Betar and Abdou (2010).

With respect to the effect of interaction between FYM rates and NPK levels, data presented in Table (4) showed that potato plants fertilized with 40 m<sup>3</sup> FYM rate and 100% NPK level resulted in the highest significant number of leaves/plant in both seasons, but, these increases in leaf area and plant fresh weight were significant in the first season only. On the other hand, the highest foliage dry weight/plant was obtained from plants received 50% NPK followed by 75% NPK under 40 m<sup>3</sup> FYM rate in both seasons. It is notable that, there were no significant differences between plant dry weight values when plants received 50% NPK or 75% NPK levels under application of 30 m<sup>3</sup> or 40 m<sup>3</sup> FYM rates in both seasons. In this respect, Abd El-Kader (2002) reported that potato plants fertilized with 30 m<sup>3</sup>/fed farmyard manure and NPK level (90–85–170 kg/fed) resulted in the highest significant plant height, number of leaves/plant, foliage fresh weight/plant foliage dry weight/plant and leaf area/plant. Same results reported by Borin *et al.* (1987) Dorobantu *et al.* (1989) and El-Banna and Abd El-Salam (2000).

Also, data presented in Table (4) show that the interaction between FYM rates and foliar spray (NPK) had a positive significant effects on plant fresh weight and foliage dry weight/plant in both seasons. While, plant stem length and leaf area were affect significantly in the first season only and number of leaves/plant was affected in second season only. Data indicated also, plants treated by Hi-Fertil 1 as a macronutrient foliar spray source and fertilized with 40 m<sup>3</sup> FYM/fed gave the highest values of vegetative growth parameters followed by the same foliar spray source with 30 m<sup>3</sup> FYM/fed. in both seasons.

Regarding, the effect of interaction between NPK levels and foliar spray NPK on vegetative growth parameters, data in Table (4) indicated that this interaction had a positive significant effects on number of leaves/plant, plant fresh weight and plant dry weight in both seasons. While, the plant stem length was affected in the first season only. On the other hand, results also indicated that the highest record in plant dry weight was obtained from plants sprayed with Ferty More (5-3-43, NPK) and fertilized with 50% or 75% NPK levels in both seasons.

**Table (4): Vegetative growth characters as affected by the interactions of FYM rates x NPK levels, FYM x macronutrients foliar spray sources and NPK levels x Foliar sources during 2008/2009 (S1) and 2009/2010 (S2) winter seasons.**

| Characters        | Stem length/plant (cm)      |            | Number of leaves / plant |             | Leaf area (M <sup>2</sup> ) |            | Plant fresh weight (g) |              | Plant dry weight (g) |             |       |
|-------------------|-----------------------------|------------|--------------------------|-------------|-----------------------------|------------|------------------------|--------------|----------------------|-------------|-------|
|                   | S1                          | S2         | S1                       | S2          | S1                          | S2         | S1                     | S2           | S1                   | S2          |       |
| <b>Treatments</b> |                             |            |                          |             |                             |            |                        |              |                      |             |       |
| <b>FYM rates</b>  | <b>NPK levels</b>           |            |                          |             |                             |            |                        |              |                      |             |       |
| 2. m <sup>3</sup> | 100%                        | 57.56      | 62.56                    | 19.33       | 20.33                       | 0.35       | 0.36                   | 248.78       | 242.89               | 27.11       | 26.56 |
|                   | 75%                         | 55.33      | 57.00                    | 18.56       | 19.33                       | 0.32       | 0.35                   | 224.22       | 227.67               | 25.78       | 25.11 |
|                   | 50%                         | 54.00      | 52.11                    | 18.00       | 18.67                       | 0.32       | 0.33                   | 209.44       | 213.67               | 24.56       | 24.33 |
| 3. m <sup>3</sup> | 100%                        | 63.67      | 64.44                    | 24.11       | 24.67                       | 0.38       | 0.39                   | 271.11       | 273.67               | 30.33       | 27.89 |
|                   | 75%                         | 59.44      | 60.00                    | 20.44       | 21.44                       | 0.35       | 0.36                   | 251.89       | 254.56               | 32.67       | 30.89 |
|                   | 50%                         | 58.00      | 56.22                    | 19.78       | 21.00                       | 0.34       | 0.35                   | 236.67       | 239.00               | 34.56       | 32.67 |
| 4. m <sup>3</sup> | 100%                        | 67.22      | 66.22                    | 26.33       | 27.33                       | 0.41       | 0.41                   | 310.78       | 300.00               | 33.00       | 31.78 |
|                   | 75%                         | 63.33      | 62.56                    | 23.11       | 24.33                       | 0.38       | 0.38                   | 270.89       | 280.11               | 37.22       | 37.44 |
|                   | 50%                         | 61.11      | 59.11                    | 21.89       | 21.56                       | 0.35       | 0.36                   | 260.67       | 266.11               | 39.44       | 39.44 |
| <b>LSD 5%</b>     | <b>N.S</b>                  | <b>N.S</b> | <b>2.78</b>              | <b>3.18</b> | <b>0.02</b>                 | <b>N.S</b> | <b>16.01</b>           | <b>N.S</b>   | <b>6.78</b>          | <b>7.51</b> |       |
| <b>FYM</b>        | <b>Foliar spray sources</b> |            |                          |             |                             |            |                        |              |                      |             |       |
| 2. m <sup>3</sup> | Control                     | 49.44      | 50.11                    | 16.22       | 17.56                       | 0.27       | 0.29                   | 181.67       | 179.78               | 21.67       | 23.11 |
|                   | Hi-Fertil 1                 | 62.44      | 65.33                    | 21.67       | 21.78                       | 0.40       | 0.40                   | 269.00       | 285.00               | 24.44       | 24.33 |
|                   | Ferty More                  | 55.00      | 56.22                    | 18.00       | 19.00                       | 0.33       | 0.34                   | 231.78       | 219.44               | 31.33       | 28.56 |
| 3. m <sup>3</sup> | Control                     | 53.89      | 51.78                    | 18.22       | 18.44                       | 0.31       | 0.32                   | 214.67       | 207.89               | 27.00       | 26.56 |
|                   | Hi-Fertil 1                 | 67.89      | 69.78                    | 24.56       | 25.67                       | 0.41       | 0.42                   | 291.11       | 302.33               | 32.89       | 31.33 |
|                   | Ferty More                  | 59.33      | 59.11                    | 21.56       | 23.00                       | 0.36       | 0.36                   | 253.89       | 257.00               | 37.67       | 33.56 |
| 4. m <sup>3</sup> | Control                     | 52.22      | 55.67                    | 20.78       | 19.67                       | 0.33       | 0.34                   | 239.56       | 225.67               | 29.89       | 30.67 |
|                   | Hi-Fertil 1                 | 73.00      | 70.89                    | 27.22       | 28.33                       | 0.43       | 0.43                   | 330.67       | 342.00               | 38.22       | 37.33 |
|                   | Ferty More                  | 66.44      | 61.33                    | 23.33       | 25.22                       | 0.37       | 0.38                   | 272.11       | 278.56               | 41.56       | 40.67 |
| <b>LSD 5%</b>     | <b>7.79</b>                 | <b>N.S</b> | <b>N.S</b>               | <b>3.66</b> | <b>0.03</b>                 | <b>N.S</b> | <b>19.88</b>           | <b>19.93</b> | <b>4.05</b>          | <b>4.38</b> |       |
| <b>NPK</b>        | <b>Foliar spray sources</b> |            |                          |             |                             |            |                        |              |                      |             |       |
| 100%              | Control                     | 53.56      | 56.33                    | 19.33       | 19.78                       | 0.33       | 0.34                   | 226.11       | 218.33               | 25.00       | 26.00 |
|                   | Hi-Fertil 1                 | 71.22      | 73.56                    | 27.11       | 28.00                       | 0.43       | 0.43                   | 335.44       | 338.78               | 29.56       | 28.33 |
|                   | Ferty More                  | 63.69      | 63.33                    | 23.33       | 24.56                       | 0.38       | 0.39                   | 269.11       | 259.44               | 35.89       | 31.89 |
| 75%               | Control                     | 51.33      | 52.44                    | 18.22       | 17.89                       | 0.30       | 0.32                   | 210.11       | 205.22               | 26.22       | 26.89 |
|                   | Hi-Fertil 1                 | 68.22      | 68.44                    | 23.78       | 25.00                       | 0.41       | 0.42                   | 285.89       | 304.89               | 32.44       | 31.67 |
|                   | Ferty More                  | 58.56      | 58.67                    | 20.11       | 22.22                       | 0.35       | 0.36                   | 251.00       | 252.22               | 37.00       | 34.89 |
| 50%               | Control                     | 50.67      | 48.78                    | 17.67       | 18.00                       | 0.28       | 0.29                   | 199.67       | 189.78               | 27.33       | 27.44 |
|                   | Hi-Fertil 1                 | 63.89      | 64.00                    | 22.56       | 22.78                       | 0.39       | 0.40                   | 269.44       | 285.67               | 33.56       | 33.00 |
|                   | Ferty More                  | 58.56      | 54.67                    | 19.44       | 20.44                       | 0.34       | 0.35                   | 237.67       | 243.33               | 37.67       | 36.00 |
| <b>LSD 5%</b>     | <b>4.16</b>                 | <b>N.S</b> | <b>2.34</b>              | <b>2.60</b> | <b>N.S</b>                  | <b>N.S</b> | <b>33.66</b>           | <b>28.15</b> | <b>1.81</b>          | <b>2.59</b> |       |

Concerning the interaction effect of the studied factors (organic manure (FYM), NPK levels and NPK foliar fertilization) on vegetative growth, data in Table (5) reveal that the highest values in the most vegetative growth parameters were obtained from plants fertilized by 40 m<sup>3</sup>/fed FYM, 100% NPK level and sprayed with Hi-Fertil 1 (19-19-19 NPK). On the other hand, the interaction among 30 m<sup>3</sup>/fed FYM, 100% NPK level and Ferty More (5-3-43 NPK) gave the highest values in plant dry weight as compared with other interactions in both seasons.

Table (5): Vegetative growth characters as affected by the interaction among FYM rates, NPK levels and macronutrients foliar spray sources during 2008/2009 (S1) and 2009/2010 (S2) winter seasons.

| Characters                   |      |             | Stem length/plant (cm) |       | Number of leaves / plant |       | Leaf area (M <sup>2</sup> ) |      | Plant fresh weight (g) |        | Plant dry weight (g) |       |
|------------------------------|------|-------------|------------------------|-------|--------------------------|-------|-----------------------------|------|------------------------|--------|----------------------|-------|
|                              |      |             | S1                     | S2    | S1                       | S2    | S1                          | S2   | S1                     | S2     | S1                   | S2    |
| Treatments                   |      |             |                        |       |                          |       |                             |      |                        |        |                      |       |
| FYM NPK Foliar spray sources |      |             |                        |       |                          |       |                             |      |                        |        |                      |       |
| 20 m <sup>3</sup>            | 100% | Control     | 63.67                  | 68.67 | 22.33                    | 22.67 | 0.29                        | 0.30 | 295.67                 | 303.00 | 26.00                | 25.67 |
|                              |      | Hi-Fertil 1 | 58.67                  | 61.67 | 19.33                    | 19.67 | 0.36                        | 0.37 | 246.33                 | 232.00 | 33.00                | 30.00 |
|                              |      | Ferty More  | 50.00                  | 50.00 | 17.00                    | 17.67 | 0.27                        | 0.31 | 175.67                 | 179.00 | 21.33                | 23.00 |
|                              | 75%  | Control     | 62.67                  | 66.33 | 21.33                    | 21.67 | 0.40                        | 0.40 | 268.67                 | 285.33 | 24.67                | 24.00 |
|                              |      | Hi-Fertil 1 | 53.33                  | 54.67 | 17.33                    | 18.67 | 0.32                        | 0.33 | 228.33                 | 218.67 | 31.33                | 28.33 |
|                              |      | Ferty More  | 48.00                  | 43.00 | 15.33                    | 16.33 | 0.25                        | 0.26 | 165.00                 | 166.67 | 21.33                | 22.33 |
|                              | 50%  | Control     | 61.00                  | 61.00 | 21.33                    | 21.00 | 0.39                        | 0.39 | 242.67                 | 266.67 | 22.67                | 23.33 |
|                              |      | Hi-Fertil 1 | 53.00                  | 52.33 | 17.33                    | 18.67 | 0.31                        | 0.33 | 220.67                 | 207.67 | 29.67                | 27.33 |
|                              |      | Ferty More  | 50.00                  | 50.00 | 17.00                    | 17.67 | 0.27                        | 0.31 | 175.67                 | 179.00 | 21.33                | 23.00 |
| 30 m <sup>3</sup>            | 100% | Control     | 71.33                  | 75.00 | 28.00                    | 29.00 | 0.43                        | 0.43 | 320.00                 | 331.33 | 30.33                | 27.00 |
|                              |      | Hi-Fertil 1 | 62.00                  | 64.00 | 24.67                    | 25.67 | 0.38                        | 0.39 | 270.67                 | 264.67 | 35.33                | 30.67 |
|                              |      | Ferty More  | 51.67                  | 50.33 | 17.33                    | 17.33 | 0.30                        | 0.31 | 220.67                 | 208.00 | 27.33                | 26.67 |
|                              | 75%  | Control     | 70.00                  | 69.33 | 23.67                    | 24.33 | 0.40                        | 0.41 | 282.33                 | 296.00 | 32.67                | 32.00 |
|                              |      | Hi-Fertil 1 | 56.67                  | 60.33 | 20.33                    | 22.67 | 0.35                        | 0.36 | 252.67                 | 259.67 | 38.00                | 34.00 |
|                              |      | Ferty More  | 52.33                  | 50.67 | 17.67                    | 18.67 | 0.29                        | 0.30 | 200.67                 | 190.67 | 28.33                | 28.00 |
|                              | 50%  | Control     | 62.33                  | 65.00 | 22.00                    | 23.67 | 0.40                        | 0.40 | 271.00                 | 279.67 | 35.67                | 34.00 |
|                              |      | Hi-Fertil 1 | 59.33                  | 63.00 | 19.667                   | 20.67 | 0.34                        | 0.35 | 238.33                 | 246.67 | 39.67                | 36.00 |
|                              |      | Ferty More  | 50.00                  | 50.00 | 17.00                    | 17.67 | 0.27                        | 0.31 | 175.67                 | 179.00 | 21.33                | 23.00 |
| 40 m <sup>3</sup>            | 100% | Control     | 78.67                  | 77.00 | 31.00                    | 32.33 | 0.46                        | 0.45 | 390.67                 | 382.00 | 32.33                | 31.33 |
|                              |      | Hi-Fertil 1 | 70.33                  | 64.33 | 26.00                    | 28.33 | 0.40                        | 0.40 | 290.33                 | 281.67 | 39.33                | 35.00 |
|                              |      | Ferty More  | 52.33                  | 57.00 | 20.33                    | 18.67 | 0.32                        | 0.33 | 234.00                 | 228.67 | 30.00                | 31.00 |
|                              | 75%  | Control     | 72.00                  | 69.67 | 26.33                    | 29.00 | 0.43                        | 0.44 | 306.67                 | 333.33 | 40.00                | 39.00 |
|                              |      | Hi-Fertil 1 | 65.67                  | 61.00 | 22.67                    | 25.33 | 0.37                        | 0.38 | 272.00                 | 278.33 | 41.67                | 42.33 |
|                              |      | Ferty More  | 51.67                  | 52.67 | 20.00                    | 19.00 | 0.30                        | 0.31 | 233.33                 | 212.00 | 32.33                | 32.00 |
|                              | 50%  | Control     | 68.33                  | 66.00 | 24.33                    | 23.67 | 0.40                        | 0.41 | 294.67                 | 310.67 | 42.33                | 41.67 |
|                              |      | Hi-Fertil 1 | 63.33                  | 58.67 | 21.33                    | 22.00 | 0.36                        | 0.37 | 254.00                 | 275.67 | 43.67                | 44.67 |
|                              |      | Ferty More  | 50.00                  | 50.00 | 17.00                    | 17.67 | 0.27                        | 0.31 | 175.67                 | 179.00 | 21.33                | 23.00 |
| LSD at 5 %                   |      |             | 6.16                   | 7.29  | N.S                      | 2.32  | N.S                         | N.S  | 32.73                  | 22.10  | 3.41                 | 3.43  |

**Yield and its components:**

Data presented in Table (6) indicated that tuber weight/plant, marketable yield and total yield were increased significantly by application of FYM at 40 m<sup>3</sup>/fed. in both seasons. These increases may be related to the higher contents of macro-elements (NPK) in FYM (Table 2) and this led to an increase of the metabolism activity, its led to increasing of plant growth (Table 3) consequently, increasing total yield. Suresh and Ramanathan (2001), Awad *et al.* (2002), Singh and Gupta (2005), Colla *et al.* (2005), El Morsy *et al.* (2006) and El-Dissoky (2008) reported that number of tuber plant, average tuber weight, marketable and total yield were significantly increased by application of farmyard manure.

Concerning the effect of NPK levels on yield and its components, data in Table (6) show that increasing NPK level up to 100% NPK level, recorded the highest values in total yield and its components in both seasons. Similar



results were obtained by Maier *et al.* (1994), Minhas and Sood (1994), Arisha and Bardisi (1999), Kushwah and Banafar (2003) and Abd El-Aal *et al.* (2008) indicated that the potato yield was increased with increasing NPK rate.

Regarding, the effect of macronutrients foliar spray sources on total yield and its components, data in Table (6) indicate that plants sprayed with Ferty More (5-3-43, NPK) followed by Hi-Fertil 1(19-19-19 NPK) produced the highest tubers yield/plant, marketable and total yield/fed. in both seasons of study. These results may be related to increasing the activity of plant metabolism, which reflected on tubers yield and enhance tuber quality. Similarly, Boliglowa and Dzienia (1999), Singh *et al.* (1996) and Ali and Mishra (2001) found that marketable tubers yield and total yield were significantly increased by foliar application of macronutrients.

**Table (6): Total yield and its components of potato plants as affected by FYM rates, NPK levels and macronutrients foliar spray sources during 2008/2009 (S1) and 2009/2010 (S2) winter seasons.**

| Characters<br>Treatments     | Tubers weight/plant (g) |        | Marketable yield (ton/fed.) |       | Unmarketable yield (ton/fed.) |       | Total yield (ton/fed.) |       |
|------------------------------|-------------------------|--------|-----------------------------|-------|-------------------------------|-------|------------------------|-------|
|                              | S1                      | S2     | S1                          | S2    | S1                            | S2    | S1                     | S2    |
| <b>FYM rates:</b>            |                         |        |                             |       |                               |       |                        |       |
| 20 m <sup>3</sup>            | 428.63                  | 428.63 | 12.99                       | 12.99 | 0.902                         | 0.999 | 13.89                  | 14.19 |
| 30 m <sup>3</sup>            | 486.07                  | 492.33 | 12.99                       | 14.22 | 0.814                         | 0.771 | 14.81                  | 14.88 |
| 40 m <sup>3</sup>            | 532.78                  | 538.78 | 14.77                       | 14.08 | 1.278                         | 1.349 | 16.03                  | 15.93 |
| LSD at 5%                    | 3.19                    | 1.14   | 0.04                        | 0.00  | 0.320                         | 0.014 | 0.04                   | 0.00  |
| <b>NPK levels:</b>           |                         |        |                             |       |                               |       |                        |       |
| 100%                         | 489.34                  | 490.73 | 14.12                       | 14.08 | 1.033                         | 1.001 | 15.14                  | 15.13 |
| 75%                          | 480.48                  | 488.80 | 12.90                       | 12.99 | 0.980                         | 0.997 | 14.93                  | 14.99 |
| 50%                          | 478.77                  | 480.41 | 12.79                       | 12.90 | 0.977                         | 0.971 | 14.77                  | 14.87 |
| LSD at 5%                    | 1.71                    | 1.20   | 0.04                        | 0.02  | 0.414                         | 0.209 | 0.04                   | 0.02  |
| <b>Foliar spray sources:</b> |                         |        |                             |       |                               |       |                        |       |
| Control                      | 421.96                  | 418.77 | 12.02                       | 12.12 | 1.104                         | 1.173 | 14.13                  | 14.29 |
| Hi-Fertil 1                  | 492.02                  | 499.37 | 14.00                       | 14.14 | 1.030                         | 1.047 | 15.08                  | 15.19 |
| Ferty More                   | 534.00                  | 541.80 | 14.78                       | 14.71 | 0.847                         | 0.800 | 15.62                  | 15.62 |
| LSD at 5%                    | 1.40                    | 1.40   | 0.04                        | 0.02  | 0.690                         | 0.409 | 0.04                   | 0.02  |

With respect to the effect of interaction between FYM rates and NPK levels, data in Table (7) showed that potato plants fertilized with 40 m<sup>3</sup> FYM rate and 50% or 75% NPK levels resulted in the highest significant tuber weight/plant and total yield. The values in both treatments were not significant differences with each other. These results were true during both seasons. Similar results were obtained by Singh *et al.* (1996) and Sadej *et al.* (2004).

Also, data presented in Table (7) showed that the interaction between FYM rates and macronutrients foliar spray sources had a positive significant effects on total yield and its components in both seasons. Data indicated that, plants sprayed with Ferty More (5-3-43, NPK) followed by Hi-Fertil 1 (19-19-19, NPK) and fertilized with 40 m<sup>3</sup> FYM/fed. gave the highest tubers weight/plant and total yield.

Table (7): Total yield and its components of potato plants as affected by the interactions of FYM rates x NPK levels, FYM x macronutrients foliar spray sources and NPK levels x Foliar sources during 2008/2009 (S1) and 2009/2010 (S2) winter seasons.

| Characters        | Tubers weight/plant (g)     |        | Marketable yield (ton/fed.) |       | Unmarketable yield (ton/fed.) |       | Total yield (ton/fed.) |       |       |
|-------------------|-----------------------------|--------|-----------------------------|-------|-------------------------------|-------|------------------------|-------|-------|
|                   | S1                          | S2     | S1                          | S2    | S1                            | S2    | S1                     | S2    |       |
| <b>Treatments</b> |                             |        |                             |       |                               |       |                        |       |       |
| <b>FYM rates</b>  | <b>NPK levels</b>           |        |                             |       |                               |       |                        |       |       |
| 2. m <sup>3</sup> | 100%                        | 472.02 | 472.89                      | 13.71 | 13.03                         | 0.903 | 1.100                  | 14.77 | 14.74 |
|                   | 75%                         | 410.89 | 424.89                      | 12.90 | 13.12                         | 0.904 | 0.978                  | 13.80 | 14.09 |
|                   | 50%                         | 408.00 | 398.06                      | 12.30 | 12.91                         | 0.803 | 0.920                  | 13.20 | 13.84 |
| 3. m <sup>3</sup> | 100%                        | 011.77 | 010.40                      | 14.01 | 14.81                         | 0.793 | 0.028                  | 10.31 | 10.33 |
|                   | 75%                         | 483.78 | 488.00                      | 14.10 | 14.13                         | 0.777 | 0.790                  | 14.87 | 14.82 |
|                   | 50%                         | 472.78 | 473.00                      | 13.39 | 13.70                         | 0.882 | 0.774                  | 14.27 | 14.47 |
| 4. m <sup>3</sup> | 100%                        | 494.33 | 493.06                      | 14.13 | 13.91                         | 1.304 | 1.022                  | 10.48 | 10.43 |
|                   | 75%                         | 041.78 | 003.77                      | 14.80 | 14.73                         | 1.284 | 1.331                  | 17.13 | 17.07 |
|                   | 50%                         | 070.22 | 079.11                      | 10.33 | 10.11                         | 1.177 | 1.194                  | 17.49 | 17.30 |
| <b>LSD at 5%</b>  |                             | 72.20  | 70.32                       | 1.93  | 1.09                          | 0.199 | 0.133                  | 1.78  | 0.99  |
| <b>FYM rates</b>  | <b>Foliar spray sources</b> |        |                             |       |                               |       |                        |       |       |
| 2. m <sup>3</sup> | Control                     | 373.89 | 371.06                      | 12.07 | 12.43                         | 0.999 | 1.084                  | 13.07 | 13.02 |
|                   | Hi-Fertil 1                 | 433.84 | 439.00                      | 13.09 | 13.40                         | 0.891 | 0.990                  | 13.99 | 14.40 |
|                   | Ferty More                  | 478.18 | 470.78                      | 13.80 | 13.73                         | 0.821 | 0.919                  | 14.71 | 14.70 |
| 3. m <sup>3</sup> | Control                     | 429.89 | 427.06                      | 13.27 | 13.41                         | 0.794 | 0.793                  | 14.07 | 14.20 |
|                   | Hi-Fertil 1                 | 494.33 | 003.89                      | 14.18 | 14.40                         | 0.831 | 0.744                  | 10.01 | 10.00 |
|                   | Ferty More                  | 034.00 | 047.06                      | 14.00 | 14.83                         | 0.818 | 0.040                  | 10.37 | 10.38 |
| 4. m <sup>3</sup> | Control                     | 472.11 | 407.89                      | 13.76 | 13.03                         | 1.018 | 1.711                  | 10.28 | 10.10 |
|                   | Hi-Fertil 1                 | 047.89 | 000.22                      | 14.87 | 14.72                         | 1.384 | 1.497                  | 17.20 | 17.12 |
|                   | Ferty More                  | 091.33 | 703.22                      | 10.78 | 10.08                         | 0.902 | 0.938                  | 17.08 | 17.02 |
| <b>LSD at 5%</b>  |                             | 21.30  | 20.08                       | 0.30  | 0.00                          | 0.412 | 0.401                  | 0.24  | 0.22  |
| <b>NPK levels</b> | <b>Foliar spray sources</b> |        |                             |       |                               |       |                        |       |       |
| 100%              | Control                     | 430.44 | 420.22                      | 13.27 | 13.33                         | 1.131 | 1.230                  | 14.40 | 14.07 |
|                   | Hi-Fertil 1                 | 497.39 | 004.06                      | 14.20 | 14.12                         | 1.073 | 1.127                  | 10.32 | 10.20 |
|                   | Ferty More                  | 040.19 | 047.11                      | 14.82 | 14.80                         | 0.907 | 0.793                  | 10.73 | 10.09 |
| 75%               | Control                     | 414.00 | 417.44                      | 13.02 | 13.12                         | 1.111 | 1.101                  | 14.13 | 14.28 |
|                   | Hi-Fertil 1                 | 497.11 | 007.22                      | 14.00 | 14.14                         | 1.032 | 1.023                  | 10.08 | 10.17 |
|                   | Ferty More                  | 031.33 | 043.89                      | 14.78 | 14.71                         | 0.813 | 0.814                  | 10.09 | 10.02 |
| 50%               | Control                     | 421.44 | 419.33                      | 12.81 | 12.92                         | 1.079 | 1.102                  | 13.88 | 14.03 |
|                   | Hi-Fertil 1                 | 482.00 | 487.33                      | 13.83 | 14.17                         | 1.012 | 0.987                  | 14.84 | 10.10 |
|                   | Ferty More                  | 032.00 | 034.00                      | 14.43 | 14.74                         | 0.821 | 0.790                  | 10.20 | 10.43 |
| <b>LSD at 5%</b>  |                             | 21.20  | 29.09                       | 0.13  | 0.17                          | 0.081 | 0.104                  | 0.14  | 0.13  |

Concerning the effect of the interaction between NPK levels and macronutrients foliar spray sources on total yield and its components, data in Table (7) indicated that the interaction had a positive significant effects on total yield and its components in both seasons. The highest total yield, tuber weight/plant and marketable yield were obtained from plants received 75 or 50 % NPK level and sprayed with Ferty More in both seasons. It is notable that, the differences between tubers weight/plant and total yield were not significant when plants sprayed with Ferty More and fertilized by 75% or 50% NPK levels in both seasons.

Regarding, the interaction effect of the studied factors (FYM rates, NPK levels and macronutrients foliar spray sources) on total yield and its components, data presented in Table (8) reveal that tubers weight/plant, marketable yield, unmarketable yield and total yield were affected by the three ways interactions, in both seasons. Generally, the highest tuber weight/plant and total yield were obtained from plants fertilized by 40 m<sup>3</sup>/fed. FYM, 50% or 75% NPK levels and sprayed with Fert More as a source of foliar NPK followed by the interaction among 40 m<sup>3</sup>/fed. FYM, 50% or 75% NPK levels and sprayed with Hi-Fertil 1. The values in both treatments were not significant differences with each other. These results were true during both seasons. These results may be attributed to the favorable effect of the combination between the three factors of this study which results in good healthy of plant vegetative growth parameters (Tables 3, 4 and 5) consequently, increasing total yield.

**Table (8): Total yield and its components of potato plants as affected by the interaction among FYM rates, NPK levels and macronutrients foliar spray sources during 2008/2009 (S1) and 2009/2010 (S2) winter seasons.**

| Characters        |                   |                      | Tuber weight/plant (g) |        | Marketable yield (ton/fed.) |       | Unmarketable yield (ton/fed.) |       | Total yield (ton/fed.) |       |       |
|-------------------|-------------------|----------------------|------------------------|--------|-----------------------------|-------|-------------------------------|-------|------------------------|-------|-------|
|                   |                   |                      | S1                     | S2     | S1                          | S2    | S1                            | S2    | S1                     | S2    |       |
| Treatments        |                   |                      | S1                     | S2     | S1                          | S2    | S1                            | S2    | S1                     | S2    |       |
| FYM               | NPK               | Foliar spray sources |                        |        |                             |       |                               |       |                        |       |       |
| 20 m <sup>3</sup> | 100%              | Control              | 409.00                 | 393.67 | 12.80                       | 12.96 | 1.133                         | 1.226 | 13.98                  | 14.19 |       |
|                   |                   | Hi-fertil 1          | 477.80                 | 476.00 | 13.78                       | 13.77 | 0.903                         | 1.120 | 14.68                  | 14.89 |       |
|                   |                   | Ferty More           | 009.22                 | 019.00 | 14.49                       | 13.96 | 0.823                         | 0.970 | 10.31                  | 14.94 |       |
|                   | 75%               | Control              | 300.00                 | 374.33 | 11.89                       | 12.30 | 0.963                         | 1.000 | 12.80                  | 13.30 |       |
|                   |                   | Hi-fertil 1          | 422.33                 | 438.00 | 13.00                       | 13.31 | 0.916                         | 0.943 | 13.97                  | 14.26 |       |
|                   |                   | Ferty More           | 470.33                 | 472.33 | 13.76                       | 13.74 | 0.833                         | 0.910 | 14.09                  | 14.60 |       |
|                   | 50%               | Control              | 307.67                 | 306.67 | 11.46                       | 12.03 | 0.900                         | 0.976 | 12.36                  | 13.01 |       |
|                   |                   | Hi-fertil 1          | 411.33                 | 403.00 | 12.46                       | 12.22 | 0.803                         | 0.923 | 13.26                  | 14.10 |       |
|                   |                   | Ferty more           | 400.00                 | 436.00 | 13.14                       | 13.49 | 0.806                         | 0.876 | 13.94                  | 14.36 |       |
|                   | 30 m <sup>3</sup> | 100%                 | Control                | 440.00 | 422.67                      | 13.87 | 14.00                         | 0.733 | 0.743                  | 14.00 | 14.74 |
|                   |                   |                      | Hi-fertil 1            | 030.33 | 043.00                      | 14.67 | 14.91                         | 0.816 | 0.016                  | 10.49 | 10.42 |
|                   |                   |                      | Ferty more             | 064.67 | 070.67                      | 10.00 | 10.47                         | 0.930 | 0.426                  | 10.93 | 10.89 |
| 75%               |                   | Control              | 431.33                 | 423.00 | 13.00                       | 12.44 | 0.826                         | 0.833 | 14.33                  | 14.27 |       |
|                   |                   | Hi-fertil 1          | 492.67                 | 496.33 | 14.17                       | 14.28 | 0.773                         | 0.700 | 14.94                  | 14.98 |       |
|                   |                   | Ferty more           | 027.33                 | 044.67 | 14.62                       | 14.78 | 0.703                         | 0.086 | 10.32                  | 10.70 |       |
| 50%               |                   | Control              | 418.33                 | 424.00 | 12.43                       | 12.70 | 0.923                         | 0.903 | 13.30                  | 13.60 |       |
|                   |                   | Hi-fertil 1          | 460.00                 | 472.33 | 13.69                       | 14.02 | 0.903                         | 0.766 | 14.60                  | 14.78 |       |
|                   |                   | Ferty more           | 010.00                 | 024.33 | 14.04                       | 14.34 | 0.820                         | 0.723 | 14.86                  | 14.97 |       |
| 40 m <sup>3</sup> |                   | 100%                 | Control                | 442.33 | 424.33                      | 13.09 | 12.98                         | 1.226 | 1.836                  | 14.22 | 14.82 |
|                   |                   |                      | Hi-fertil 1            | 494.00 | 494.67                      | 14.31 | 13.79                         | 1.470 | 1.746                  | 10.78 | 10.03 |
|                   |                   |                      | Ferty more             | 046.67 | 001.67                      | 14.98 | 14.96                         | 0.966 | 0.983                  | 10.90 | 10.94 |
|                   | 75%               | Control              | 400.67                 | 412.00 | 13.66                       | 13.63 | 1.043                         | 0.070 | 10.20                  | 10.21 |       |
|                   |                   | Hi-fertil 1          | 073.33                 | 084.33 | 14.93                       | 14.84 | 1.406                         | 1.476 | 16.34                  | 16.32 |       |
|                   |                   | Ferty more           | 096.33                 | 714.67 | 10.90                       | 10.71 | 0.903                         | 0.446 | 16.80                  | 16.60 |       |
|                   | 50%               | Control              | 488.33                 | 477.33 | 14.03                       | 13.99 | 1.386                         | 1.426 | 10.92                  | 10.42 |       |
|                   |                   | Hi-fertil 1          | 076.33                 | 086.67 | 10.30                       | 10.24 | 1.276                         | 1.270 | 16.62                  | 16.01 |       |
|                   |                   | Ferty more           | 731.00                 | 743.33 | 16.10                       | 16.08 | 0.836                         | 0.886 | 16.94                  | 16.97 |       |
|                   | LSD at 5 %        |                      |                        | 39.06  | 33.16                       | 0.07  | 0.11                          | 0.143 | 0.137                  | 0.20  | 0.11  |

## CONCLUSION

From the results of this study, it could be concluded that, application 40 m<sup>3</sup>/fed FYM + 50% NPK level with spraying the plants with Ferty More (5-3-43, NPK) or Hi-Fertil 1(19-19-19, NPK) are the recommended treatments for increasing potato yield, improving tuber quality of potato, lowering cost production (as a result of saving half of the added NPK-fertilizer) and will also decrease the pollution of environment under similar conditions to this work.

## REFERENCES

- Abd El-Aal, S. F.; A.M. Shaheen;. and F. A. Rizk (2008).The effect of foliar application of GA3 and soil dressing of NPK at different levels on the plant productivity of potatoes (*Solanum tuberosum* L.). Res. J of agric and Biological Sci., 4(5): 384-391.
- Abd El-Kader, A.I. (2002). Effect of some organic and mineral fertilizers on some potato cultivars (*Solanum tuberosum* L.) M. Sc. Thesis Fac. of Agric., Mans. Univ., Egypt.
- Abdel-Ati, Y.Y. (1998). Yield and quality of potato as affected by phosphorus chicken manure and seed tuber size. Assiut J. of Agric. Sci. 29(5): 129-147.
- Abou-Hussein, S.D. (1995). Studies on potato fertigation in newly reclaimed lands. M.Sc.Thesis Fac. Agric. Ain-Shams Univ., Cairo, Egypt.
- Abou-Hussein, S.D.; I. El-Oksh; T. El-Shorbagy and U.A. El-Bahiry (2002 b). Effect of chicken manure, compost and biofertilizers on vegetative growth, tuber characteristics and yield of potato crop. Egypt. J. Hort., 29 (1):135-149.
- Abou-Hussein, S.D.; U.A. El-Bahiry , I. El-Oksh and M.A. Kalafallah (2002 a). Effect of compost, biofertilizer and chicken manure on nutrient content and tuber quality of potato crops. Egypt. J. Hort. 29(1):117-133.
- Abou-Hussein, S.D.; T. El-Shorbagy and A.F. Abou-Hadid (2003). Effect of cattle and chicken manure with or without mineral fertilizers on tuber quality and yield of potato crops. Acta Hort. (ISHS), 608 : 95 – 100.
- Al-Betar, A. and M Abdou (2010). Effect of foliar nutrition on growth, green and fodder yield of sweet corn. J. Plant Production, Mans. Univ., 1(5): 769-777.
- Ali, M. and J. P. Mishra (2001). Effect of foliar nutrition of boron and molybdenum on chickpea. Ind. J. pulses 14:41-43. Alvaret- Tinault MC, Leal A, Martinez LR. 1980. Iron-manganese interaction and its relation to boron levels in tomato plants. Plant Soil, 55:377-388.
- Al-Moshileh, A.M. and A.E. Moftah, (2005). Effects of chemical and biological fertilizers on growth, yield and tuber quality of potatoes grown in sandy soils under semi-arid conditions. J. Agric. Sci., Mans. Univ., 30(9): 5459-5470.
- Arisha, H. M. and A. Bardisi. (1999). Effect of mineral and organic fertilizers on growth, yield and tuber quality of potato under sandy soil conditions. Zagazig J. Agric. Res., 26 (2) : 391 -409.

- Awad, E.M.; E.A.A. Tartoura; H.M. El-Foly and A.I. Abdel-Fattah (2002). Response of potato growth, yield and quality to farmyard manure, sulphur and gypsum levels application. Conf. Hort. Sci., 10–12 Sept. 2002, Kafr El-Sheikh, Tanta Univ, Egypt.
- Ayyub, C. M. ; A. Muhammad; A Waqar;Z Khurram, and M. I. Khan (2006). Response of potato to nitrogen application methodologies. Pakistan, J. Agric. Sci.; 43(1/2): 45-49.
- Black, W.N. and R.P. White (1973). Effects of nitrogen, phosphorus and potassium manure factorially applied to potatoes in long term study. Candian J. Soil Sci., 53:205.
- Boligowa, E. and S. Dzienia (1999). Efferctiveness of soil tillage systems and organic fertilization for potato. Folia Univ., Agric. Stetinensis, Agric. (74): 191-195.
- Boote, J. B.; R. N Gallaher.; W.K Robertson., and L. C. Hammons (1978). Effect of foliar fertilization on photosynthesis, leaf nutrition and yield of soybean. Agron J., 70: 786-791.
- Borin, M.; C. Giupponi and I. Osele (1987). The effects of organic and mineral fertilizers and soil type on potato tuber formation. Information, Agrario, 43: 91-92.
- Colla, G.; M. Cardare, Y. Roupael; A. Fiorilla; B. Cirica (2005). Response of potatoes to organic fertilizer rate, yield, tuber quality and nutrient uptake. Acta, Hort; , 687:73-77.
- Doeing, H.W. (1986). Foliar fertilization. An effect method for controlling micronutrients deficiencies on calcareous and saline soil in arid lands. In: Proc. Symp. Application of Special Fertilizers. Alex. 21-23 Feb., Egypt.
- Dorobantu, N.; V. Birnaure; R. Gaina and L. Geamanu (1989). Influence of fertilizers on total leaf area of potatoes and tuber yields. Agronomie, 32 (1) : 54 – 65.
- El-Banna, E. N. and H. Z. Abd El-Salam (2000). Effect of rock phosphate and superphosphate application with organic manures on growth, yield and quality of potato (*Solanum tuberosum* L.) J. Agric. Sci. Mans. Univ., 25 (8) : 5343 – 5352.
- El-Dissoky, R.A. (2008). Studies on the use efficiency of potassium and organic fertilizers on potatoes and its role in improving soil properties. Ph. D. Thesis. Fac. of Agric., Mans, Univ., Egypt.
- El-Kassas, A.L. (2005). Effect of organic manures and bio fertilizers on growth, yield and fruit quality of tomato under low plastic tunnels in north sinal. J. Agric. Sci. Mans, Univ., 30 (8): 4729 – 4751.
- El-Morsy, A.H.A.; E. N. El-Banna and M.M.B. Shokr (2006). Effect of sources of organic manures and foliar spray with some micronutrients on productivity and quality of potato crop (*Solanum tuberosum* L.). J. Agric. Sci. Mans Univ., 31 (6): 3859 – 3868.
- El-Nagar, E.M. (1996). Effect of applying some organic residues to sandy and calcareous soils on growth and composition of some plant M. Sc. Thesis Fac. of Agric., Mans, Univ., Egypt.

- El-Sirafy Z. M.; K. A. Abbady; A. M. El-Ghamry and R. A. El- Dissoky. (2008). Potato yield quality, quantity and productivity as affected by soil and foliar potassium application. *Res. J. of Agric. and Biological Sci.*, 4(6): 912-922.
- Gardener, F.P.; R.B. Peraco and R.L. Mitchell (1985). *Physiological of Crop Plants*. First Ed. Iowa State Univ. Press, Ames, 277 p.
- Gomez, K.A. and A.A. Gomez (1984). "Statistical procedure for Agric. Res." 2<sup>nd</sup>. ED. John-Wiley and Sons. Inc. New Yourk, 680 pages.
- Jackson, M.L. (1973). "Soil Chemical Analysis". Prentice-Hall of India, Private limited, New Delhi, PP: 111-204.
- Jasiwal, V. P. (1995). Effect of cost reduction schedules on yield and economics of potato production. *J. Indian potato Assoc.*, 22(3/4): 162-165.
- Joudu, J.; A. Lohmus,; P. Laaniste.; V. Eremeev.; and A. Makke, (2001). The effect of growth-period and calcium nitrate fertilization on potato yield and relative leaf chlorophyll content. *Proceedings, Inter. Conf. Sustainable, Agric. in Baltic-States, Tartu, Estonia*, (6): 28-30 and 44-47.
- Karadogan, T. (1996). Effects of manure and fertilizer application on yield, yield components and quality of potatoes. *Turkish J. Agric. & Forestry*, 19 (5) : 373 – 377. (C.A. Potato Abst., 21 (1) : 19).
- Kolbe, H.; S. Meineke and W.L. Zhang (1995). Differences in organic and mineral fertilization on potato yield and chemical composition compared to model calculations *Agrobiol. Res.*, 48 (1): 63 – 73.
- Koller, H. R. C. (1972). Leaf area- leaf weight relationship in the soybean. *Canopy Crop Sci.*, 12: 180-183.
- Kushwah, S. S. and R. N. S. Banafar (2003). Influence of different N and P levels with and without bio-fertilizer on N, P content, uptake and yield of potato cv. Kufri jyoti. *J. of the Indian Potato Association*, 30: 3/4, 321-324.
- Maier, N.A.; A.P. Dahlendurg and C.M.J. Williams (1994). Effect of nitrogen, phosphorus and potassium on yield and petiolar nutrient concentration of potato cv. Kennebec and Atlantic. (C.A. potato Abst., 20(3): 133).
- Marschner, H. (1995). *Mineral Nutrition of Higher plants*. 2<sup>nd</sup> Ed. Academic press, Harcourt Brace and company, Publishers. London, San Diego, New York, Boston, Sydney, Tokyo, Toronto, 864 pp.
- Minhas, R.S. and A. Sood (1994). Effect of inorganic and organic materials on the yield and uptake by three crops in a rotation on an acid alfisol. *J. Indian Soc. Soil Sci.*, 42 (2): 257-260.
- Rabie, A. R. (1996). Effect of some potato cultural practices on potato production for processing. M. Sc. Thesis. Fac. of Agric., Ain Shams Univ., Cairo.
- Radwan, E.A. and A.A. Tawfik (2004). Effect of sulphur, manganese and zinc on growth, yield and quality of potato. *J. Agric .Sci. Mans. Univ.*, 29 (3) : 1423 – 1431.

- Sadej, W.; K. Przekwas and J. Bartoszewicz (2004). Effects of long term differentiated fertilization on changes in the yield and chemical composition of potato tubers. Ann. Univ. Mariae Curie Skodowska Sectio E. Agric., 59(1): 83-92.
- Sarhan, S. H.; H.K. Zaki and E. N. El-Banna (2004). Impact of organic and inorganic fertilization on yield, tuber contents and some heavy metals concentration in potato tubers. J. Agric. Sci. Mans. Univ., 29(5): 2753-2760.
- Singh, S.P.; V. S. Kushwah (2006). Effect of intergrated use of organic and inorganic sources of nutrients on potato (*Solanum tuberosum* L.). Product. Indian J. Agronomy. 51(3): 236-238.
- Singh, S. K.; V. K. Gupta (2005). Influence of farmyard, nitrogen and biofertilizer on growth, tuber yield of potato under rainfed condition in East Khasi Hill District of Meghalaya. Agric. Sci. Digest., 25(4):281-283.
- Singh-S.P; V. Singh; R. Lakhan and R. Lakhan (1996). Effet of phosphorus and farmyard manure application on yield, content and up take of nitrogen, phosphorus and sulphur (*Solanum tuberosum* L.) Indian j. of Agronomy. 41 (4): 630–632.
- Suresh, S. and K.K. Ramanathan (2001). Integrated potassium nutrition for sweet potato. Haryana, J. Hort. Sci., 30(1/2):100-101.

### إستجابة نمو ومحصول وقابلية تخزين البطاطس لبعض معدلات السماد العضوي والرش ببعض العناصر الكبرى:-

#### ١- النمو الخضري والمحصول ومكوناته.

هالة عبدالغفار السيد\*، عبدالله حلمي على المرسي\*\* و حمادة ماهر بدير المتولى\*\*

\* قسم الخضار والزينة- كلية الزراعة- جامعة المنصورة - مصر

\*\* قسم بحوث البطاطس والخضار خضرية التكاثر - معهد بحوث البساتين - مركز البحوث الزراعية ، الجيزة - مصر.

تُفذت تجربتان حقليتان على نباتات البطاطس صنف اسبونتتا في قرية كفر ميت فارس بالقرب من المنصورة، محافظة الدقهلية خلال موسمي الزراعة الشتوية ٢٠٠٨/٢٠٠٩ و ٢٠٠٩/٢٠١٠ م وذلك لدراسة تأثير استخدام بعض معدلات سماد المزرعة (٢٠ م<sup>٢</sup> ، ٣٠ م<sup>٢</sup> و ٤٠ م<sup>٢</sup> للفدان) وبعض مستويات النتروجين والفوسفور والبوتاسيوم (١٠٠٪ ، ٧٥٪ و ٥٠٪ من المعدل الموصى به للفدان) كل منها منفرداً أو مع الرش الورقي لنبات البطاطس ببعض مصادر العناصر الكبرى (كنترول، هاي فرتيل ١ و فيرتي مور) على النمو الخضري، المحصول ومكوناته.

استخدم لاجراء التجربة تصميم القطاعات الكاملة العشوائية بنظام القطع المنشقة مرتين في ثلاث مكررات وتم توزيع معدلات سماد المزرعة في القطع الرئيسية و مستويات النتروجين والفوسفور والبوتاسيوم في القطع الشقية وخصبت القطع التحت شقية لمعاملات الرش الورقي ببعض مصادر العناصر الكبرى. ويمكن تلخيص النتائج المتحصل عليها فيما يلي:-

أوضحت النتائج أن التسميد البلدي بصفة عامة أثر معنوياً على صفات النمو الخضري والمحصول ومكوناته. ولقد أدى استخدام ٤٠ م<sup>٢</sup> للفدان من سماد المزرعة متبوعاً بـ ٣٠ م<sup>٢</sup> للفدان إلى زيادة في طول النبات، عدد الأوراق للنبات، المساحة الورقية للنبات والوزن الطازج للنبات. وكذلك أدى إلى زيادة معنوية للمحصول الكلي للفدان ومتوسط وزن الدرنة للنبات.

أوضحت النتائج أيضاً، أن زيادة السماد الكيماوى المستخدم حتى ١٠٠٪ من المعدل الموصى به للفدان أدى إلي زيادة معنوية في كل صفات النمو الخضري ماعدا الوزن الجاف للنبات، وكذلك زيادة متوسط وزن الدرناات لكل نبات، المحصول التسويقي و المحصول الكلي للفدان.

وتشير النتائج ان الرش الورقى للنباتات بالمصادر المختلفة للعناصر الكبرى ادى إلي زيادة معنوية في جميع الصفات المدروسة. معظم القياسات الخضريه زادت عند رش النباتات بمركب هاى فرتيل ١ (١٩-١٩ ن فو بو)، بينما أدى رش النباتات بمركب فيرتى مور (٥-٣-٤٣، ن فو بو) إلي زيادة معنوية في كل من الوزن الجاف لعرش النبات و المحصول الكلي و المحصول التسويقي و كذلك متوسط وزن الدرناات للنبات.

وعموما كان تأثير التفاعل بين العوامل الثلاثة المدروسة (معدلات السماد العضوى، مستويات التسميد الكيماوى والرش الورقى بمصادر العناصر الكبرى) واضحاً على جميع الصفات المدروسة. وكانت أفضل النتائج باستخدام ٤٠ م<sup>٢</sup> للفدان من سماد المزرعة و التسميد الكيماوى بمستوى ٥٠٪ من الموصى به للفدان مع الرش الورقى بمركب الـ فيرتى مور ويليه التفاعل باستخدام ٤٠ م<sup>٢</sup> للفدان من سماد المزرعة و التسميد الكيماوى بمستوى ٥٠٪ من الموصى به للفدان مع الرش الورقى بمركب هاى فرتيل ١، حيث أدى هذا التفاعل إلي زيادة فى وزن الدرناات للنبات و المحصول الكلى و المحصول التسويقي للفدان كما أدى إلي تخفيض المحصول غير التسويقي. ولذلك يمكن التوصية باستخدام هذه المعاملات، لرفع إنتاجية البطاطس وتحسين جودة الدرناات، وكذلك خفض تلوث البيئة تحت ظروف محافظة الدقهلية.

#### قام بتحكيم البحث

كلية الزراعة – جامعة المنصورة  
كلية الزراعة – جامعة بنها

أ.د / السيد احمد احمد طرطوره  
أ.د / نادية سعد شفشق