



## Diterpenoids Profile of *E. paralias* and *E. geniculata* Using UPLC-ESI/MS Spectrometry



Afaf E. Abdel-Ghani<sup>a</sup>, Sayed A. El-Toumy<sup>b</sup>, Wagdi I. A. El-DougDoug<sup>c</sup>, Hanaa M. Hassan<sup>d\*</sup>, Wafaa H. B. Hassan<sup>e</sup>

<sup>a,e</sup>Department of Pharmacognosy Faculty of Pharmacy, Zagazig University, 44519 Zagazig

<sup>b</sup>Chemistry of Tannins Department, National Research Centre, Dokki, Cairo, 12622 Egypt

<sup>c</sup>Department of Chemistry, Faculty of Science, Benha University, 13518 Benha

<sup>d</sup>Pharmacy department Benha Educational Hospital, 13518 Benha

<sup>e</sup>Department of Pharmacognosy Faculty of Pharmacy, Zagazig University.

### Abstract

Euphorbia species contain series of macrocyclic diterpenoids which present as mono, di and sometimes triesters and were reported to possess several and diverse biological activities. The aim herein was to use the ultra-performance coupled to the positive ion mode electron spray ionization mass spectrometry (UPLC-ESI-MS) as an analytical tool for rapid screening and tentative identification of diterpenoids in the methylene chloride fractions of *Euphorbia paralias* and *Euphorbia geniculata*. The identification of compounds was based on ESI-MS and ESI-MS/MS fragmentation pattern, in addition to comparison with reported literature. Five classes were determined including tigliane, ingenane, lathyrane, daphnane and jatrophone diterpenoids esters. Each class of the detected diterpenoids obtained its characteristic ESI-MS/MS fragmentation which facilitates its identification. As a result, a total of fifty-two diterpenoids were identified and characterized in *Euphorbia paralias* and *Euphorbia geniculata*. All of these compounds are first report in the two plants by this method. Peak 35 was tentatively identified as ingenol-20-hydroxyheptanoate which expected to be a new compound. The results demonstrated that the used method could be a rapid, and an effective analytical tool for screening and characterization of diterpenoids in the complex systems of plant extracts.

Keywords: Euphorbia, Terpenoids, UPLC-ESI-MS, ingenanes, and tiglianes.

### 1. Introduction

The genus *Euphorbia* is one of the largest genera with more than 2000 species worldwide [1]. *Euphorbia* species are characterized by the presence of a vast array of macrocyclic diterpenoids with various types of carbon skeletons. Several hundred diterpene compounds were isolated from the *Euphorbia* species with more than twenty different skeletons [2,3] as jatrophone (5/12-bicyclic ring system), ingenane (5/7/7/3-tetracyclic ring system), tigliane (5/7/6/3 tetracyclic ring structure), daphnane (5/7/6-tricyclic

system), segetane (5/9/5 tricyclic ring system) and skeleton), myrsinane (5/7/6-fused tricyclic ring lathyrane (5/11/3-tricyclic ring system) [2]. These macrocyclic diterpenoids are valuable due to their diverse biological and therapeutic activities as cytotoxic [4], antitumor [5,6], multi-drug-resistance-reversing [7], immuno-modulatory [8], antimicrobial [9], anti-inflammatory [10], pesticidal, pro-inflammatory [3], molluscicidal [11], vascular-relaxing, and neuroprotective activities [10,12]. So the authors aimed in this work to study the UPLC-ESI-MS/MS analysis of nonpolar fractions of *E. paralias* and *E. geniculata* to detect their diterpene constituents.

\*Corresponding author : Hanaa Mohamed Hassan, e-mail: [hanaazain111@yahoo.com](mailto:hanaazain111@yahoo.com).

Receive Date: 03 March 2020, Revise Date: 31 March 2020, Accept Date: 22 April 2020

DOI: 10.21608/ejchem.2020.25113.2486

©2020 National Information and Documentation Center (NIDOC)

## Material and methods:

### Plant material:

The aerial parts of *Euphorbia paralias* L. (Euphorbiaceae) used in this work was collected in the flowering stage on May 2015 from the North beach of Alexandria, Egypt, while *Euphorbia geniculata* Ortega was collected in August 2015 from the roadsides and beside farms in the vicinity of Benha, Qalubya Governorate, Egypt. The identification was kindly verified by Dr. Ahmed Abd El-Razik respective lecturer of Plant Taxonomy, Department of Botany, Faculty of Science, Banha University, Banha, Qalubya Governorate, Egypt. The voucher specimens (no. S303 (for *E. paralias*) and S304 (for *E. geniculata*)) were deposited in National Research Center, El Doki, Egypt. The aerial parts of *E. paralias* L. and *E. geniculata* Ortega were air-dried and ground into coarse particles till use.

### Extraction of plant material:

The air-dried powdered aerial parts of *E. paralias* and *E. geniculata* (200 gm of each) were extracted by cold maceration with 70 % methanol (1 L for each) till complete exhaustion. The combined methanolic extract of each was evaporated under reduced pressure at 45 °C to give 32 and 30 gm respectively, of greenish brown viscous residue. Each residue was suspended in MeOH-H<sub>2</sub>O mixture (150 ml, 1:9 v/v) and extracted with methylene chloride (5 x 100 ml). The combined methylene chloride fractions of *E. paralias* and *E. geniculata* were separately washed with distilled water, dried over anhydrous sodium sulphate, then distilled off under vacuum at 45 °C to afford 1.5 and 1.2 g respectively.

### UPLC- ESI- MS/MS analysis:

The methylene chloride fraction samples of *E. paralias* and *E. geniculata* were prepared as solutions of 100 µg/mL using HPLC grade methanol, filtered using a membrane disc filter (0.2 µm) then subjected to LC-ESI-MS analysis. The injection volume was 10 µL. The separation of samples was performed on UPLC instrument with ESI-MS detector equipped with reverse phase C-18 column (ACQUITY UPLC – BEH 2.1 × 50 mm, 1.7 µm). Mobile phase elution was made with the flow rate of 0.2 ml/min using gradient mobile phase comprising two eluents: eluent A is H<sub>2</sub>O acidified with 0.1 % formic acid and eluent B is MeOH acidified with 0.1 % formic acid. Elution was performed using the following gradient: 20 % B, 0-1 min; 20-90 % B, 1-18 min; 20 % B, 18-20 min. Mass spectra were detected in the ESI-MS positive ion mode between *m/z* 50-900 at 30 eV capillary cone and capillary voltage 3 kV. The peaks and spectra were processed using the Maslynx 4.1 software and tentatively identified by comparing its retention time

(Rt) and mass spectrum with reported data. For fragmentation collision energy 40 eV was used.

## Results:

Electrospray ionization mass spectroscopy is an important, sensitive and reliable technique used for the analysis and rapid tentative identification of various classes of compounds. This helped in studying the fragmentation pathways of five classes of diterpenoids, ingenane, daphnane, tiglane, lathyrane and jatrophane, in methylene chloride fractions of two *Euphorbia* species. As a result, a total of fiftytwo diterpenoids (Fig. 1-13, Table 1) in *E. paralias* and *E. geniculata* were screened. The tentative identification of these compounds was based on matching the molecular ion peaks detected in positive ESI-MS and MS<sup>2</sup> fragmentation pattern with the previous literature. The major compounds in *E. paralias* were phorbol-12-tetradecanoate (4.93 %), 20-Oxo-phorbol-12, 13-dibutyrate (4.67%) and 20-Oxo-13-deoxy phorbol-12-propionate (4.18%), while the major compounds in *E. geniculata* were 20-Oxo-phorbol-12, 13-dibutyrate, 20-Oxo-13-deoxy phorbol-12-propionate and 13-*O*-isobutyryl-12-deoxyphorbol-20-acetate in ratio of 10.4%, 8.87% and 7.74 % respectively.

### Identification of diterpenoids in Methylene Chloride Fractions of *E. paralias* and *E. geniculata*

Depending on the class of core structure and the substitution pattern, each class of diterpenoids gave either the molecular adduct ion [M<sup>+</sup>+Na] and sometimes [M<sup>+</sup>+H]/ [M<sup>+</sup>+K]. In the MS/MS of ingenanes (ingenol esters), the molecular adduct ion gave characteristic product ions corresponding to the loss of angelic, acetic, benzoic, or hydroxyheptanoic acid, followed by the loss of water to yield the prominent characteristic fragment ion at *m/z* 295 (ingenol residual structure), the fragment at *m/z* 267 was sometimes observed which showed the loss of C=O. The tiglane class (phorbol esters) produced fragment ions showing the subsequent losses of the acyl moieties as acetic, benzoic, decanoic acids, N-methyl anthranilic moiety etc., in addition to the loss of three molecules of water leaving the characteristic fragment ion at *m/z* 293 (phorbol residual structure). While lathyrane diterpenes showed [M<sup>+</sup>+H], [M<sup>+</sup>+Na] and [M<sup>+</sup>+K] in ESI-MS. In ESI-MS/MS spectrum of lathyrane diterpenes the loss of acyl moieties with the subsequent loss of water molecules were observed producing the diagnostic fragment ion at *m/z* 281 (for lathyrane skeleton). Furthermore, the jatrophane type showed the MS<sup>2</sup> fragment ions at *m/z* 327, 309, 299 and 281.




Figure 1: LC-ESI-MS profiles of methylene chloride fractions of *E. paralias* (A) and *Egeniculata* (B) (Positive mode).

Table 1: The diterpenoids identified in Methylene Chloride Fractions of *E. paralias* and *Egeniculata*:

Peak No	Rt (min)	[M <sup>+</sup> +Na] (M <sup>+</sup> )	Daughter fragments (m/z) of MS <sup>2</sup>	E.p % area	E.g % area	Proposed compound
<b>Ingenane diterpenoids (Ingenol esters):</b>						
1	0.74	453 (430)	353, 295	-	+	Ingenol-3- mebutate (Ingenol-3- angelate)
23	15.9	579 (556)	457, 335, 317, 313, 307, 295, 285	-	+	Ingenol-3,20-dibenzoate
27	17.32	579(556)	457, 335, 317, 313, 307, 295, 285	-	+	Ingenol-3,20-dibenzoate isomer
28	18.60	579(556)	457, 335, 317, 313, 307, 295, 285	-	1.17	Ingenol-3,20-dibenzoate isomer
29	18.86	579(556)	457, 335, 317, 313, 307, 295, 285	-	1.74	Ingenol-3,20-dibenzoate isomer
30	19.45	579 (556)	457, 335, 317, 313, 307, 295, 285	-	+	Ingenol-3,20-dibenzoate isomer
33	20.15	437 [M <sup>+</sup> +Na], 453 [M <sup>+</sup> +K](414)	353, 313, 295, 267, 255	-	1.19	3-Deoxy-3-oxoingenol 20-butenate
34	20.34	579[M <sup>+</sup> +Na]	457, 335, 317, 313, 307, 295, 285	-	+	Ingenol-3,20-dibenzoate isomer
35	20.82	515 [M <sup>+</sup> +K], 499[M <sup>+</sup> +Na], 477 [M <sup>+</sup> +H] (476)	313, 295, 267	1.63	-	Ingenol-20-hydroxyheptanoate
<b>Tigliane diterpenoids (Phorbol and phorbol esters)</b>						
2	0.82	541(518)	369, 351, 333, 329, 315, 311, 293	1.19	+	Phorbol-12-decanoate
3	1.69	387(364)	387, 369, 351, 333, 315, 311, 293	0.77	-	Phorbol
4	8.89	429 (406)	369, 351, 333, 329, 315, 311, 293	0.40	+	Phorbol-12-acetate
5	9.82	429 (406)	369, 351, 333, 329, 315, 311, 293	0.15	+	Phorbol-13-acetate
6	11.05	471 (448)	411, 351, 333, 329, 315, 311, 293	0.36	0.32	Phorbol-12,13-diacetate
7	11.50	471 (448)	411, 351, 333, 329, 315, 311, 293	-	+	Phorbol-12, 13-diacetate isomer
9	11.78	562 (539)	411, 502, 351, 333, 329, 315, 311, 293	0.30	0.44	12-O-(N-methylanthranilate) phorbol-13-acetate

10	11.80	562 (539)	411, 502, 351, 333, 329, 315, 311, 293	+	+	12- <i>O</i> -( <i>N</i> -methylantranilate) phorbol-13-acetate isomer
11	11.85	562 (539)	411, 502, 351, 333, 329, 315, 311, 293	+	+	12- <i>O</i> -( <i>N</i> -methylantranilate) phorbol-13-acetate isomer
13	12.30	583 (560)	467, 351, 333, 329, 315, 311, 293	0.45	-	Phorbol-12,13-dihexanoate
14	12.82	541 (518)	387, 369, 351, 333, 329, 315, 311, 293	+	+	Phorbol-12-decanoate isomer
15	13.03	531 (508)	489, 471, 371, 335, 317, 313, 295	+	+	13- <i>O</i> -phenylacetyl-12-deoxyphorbol-20-acetate
16	13.20	541 (518)	387, 369, 351, 333, 329, 315, 311, 293	+	-	Phorbol-12-decanoate isomer
17	14.10	429 (406)	369, 351, 333, 329, 315, 311, 293	0.38	+	Phorbol-12-acetate isomer
19	15.02	597 (574)	369, 351, 333, 329, 315, 311, 293	4.93	-	Phorbol-12-tetradecanoate
20	15.18	597 (574)	369, 351, 333, 329, 315, 311, 293	1.97	-	Phorbol-13-tetradecanoate
21	15.63	695 (672)	523, 351, 333, 329, 315, 311, 293	0.95	+	Phorbol-12,13-didecanoate
22	15.82	429 (406)	369, 351, 333, 329, 315, 311, 293	2.18	-	Phorbol-13-acetate isomer
24	16.08	695 (672)	523, 351, 333, 329, 315, 311, 293	3.18	-	Phorbol-12,13-didecanoate isomer
25	16.78	597 (574)	369, 351, 333, 329, 315, 311, 293	1.97	-	Phorbol-13-tetradecanoate isomer
26	16.93	597 (574)	369, 351, 333, 329, 315, 311, 293	2.74	-	Phorbol-13-tetradecanoate isomer
31	19.61	483 (460)	335, 317, 313, 307, 295	-	7.74	13- <i>O</i> -isobutyryl-12-deoxyphorbol-20-acetate
36	21.01	425 (402)	347, 331, 291 (100%)	4.18	8.87	20-Oxo-13-deoxy phorbol-12-propionate
37	21.31	535 (512)	495, 395, 313, 295, 267	1.79	-	12-Deoxyphorbol-13, 20 diangelate
38	24.20	639 (616)	579, 411, 351, 333, 329, 315, 311, 293	-	+	12- <i>O</i> -Tetradecanoylphorbol-13-acetate (TPA)
39	24.28	525 (502)	507, 437, 349, 331, 309, 291	4.67	10.4	20-Oxo-phorbol-12,13-dibutyrate
40	24.78	525 (502)	507, 437, 349, 331, 309, 291	-	+	20-Oxo-phorbol-12,13-dibutyrate isomer
41	24.98	525 (502)	507, 437, 349, 331, 309, 291	-	+	20-Oxo-phorbol-12,13-dibutyrate isomer
42	25.0	639 (616)	411, 351, 333, 329, 315, 311, 293	-	+	12- <i>O</i> -tetradecanoylphorbol-13-acetate (TPA) isomer
43	25.61	531 (508)	489, 471, 371, 335, 317, 313, 295	-	1.97	13- <i>O</i> -phenylacetyl-12-deoxyphorbol-20-acetate
44	26.34	531 (508)	487, 471, 371, 335, 317, 313, 295	-	1.0	13- <i>O</i> -phenylacetyl-12-deoxyphorbol-20-acetate isomer
45	27.57	471 (448)	411, 351, 333, 329, 315, 311, 293	-	+	Phorbol-12, 13-diacetate isomer
46	27.67	541 (518)	387, 369, 351, 333, 329, 315, 311, 293	2.11	0.86	Phorbol-12-decanoate isomer
47	27.69	471 (448)	411, 351, 333, 329, 315, 311, 293	-	+	Phorbol-12,13-diacetate isomer
48	27.73	531 (508)	489, 471, 371, 335, 317, 313, 295	-	+	13- <i>O</i> -phenylacetyl-12-deoxyphorbol-20-acetate isomer
49	28.11	471 (448)	411, 351, 333, 329, 315, 311, 293	-	+	Phorbol-12, 13-diacetate isomer
51	28.58	695 (672)	387, 523, 351, 333, 329, 315, 311, 293	-	1.02	Phorbol-12,13-didecanoate isomer
52	28.70	541 (518)	369, 351, 333, 329, 315, 311, 293	2.18	4.47	Phorbol-12-decanoate isomer
<b><i>Lathyrane diterpenoids (Lathyrrol and hydroxylathyrrol esters)</i></b>						
8	11.69	539 [M <sup>+</sup> +H],561 [M <sup>+</sup> +Na], 577[M <sup>+</sup> +K](538)	461, 377, 317, 299, 281, 259	+	-	5-Pentanoyl-3-benzoyl-hydroxylathyrrol
12	12.14	539 [M <sup>+</sup> +H],561.3[M <sup>+</sup> +Na], 577[M <sup>+</sup> +K] (538)	461, 377, 317, 299, 281, 259	0.35	-	5-Pentanoyl-3-benzoyl-hydroxylathyrrol isomer

32	19.94	561 [M <sup>+</sup> +K], 545 [M <sup>+</sup> +Na], 523 [M <sup>+</sup> +H] (522)	463, 403, 375, 341, 299, 281 (100%)	0.37	-	Euphorbia factor L 3
18	14.16	647 [M <sup>+</sup> +K], 631 [M <sup>+</sup> +Na] (608)	591, 549, 531, 489, 471, 429, 411, 387, 369, 341, 327, 309, 299, 281	+	-	3,5,7,8,15, pentacetyl-2-hydroxy-9,14 dioxo-jatropha-6(16), -11(12)-diene
<i>Jatrophone diterpene ester</i>						
50	28.30	651 [M <sup>+</sup> +Na] (628)	507, 441, 371, 355, 333	-	1.1	Resiniferatoxin
<i>Daphnane diterpene ester</i>						

+ = present    - = absent    E.p = *Euphorbia paralias*    E.g = *Euphorbia geniculata*

### Ingenol esters:

Nine ingenol esters were tentatively identified in the methylene chloride fractions of *E. paralias* and *E. geniculata*. Eight compounds including ingenol-3-mebutate (ingenol-3-angelate), ingenol-3, 20-dibenzoate with its five isomers, in addition to 3-deoxy-3-oxoingenol 20-butenolate were identified in *Egeniculata* and one in *E. paralias* (ingenol-20-hydroxyheptanoate).

Peak 1 (Rt 0.74 min.) generated [M<sup>+</sup>+ Na] at *m/z* 453. It gave diagnostic fragment ions at *m/z* 353 [M<sup>+</sup>+Na-100 amu (angelic acid)] and 295 [M<sup>+</sup>+H-100 amu (angelic acid)-36 amu] corresponding to the loss of angelic acid ion and two molecules of water (*m/z* 36). Thus, it was characterized tentatively as ingenol-3-mebutate [13]. While peaks 23, 27-30 and 34 eluted at Rt. 15.90, 17.32, 18.60, 18.86, 19.45 and 20.34 min. were observed to possess the same ESI-MS<sup>1</sup> at *m/z* 579 [M<sup>+</sup>+Na]. The dominant ESI-MS<sup>2</sup> fragment ions at *m/z* 457 [M<sup>+</sup>+Na- C<sub>7</sub>H<sub>5</sub>O<sub>2</sub>], 335 [M<sup>+</sup>+Na-2 C<sub>7</sub>H<sub>5</sub>O<sub>2</sub>], and 317 [M<sup>+</sup>+Na-2 C<sub>7</sub>H<sub>5</sub>O<sub>2</sub>-H<sub>2</sub>O], showing the subsequent losses of one and two benzoic acid moieties then one molecule of H<sub>2</sub>O. the deprotonated base peak fragment ion at *m/z* 295 [M<sup>+</sup>+H-2C<sub>7</sub>H<sub>5</sub>O<sub>2</sub>-H<sub>2</sub>O] was corresponding to the loss of two molecules of benzoic acid and one molecule of water and corresponding to the residual ingenol skeleton. Consequently, peak 23 and its isomer were tentatively characterized as ingenol-3, 20-dibenzoate [13].

In addition, peak 35 (Rt, 20.82 min) displayed molecular ion peaks at *m/z* 515 [M<sup>+</sup>+K], 499 [M<sup>+</sup>+Na], 477 [M<sup>+</sup>+H]. Its further fragmentation resulted in [M<sup>+</sup>+H- C<sub>7</sub>H<sub>13</sub>O<sub>3</sub>-H<sub>2</sub>O] at *m/z* 313 for the loss of hydroxyheptanoic acid and one molecule of water. The prominent fragment ion at *m/z* 295 [M<sup>+</sup>+H-C<sub>7</sub>H<sub>13</sub>O<sub>2</sub>-2H<sub>2</sub>O] showing the loss of hydroxyheptanoic acid and two molecules of water and is characteristic for ingenol skeleton [4,13]. Furthermore, the remaining fragment at *m/z* 267 showed the loss of C=O group. By comparing the fragmentation pattern with the previous literature, the nucleus of peak 35 is ingenol while the substitution is hydroxyheptanoate which did not previously reported. Hence, peak 35 was tentatively identified as ingenol-20-hydroxyheptanoate which expected to be a new compound. Similarly, peak 33 eluted at Rt 20.15 min generated its ESI-MS at *m/z* 437 [M<sup>+</sup>+Na], and MS<sup>2</sup> fragment ion at *m/z* 353 [M<sup>+</sup>+Na-C<sub>4</sub>H<sub>5</sub>O<sub>2</sub>] which was produced by the loss of butenoic acid moiety, while the product ion at *m/z* 313 [M+H-C<sub>4</sub>H<sub>5</sub>O<sub>2</sub>-H<sub>2</sub>O] showed the loss of one molecule of water, furthermore the fragment ion at *m/z* 295 [M<sup>+</sup>+H-C<sub>4</sub>H<sub>5</sub>O<sub>2</sub> (86 amu) -2H<sub>2</sub>O (36 amu)] was resulted from the loss of two molecules of H<sub>2</sub>O leaving the residual ingenol skeleton [4,13]. The next fragment ion at *m/z* 267 [M<sup>+</sup>+H-C<sub>4</sub>H<sub>5</sub>O<sub>2</sub>-2H<sub>2</sub>O-CO] which showed the loss of carbonyl group was also observed. So peak 33 was identified as 3-deoxy-3-oxoingenol 20-butenolate [4,13].




Figure 2: The chemical structures of ingenol esters identified in methylene chloride fractions of *E. paralias* and *E. geniculata*





Figure 3: Positive ESI-MS/MS spectrum of ingenol-3, 20- dibenzoate (peak 23,  $m/z$  579[M<sup>+</sup>+Na])



Scheme 1: Characteristic ESI-MS/MS fragmentation pattern of ingenol-3, 20-dibenzoate in positive ion mode

#### Phorbol esters:

##### Phorbol and Phorbol monoesters

Fourteen (one phorbol, and thirteen phorbol monoester) diterpenoids were determined in both *Euphorbia* species including phorbol-12-tetradecanoate, phorpol-12-acetate, phorpol-13-acetate and phorpol-12-decanoate and their isomers. Peak 3, (Rt, 1.6 min), its ESI-MS spectrum showed a molecular ion peak at  $m/z$  387 [M<sup>+</sup>+Na] and MS<sup>2</sup> fragment ions at  $m/z$  369 [M<sup>+</sup>+Na-H<sub>2</sub>O], 351 [M<sup>+</sup>+Na-2H<sub>2</sub>O] and 333 [M<sup>+</sup>+Na-3H<sub>2</sub>O] were yielded due to losses of one, two and three molecules of water respectively. Base peak fragment at  $m/z$  293 [M<sup>+</sup>+H-4H<sub>2</sub>O] was observed. From this behavior in ESI-MS/MS and through comparison with literature, peak

3 was identified as phorbol [13]. Peaks 19, 20, 25 and 26 eluted at Rt, 15.02, 15.18, 16.78, 16.93 min exhibited the same MS<sup>1</sup> and MS<sup>2</sup> data. They gave molecular ion peak at  $m/z$  597 [M<sup>+</sup>+Na]. Series of MS<sup>2</sup> fragment ions showing the loss of tetradecanoic acid and several molecules of water at  $m/z$  369 [M<sup>+</sup>+Na-tetradecanoic acid], 351 [M<sup>+</sup>+Na-tetradecanoic acid-H<sub>2</sub>O], 333 [M<sup>+</sup>+Na-tetradecanoic acid-2H<sub>2</sub>O], 315 [M<sup>+</sup>+Na-tetradecanoic acid-3H<sub>2</sub>O] and 293 [M<sup>+</sup>+H-tetradecanoic acid-3 H<sub>2</sub>O] were observed. Accordingly peaks 19 was tentatively identified as phorbol-12-tetradecanoate while peak 20 as its isomers (peaks 25 and 26) were tentatively identified phorbol-13-tetradecanoate [13].

Peaks 2, 14, 16, 46 and 52 (Rt, 0.82, 12.82, 13.20, 27.67, 28.70) were tentatively identified as phoropol-12-decanoate and its four isomers [11]. Where the ESI-MS spectra displayed the same molecular ion peak at  $m/z$  541  $[M^+Na]$  and MS<sup>2</sup> spectra produced a fragment ion showing the loss of decanoate moiety at  $m/z$  387 followed by several fragment ions at  $m/z$  369  $[M^+Na-decanoate-H_2O]$ , 351  $[M^+Na-decanoate-2H_2O]$  and 333  $[M^+Na-decanoate-3H_2O]$  which showed repeated loss of 18 amu for loss of several molecules of water.

Peaks 4, 5, 17 and 22, (Rt, 8.89, 9.82, 14.1, 15.82 min) possessed the same molecular ion peak at  $m/z$  429  $[M^+Na]$  and displayed MS<sup>2</sup> fragment ions at  $m/z$  387  $[M^+Na-COCH_3]$ , 369  $[M^+Na-CH_3CO+H_2O]$ , 351  $[M^+Na-CH_3CO-2H_2O]$ , 333  $[M^+Na-CH_3CO-3H_2O]$  and 315  $[M^+Na-CH_3CO-4H_2O]$ . In addition, a base peak fragment at  $m/z$  293  $[M^+H-CH_3COOH-4H_2O]$  was observed. Thus peak 4 and its isomer (peak 17) were tentatively identified as phoropol-12-acetate, while peak 5 and its isomer (peak 22) were tentatively identified as phoropol-13-acetate [13]. Figure 5 showed the fragmentation pattern of phoropol-12-acetate (Fig. 5, Scheme 2).

#### Phorbol diesters:

Fifteen phorbol diesters were identified in both euphorbia species (13 in *E. paralias* and 15 in *E. geniculata*) as follow:

Peaks 6, 7, 45, 47 and 49 (Rt, 11.05, 11.50, 27.57, 27.69 and 28.11 min) produced the same molecular ion peak at  $m/z$  471  $[M^+Na]$  and all of their molecular ion peaks showed series of MS<sup>2</sup> fragment ions at  $m/z$  411  $[M^+Na-CH_3COOH]$ , 351  $[M^+Na-2CH_3COOH]$ , 333  $[M^+Na-2CH_3COOH-H_2O]$ , 315  $[M^+Na-2CH_3COOH-2H_2O]$ , 293  $[M^+H-2CH_3COOH-2H_2O]$ , which conferred the identification of these compounds as phorbol-12,13-diacetate and its four isomers (Fig. 4) [13]. In addition, peaks 9, 10 and 11 (Rt, 11.78, 11.80, 11.85 min.) showed the same  $[M^+Na]$  ion at  $m/z$  562. These peaks generated  $[M^+Na-CH_3COOH]$  at  $m/z$  502 by losing acetic acid, beside the fragment ion at  $m/z$  411 showing the loss of  $C_8H_8NO_2$  (*N*-methylantranilate). Other MS/MS fragments at  $m/z$  333 corresponded to subsequent loss of acetic acid moiety plus one molecule of water after loss of *N*-methylantranilate. Moreover, the fragment ion at  $m/z$  293 for  $[M^+H-2H_2O+CH_3COOH+C_8H_8NO_2]$  was produced. This fragmentation of peak 9 (Fig. 6) was in a good agreement with that for 12-*O*-(*N*-methylantranilate) phorbol-13-acetate (Fig. 6, scheme 3) which is known as sapintoxin D [13] and its isomers. While peaks 21, 24 and 51 eluted at Rt, 15.63, 16.08, 28.58 min were identified as phorbol-12, 13-didecanoate and its isomers from MS<sup>1</sup> and MS<sup>2</sup> spectral data (Table 1). Prominent MS<sup>1</sup> pseudo molecular ion peak at  $m/z$  695 was formed. The MS<sup>2</sup> product ions at  $m/z$  523

$[M^+Na-C_{10}H_{19}O_2]$  and 351  $[M^+Na-2C_{10}H_{19}O_2]$  corresponding to neutral losses of one and two molecules of decanoic acid. In addition to fragment ions at  $m/z$  333  $[M^+Na-2C_{10}H_{19}O_2-H_2O]$  and 315  $[M^+Na-2C_{10}H_{19}O_2-2H_2O]$  which showed the subsequent losses of one and two molecules of water. The base peak fragment at  $m/z$  293  $[M^+H-2C_{10}H_{19}O_2-2H_2O]$  was also observed. The mass data of these compounds are in a good agreement with those of phorbol-12, 13-didecanoate [13]. Peak 13 (Rt, 12.30) gave a pseudo-molecular ion peak at  $m/z$  583  $[M^+Na]$  and a fragment ion at  $m/z$  467  $[M^+Na-C_6H_{11}O_2]$  which involve loss of one hexanoic acid, further fragmentation gave  $m/z$  351  $[M^+Na-2C_6H_{11}O_2]$  for further loss of one hexanoic acid molecule. Moreover, series of fragments were produced at  $m/z$  333  $[M^+Na-2\text{ hexanoic acid-H}_2O]$ , 315  $[M^+Na-2\text{ hexanoic acid-2H}_2O]$  and 293  $[M^+H-2\text{ hexanoic acid-2H}_2O]$ . Consequently, compound 13 was identified as phorbol-12,13-dihexanoate [13].

Similarly, peaks 38 and 42 (Rt, 24.20, 25.0 min) exhibited the same molecular ion peak at  $m/z$  639  $[M^+Na]$ . MS<sup>2</sup> yielded fragment ion peaks at  $m/z$  579  $[M^+H-2H_2O]$  and 411  $[M^+Na-C_{14}H_{27}O_2]$  which showed the loss of two molecules of water and one molecule of tetradecanoic acid respectively. Further fragmentation gave  $[M^+Na-C_{14}H_{27}O_2-CH_3COOH]$  at  $m/z$  351 showing the loss of one molecule of acetic acid and tetradecanoic acid. The base peak fragment at  $m/z$  293  $[M^+H-acetic\ acid\ and\ tetradecanoic\ acid-2H_2O]$  was observed. Therefore, peak 38 was tentatively identified as 12-*O*-tetradecanoylphorbol-13-acetate (TPA) and its isomer [13].

#### Deoxyphorbol esters:

Six deoxyphorbol esters were characterized in both Euphorbia species (three in *E. paralias* and five in *E. geniculata*). Peaks 15, 43, 44 and 48 with time retention 13.03, 25.61, 26.34 and 27.73 min, respectively, gave the same molecular ion peaks at  $m/z$  531  $[M^+Na]$ . MS<sup>2</sup> fragment ions produced at  $m/z$  489  $[M^+Na-CH_3CO]$ , 471  $[M^+Na-CH_3COOH]$ , 335  $[M^+Na-CH_3COOH-C_8H_7O_2]$  showed the loss of acetyl, acetic acid and acetic acid with phenylacetyl moieties respectively. The fragment at  $m/z$  295  $[M^+H-CH_3COOH-C_8H_7O_2-H_2O]$  was observed. Accordingly peak 15 was tentatively characterized as 13-*O*-phenylacetyl-12-deoxyphorbol-20-acetate and its isomers [13]. While peak 31 (Rt, 19.61 min.) exhibited a molecular ion peak at  $m/z$  483  $[M^+Na]$  and MS<sup>2</sup> fragment at  $m/z$  335 which showed the loss of acetic and isobutyric acid moieties. It gave a fragment ion at  $m/z$  317 owing to the subsequent loss of one molecule of water. The base peak fragment ion at  $m/z$  295  $[M^+H-acetic\ acid- isobutyric\ acid-H_2O]$  was observed. These data confirmed the existence of butyryl and acetyl moieties. Therefore, peak 31 was tentatively identified as 13-*O*-isobutyryl-12-deoxyphorbol-20-


acetate [13]. Similarly, peak 37, (Rt, 21.31 min), its ESI-MS spectrum showed  $m/z$  535 [ $M^+ + Na$ ], 551 [ $M^+ + K$ ] and 513 [ $M^+ + H$ ]. The dominant characteristic fragment ions at  $m/z$  495 [ $M^+ + H - H_2O$ ] showing the loss of one molecule of water,  $m/z$  395 [ $M^+ + H - H_2O - C_4H_7O_2$ ] exhibited the loss of angelic acid and water  $m/z$  295 [ $M^+ + H - H_2O - 2C_4H_7COOH$ ] showing loss of two angelic acid moieties and one molecule of water, the loss of C=O moiety was observed from fragment at  $m/z$  267 [ $M^+ + H - H_2O - 2C_4H_7COOH - CO$ ]. Thus peak 37 was tentatively identified as 12-deoxyphorbol-13, 20 diangelate (Fig. 7) [13].

### 20-oxo-phorbol esters

One 20-oxo-phorbol ester was tentatively identified in *E. paralias* and three in *E. geniculata* as follow:

Peak 36 (Rt, 20.01 min) showed a molecular ion peak at  $m/z$  425 [ $M^+ + Na$ ]. The ESI-MS/MS product

ion at  $m/z$  331 [ $M^+ + H - C_3H_5O_2$ ] is produced by cleavage of  $C_3H_5O_2$  of propionic acid moiety. The base peak fragment at  $m/z$  291 showed the subsequent loss of 40 amu [ $Na + H_2O$ ] from 331 which showed the loss of one molecule of water. This fragment is characteristic for 20-oxophorbol skeleton [13]. These data led to tentative identification of peak 36 as 20-oxo-13-deoxy phorbol-12-propionate. By the same manner peaks 39, 40 and 41 (Rt, 24.28, 24.78, 24.98 min.) which gave their molecular ion peak at  $m/z$  525 [ $M^+ + Na$ ] were tentatively identified as 20-oxophorbol-12, 13-dibutyrate isomers (Fig. 8, Scheme 5) [13]. The MS<sup>2</sup> fragment ions at  $m/z$  507 [ $M^+ + Na - H_2O$ ] showed the loss of one molecule of water, In addition to  $m/z$  437 [ $M^+ + Na - C_4H_8O_2$ ] showed the loss of butyric acid, 331 [ $M^+ + Na - 2C_4H_8O_2 - H_2O$ ] exhibit the loss of two molecule of butyric acid and one molecule of water.



Compound No.	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>
19, 20, 25, 26	O-Tetradecanoate	OH	OH
2, 14, 16, 46	O- decanoate	OH	OH
3	OH	OH	OH
4, 17	O-acetate	OH	OH
5, 22	OH	O-acetate	OH
6, 7, 45, 47, 49	O-acetate	O-acetate	OH
9-11	N-methyl anthranilate	O-acetate	OH
21, 24, 51	O- decanoate	O- decanoate	OH
13	O-Hexanoate	O-Hexanoate	OH
38, 42	O-Tetradecanoate	O-acetate	OH
15, 43, 44, 48	H	O-Phenylacetyl	O-Acetate
31	H	O-Isobutyryl	O-Acetate
36	O-Propionyl	H	O
37	H	O-Angelate	O-Angelate
39, 40, 41	O- Butyrate	O- Butyrate	O

Figure 4: The chemical structures of phorbol esters identified in methylene chloride fractions of *E. paralias* and *Egeniculata*







Figure 5: Positive ESI-MS/MS spectrum of phorpol-12-acetate (peak 4  $m/z$  429 $[M^++Na]$ )



Scheme 2: Characteristic ESI-MS/MS fragmentation pattern of phorpol-12-acetate in positive ion mode





Figure 6: Positive ESI-MS/MS spectrum of 12-*O*-(*N*-methylantranilate) phorbol-13-acetate (peak 9  $m/z$  562 $[M^++Na]$ )



Scheme 3: Characteristic ESI-MS/MS fragmentation pattern of 12-O-(N-methylantranilate) phorbol-13-acetate in positive ion mode





Figure 7: Positive ESI-MS/MS spectrum of 12-deoxyphorbol-13, 20 diangelate (peak 37  $m/z$  535  $[M^+ + Na]$ )



Scheme 4: Characteristic ESI-MS/MS fragmentation pattern of 12-deoxyphorbol-13, 20-diangelate in positive ion mode





Figure 8: Positive ESI-MS/MS spectrum of 20-oxo-phorbol-12, 13-dibutyrate (peak 39  $m/z$  525 $[M^+Na]$ )



Scheme 5: Characteristic ESI-MS/MS fragmentation pattern of 20-oxo-phorbol-12, 13-dibutyrate in positive ion mode

### Lathyrol and hydroxylathyrol esters

Only one lathyrol (Euphorbia Factor L<sub>3</sub>) and two hydroxylathyrol (5-pentanoyl-3-benzoyl-hydroxylathyrol and its isomer) were determined in *E. paralias* while it is completely absent in *E. geniculata*.

Peaks 8 and 12 (Rt, 11.69 and 12.14 min) have the same molecular ion peaks at 567  $[M^+K]$ , 561  $[M^+Na]$ ,  $m/z$  539  $[M^+H]$ . The MS<sup>2</sup> diagnostic fragment ion appear at  $m/z$  461 for  $[M^+Na-C_5H_9O_2]$  which showed the loss of pentanoic acid and the fragment ion at  $m/z$  317  $[M^+H-C_7H_5O_2-C_7H_5O_2]$  showed the loss of pentanoic and benzoic acid moieties as shown in (Fig 9). The fragment ions at  $m/z$  299 and 281 showed the loss of two molecules of water which are diagnostic fragments corresponding to the

residual lathyrene skeleton [14]. From these results and through comparison with previous reports it was concluded that peaks 8 and 12 were tentatively identified as 5-pentanoyl-3-benzoyl-hydroxylathyrol and its isomer [12]. Similarly, peak 32 (Rt, 19.94 min) was identified as Euphorbia Factor L<sub>3</sub> which has a molecular ion peak at  $m/z$  561  $[M^+K]$ , 545  $[M^+Na]$ , 523  $[M^+H]$  (Fig. 10, Scheme 6). It displayed MS<sup>2</sup> fragment ions at  $m/z$  463  $[M^+H-CH_3COOH]$ , 403  $[M^+H-2CH_3COOH]$ , 375  $[403-CO]$ , 341, 281(100%)  $[M^+H-2CH_3COOH-C_6H_5COOH]$  due to the loss of two molecules of acetic acid and one benzoic acid moieties. These data are in a good agreement with the reported data [12].




Figure 9: The chemical structures of lathyrene diterpene esters identified in methylene chloride fractions of *E. paralias* and *E. geniculata*





Figure 10: Positive ESI-MS/MS spectrum of Euphorbia factor L<sub>3</sub> (peak 32  $m/z$  545 [M<sup>+</sup>+Na])



Scheme 6: Characteristic ESI-MS/MS fragmentation pattern of Euphorbia Factor L<sub>3</sub> in positive ion mode

**Jatrophone diterpene:**

Only one Jatrophone diterpene was determined in *E. paralias*. Compound 18 ( $R_t$  of 14.16 min) (Table 1) was identified as 3,5,7,8,15, pentacetyl-2-hydroxy-9,14-dioxo-jatropha-6(16)-11(12)-diene with molecular ion peak at  $m/z$  647 [ $M^+ + K$ ], 631 [ $M^+ + Na$ ] and MS<sup>2</sup> fragmentation displayed several fragment ions for loss of several acetic acid moieties at  $m/z$  591 [ $M^+ + H - H_2O$ ], 549 [ $M^+ + H - H_2O - COCH_3$ ], 531 [ $M^+ + H - 2H_2O - CH_3COOH$ ], 489 [ $549 - 60(CH_3COOH)$ ] 471 [ $M^+ + H - H_2O - 2CH_3COOH$ ], 429 [489-60

( $CH_3COOH$ )] and 411 [ $M^+ + H - H_2O - 3CH_3COOH$ ]. All these fragment ions showed the probability of loss of several acetyl groups as acetic acid (60 amu) or as acetyl group (42 amu). Furthermore, fragmentation showed fragment ions at  $m/z$  387, 369, 341, 327, 309, 299 and 281 corresponding to the residual jatrophone nucleus which are in a good agreement with the previously reported data of 3, 5, 7, 8, 15-pentacetyl-2-hydroxy-9,14 dioxo-jatropha-6(16)-11(12)-diene (Fig. 11 & 12 (A)) [13].




Figure 11: Positive ESI-MS/MS spectrum of 3, 5, 7, 8, 15- pentacetyl-2-hydroxy- 9, 14 dioxo-jatropha-6(16)-11(12)-diene (peak 18  $m/z$  631[ $M^+ + Na$ ])

**Daphnanes diterpenes:**

Resiniferatoxin was the only daphnanes diterpene identified in *E. geniculata*. This compound is reported to have a prolonged effect on relief of spasm and pain and more potent than capsaicin with reduced side effects at nontoxic dose and may therefore be a better therapeutic option [15]. Compound 50 ( $R_t$ . 28.30 min) gave [ $M^+ + Na$ ] at  $m/z$  651 in the MS/MS spectrum (Fig 12 (B)) yields the diagnostic transition corresponding to

the loss of  $C_7H_7O_2$  moiety at  $m/z$  507 [ $M^+ + H - C_7H_7O_2$ ], in addition to fragment ion at  $m/z$  371 [ $M^+ + H - C_9H_9O_4 - C_6H_5$ ] and  $m/z$  355 [ $M^+ + Na - C_9H_9O_4 - C_7H_7$ ] which showed loss of  $C_9H_9O_4$  for 4-hydroxy-3-methoxyphenyl acetic acid and  $C_7H_7$  for benzyl moiety, respectively. Also MS<sup>2</sup> fragment at  $m/z$  333 [ $M^+ + H - C_9H_9O_4 - C_7H_7$ ], comparing these results with the reported literature [13] peak 50 was tentatively identified as resiniferatoxin (Fig. 13).




Figure 12: The chemical structures of jatrophone (A) and daphnane (B) diterpenes identified in methylene chloride fractions of *E. paralias* and *Egeniculata*




Figure 13: Positive ESI-MS/MS spectrum of resiniferatoxin (peak 50,  $m/z$  651  $[M+Na]$ )

### Discussion:

According to the characteristic fragmentation pathways of the detected diterpenoids in comparison with literature, totally 52 compounds were preliminary identified for the first time in the methylene chloride fractions of *E. paralias* (30 compounds) and *Egeniculata* (36 compounds). It is noteworthy to say that lathyrane diterpenes and jatrophane diterpene were determined only in *E. paralias* and were absent in *Egeniculata* while daphnane diterpene was determined only in *Egeniculata*. Moreover, the tiglliane and ingenane diterpenoids were determined in both *Euphorbia* species. The compounds determined in both plants were reported to have powerful anti-chikungunya virus (anti-CHIKV) activity [2]. In particular compounds 2, 6, 9, 14, 16, 21, 23, 24, 26, 31, 38 and 42 proved to be the most promising anti-CHIKV agents till now. Chikungunya virus is spread to people by bite of infected mosquito this cause fever, joint pain and swelling, headache, muscle pain, rash and there is no vaccine to prevent this viral infection. Thus, *E. paralias* and *E. geniculata* offer interesting opportunity as potential therapeutic agents for anti-chikungunya virus.

### References:

- Hammadi R., Kusz N., Mwangi P. W., Kulmany A., Zupko I., Orvos P. et al. Isolation and pharmacological investigation of compounds from *Euphorbia matabelensis*. *Nat. prod. Com.* 2019, 1-5.
- Remy S. and Litaudon M., Macrocyclic Diterpenoids from Euphorbiaceae as A Source of Potent and Selective Inhibitors of Chikungunya Virus Replication. *Mol. J.* 2019, 24: 2336-2351.
- Shu X., Yu L., Tang Y., Zhang L., Ding A., Luo D. Bioassay-guided separation of the pro-inflammatory constituents from the roots of *Euphorbia kansui*. *J. Nat. Med.* 2010; 64:98-103.
- Yang M., Lu Z., Yu K., Yang Q., Chen X., Li Y. et al. Studies on the fragmentation pathways of ingenol esters isolated from *Euphorbia esula* using IT-MS and Q-TOF-MS/MS methods in electrospray ionization mode. *Int. J. Mass Spec.* 2012, 323-324, 55-62.
- Miyata, S., Wang L.Y., Yoshida C., and Kitanaka S. Inhibition of cellular proliferation by diterpenes, topoisomerase II inhibitor. *Bioorg. & Med. Chem.* 2006, 14: 2048-2051.
- Yoshida C., Hishiyama K., Miyazaki K., Watanabe M., Kanbe M., Yamada Y. et al. Analysis of inhibition of topoisomerase II alpha and cancer cell proliferation by ingenol EZ. *Can. Sci.* 2010, 101: 374-378.
- Jiao W., Dong W.; Li, Z., Deng M., Lu R. Lathyrane diterpenes from *Euphorbia lathyris* as modulators of multidrug resistance and their crystal structures. *Bioorg. Med. Chem.* 2009, 17: 4786-4792.
- Ayatollahi A.M., Ghanadian M., Mesaik A., Abdella O.M., Afsharypuor S., Kobarfard F. and Mirza-taheri M., New myrsinane-type diterpenoids from *Euphorbia aellenii* Rech. f. with their immunomodulatory activity. *Journal of Asian Natural Products Research*, 2010, 1020-1025.
- Avila L., Perez M., Sanchez-Duffhues G., Hernández-Galán R., Munoz E., Cabezas F. et al. Effects of diterpenes from latex of *Euphorbia lactea* and *Euphorbia laurifolia* on human immunodeficiency virus type 1 reactivation. *Phytochem.* 2010, 71: 243-248.
- Vasas A. and Hohmann J., *Euphorbia* Diterpenes: Isolation, Structure, Biological Activity, and Synthesis. (2008-2012). *Chem. Rev.* 2014, 114: 8579-8612.
- Baloch I. B., Baloch M. K. and Baloch A. K. Bioactive compounds from *Euphorbia cornigera*. *Boiss. Eur. J. Med. Chem.* 2009, 44: 3188-3194.

12. Xu, J.; Jin, D.; Guo, P.; Xie, C.; Fang, L.; Guo, Y. Three New Myrsinol Diterpenes from *Euphorbia prolifera* and Their Neuroprotective Activities. *Mol.* 2012, 17: 9520-5528.
13. Nothias-Scaglia L-F, Schmitz-Afonso I., Renucci F., Roussi F., Touboul D., Costa J. *et al.* Insights on Profiling of Phorbol, Deoxyphorbol, Ingenol and Jatrophone Diterpene Esters by High Performance Liquid Chromatography Coupled to Multiple Stage Mass Spectrometry. *J. Chromat. A.* 2015, 128-139.
14. Bicchi C., Appendino G., Cordero C., Rubiolo P., Ortelli D., and Veuthey J, UPLC-UV and HPLC-Positive –ESI-MS Analysis of the diterpenoid fraction from Caper Spurge (*Euphorbia Lathyris*) seed oil. *Phytochem. Anal.* 2001, 12: 255-262.
15. Sharma P.K., Avula B., Panda A., Wang Y-Hong, Khan IA, Murthy S. N., A Sensitive Bioanalytical Method for Quantitative Determination of Resiniferatoxin in Rat Plasma Using Ultra-High Performance Liquid Chromatography Coupled to Tandem Mass Spectrometry and its Application in Pharmacokinetic Study, *J. Pharm. and Biom. Anal.* 2019, 165, 284-291.