

## Physical and Psychological Problems among Patients Post Bariatric Surgery

Asmaa Atif Abd El-shaheed,<sup>(1)</sup> Afaf Mohamed,<sup>(1)</sup> Fatma Ata,<sup>(2)</sup> Amr Hamed

<sup>(1)</sup>Psychiatric-Mental Health Nursing, Faculty of Nursing, Ain Shams University

<sup>(2)</sup>General Surgery Faculty of Medicine-Ain Shams University

### Abstract

**Background:** Bariatric surgery is currently considered the most effective treatment option for morbid obesity; it results in greater improvements in weight loss outcomes and obesity related co-morbidities but there are several complications that can arise from weeks to years after bariatric surgery. **Aim:** this study aimed to assess physical and psychological problems among patients post bariatric surgery. **Design:** A descriptive exploratory design was conducted in this study. **Setting:** This study was conducted at surgical unit and obesity clinic in Eldemerdash Hospital. **Sample:** Convenient sample of 177 patients was obtained from surgical unit and obesity clinic in Eldemerdash Hospital affiliated to Ain Shams University Hospitals. **Tools of data collection:** Physical and psychological problem questionnaire consisted of Beck depression inventory, body image acceptance and action questionnaire, Rosenberg self-esteem scale and eating disorder questionnaire. **Results:** Physical problems less than three quarter of the studied patients had dumping syndrome, majority of the patients had nutritional deficiencies and steatorrhea and two thirds of them complained from sagging skin. Psychological problems, it was found that, depressive symptoms appeared on two fifth of the studied patients, less than three quarter of the studied patients not accept their body image, less than half of them had low self-esteem and less than three quarter of them weren't restraining their eating. **Conclusions:** Majority of the studied patients had nutritional complications and steatorrhea, two thirds of the studied patients complained from excess skin, less than three quarter of the studied patients not accept their body image, less than three quarter of them weren't restraining their eating. **Recommendations:** Further study is recommended about lifestyle modification and cognitive behavioral therapy to patients post bariatric surgery. Further study is recommended to evaluate psychological assessment before and after the surgery.

**Key words:** Bariatric Surgery, Physical Problems & Psychological Problems.

### Introduction

One of the most common problems related to lifestyle today is being overweight. Severe overweight or obesity is a key risk factor in the development of many chronic diseases. Obesity is often defined simply as a condition of abnormal or excessive fat accumulation in the fat tissues of the body leading to health hazards (World Health Organization, 2015).

The treatment of obesity includes primary prevention, dietary measures, behavior modification, pharmacotherapy and bariatric surgery (BS). Bariatric describes the medical treatment of serious overweight that is, obesity. Bariatric surgery is only employed when other methods of weight loss have been tried and failed (Mandal, 2014).

Bariatric surgery techniques are applied in various types. There are four main types of bariatric surgery: the Roux-

en-Y, adjustable gastric band, sleeve gastrectomy, and biliopancreatic diversion with duodenal switch (**Dambaugh & Ecklund, 2016**).

As with all medical procedures, postoperative complications will occur. These problems present both early and late in the postoperative course. Bariatric operations, in many instances, result in permanent alteration of a patient's anatomy, which can lead to complications at any time during the course of a patient's life as dumping syndrome, stenosis, nutritional deficiency and sagging skin (**Lim, Beekley, Johnson & Davis, 2018**).

While bariatric surgery has been shown to improve a number of metabolic conditions, some patients experience persisting disorder in psychological outcomes like depression, body image, self-esteem and eating disorders (**Jumbe, Hamlet & Meyrick, 2017**).

#### **Significance of the study:**

Bariatric surgery is designed to alter or interrupt the digestion process. A reduction in the amount of nutrients and calories absorbed enables patients to lose weight and decrease their risk for obesity-related health risks or disorders. It's estimated that 5875 of population in Egypt done bariatric surgery (**Angrisani, Santonicola, Iovino, Formisano, Buchwald & Scopinaro, (2015)**). According to **Ain shams university hospital statistical department (2015)**, 500 patients are undergone bariatric surgery in this year. In order to have a positive long-term outcome, bariatric surgery patients must be assessed for physical and psychological problems from a relapse prevention standpoint.

#### **Aim of the study:**

The present study was conducted to assess physical and psychological problems among patients post bariatric surgery.

#### **Research Question:**

What are physical and psychological problems among patient post bariatric surgery?

#### **Subjects and Methods**

##### **A-Research design:**

A descriptive exploratory design was conducted to achieve the aim of this study.

##### **B- Research Setting:**

This study was conducted in surgical units and obesity clinic at Ain Shams University Hospital affiliated to Ain Shams University.

##### **Research Subject:**

Convenient sample of 177 patients was obtained from surgical unit (77patients) and obesity clinic (100 patients) in Eldemerdash Hospital affiliated to Ain Shams University Hospitals.

##### **D-Tools for data collection:**

The study data was collected through the following tools:

##### **• Part (1): Socio-demographic sheet:**

The researcher developed this sheet to assess bariatric surgery patient's socio-demographic characteristics. It contains the following items: Age, gender, educational level, marital status, previous bariatric surgery, type of bariatric surgery and medical history.

**Part (2):** Physical problem questionnaire:

The researcher designed this tool after a thorough review of relevant literature of medical complications to assess physical problems post bariatric surgery. It was translated into Arabic by researcher and was revised by jury of seven expertise. It contained 18 items covering the main physical problems such as dumping syndrome, wound infections, stenosis, hemorrhage anastomotic leak, deep venous thrombosis and pulmonary embolism.

#### **Scoring system**

Each item had set of answers; chosen answer was given one mark. The total score equals 60. Low physical problems <30. Moderate physical problems [30-45]. High physical problems [46-60].

**Part (3):** Psychological problem questionnaire consisted of the following:

The Beck Depression Inventory-II (BDI-II): Created by Aaron Beck and was modified by researcher. It contains 16 items to assess relating symptoms of depression such as sadness, pessimism, past failure, loss of pleasure, guilty feeling, and punishment feeling and self-dislike.

#### **Scoring system:**

Each item has set of responses ranging in intensity. First response is 1, second response 2, third response 3, fourth response 4. Total score equals 50.

Mild depression [16 <24], Moderate depression [24-38] & Severe depression [>38-50].

Higher total scores indicate more severe depressive symptoms

Body image-Acceptance and action questionnaire:

Created By Emily, Sandoz & Kelly in 2006. It is modified by researcher. It contains 14 items on 5 points likert scale to assess body image concerns, acceptance and dissatisfaction such as I get on with my life even when I feel bad about my body, Worrying about my weight makes it difficult for me to live a life that I value, I care too much about my weight and body shape, I care too much about my weight and body shape and How I feel about my body has very little to do with the daily choices I make.

#### **Scoring system:**

The responses for each item were never, seldom, sometimes, frequently and always. Scores for each response ranged from 1 to 5 as the following: Never =1, seldom= 2, sometimes= 3, frequently= 4 and always= 5. Total score equals 70. Higher scores indicate more acceptances.

#### **Rosenberg self-esteem scale:**

Created by Morris Rosenberg in 1965, to assess level of self-esteem. It is a ten-item Likert-type scale with items answered on a four-point scale from strongly agrees to strongly disagree. Five of the items have positively worded statements and five have negatively worded ones. The scale measures state of self-esteem by asking the respondents to reflect on their current feeling such as on the whole, I am satisfied with myself, at times I think I am no good at all. I feel that I have a number of good qualities and I am able to do things as well as most other people.

**Scoring system:**

For items 1, 2, 4, 6, and 7: Strongly agree = 3, Agree = 2, Disagree = 1 & strongly disagree = 0.

For items 3, 5, 8, 9, and 10 (which are reversed in valence):

Strongly agree = 0, Agree = 1, Disagree = 2, strongly disagree = 3. Total score equals 30.

**Eating disorders questionnaire:**

The researcher designed this questionnaire after a thorough review of relevant literature to assess eating disorders post bariatric surgery. It was translated into Arabic by researcher. It contained 15 items covering the main items of eating disorders such as do you often feel hungry, do you eat any type of food, do you pay attention to type of food you eat, do you eat healthy food quantities, do you eat more frequent meal, do you count your food calories and do you eat and chew slowly.

**Scoring system:**

The answer for each item either yes or no, scoring as following: yes=1, no=0, the total score equals 15. Restrain [0-8], Not Restrain [9-15].

**Operational design:****Operational Design:**

The Operational design of the study includes preparatory phase, pilot study, and field work, limitations of the study and ethical considerations.

**The Preparatory Phase:**

1. Tools of data collection development. First, tools of data collection were developed and adopted by the

researcher based on the relevant literature of review.

**Tools validity and reliability:**

To achieve the criteria of trustworthiness of the tools of data collection in this study, the tools were tested for its face and content validity by jury group consisting of seven experts of psychiatric mental health nursing from Faculty of Nursing, Ain Shams University who reviewed the content of the tools for its comprehensiveness, accuracy, clarity and relevance. The required corrections and modifications were done. The items on which 85% or more of the experts had agreed were included in the proposed tool.

**Testing reliability** of proposed tools was assessed statistically through measuring their internal consistency by determining Cronbach's Alpha.

Tool	Reliability		
	Reliability Coefficient	Cronbach's Alpha	Face validity
Physical problems questionnaire	0.97	.98	95%
Beck depression inventory.	.70	0.78	96%
The Body Image - Acceptance and Action Questionnaire	0.95	0.98	95%
Eating disorder questionnaire	0.91	0.61	98%

**Pilot study (Exploratory phase):**

A pilot study was conducted to test clarity, applicability, efficiency and validity of the study tools and to estimate the required time to gather data. The pilot study was conducted over a period of one week starting from 22 of June 2018 to 29 of June 2018 on 10% (17 patients) of the expected

study subjects, all participants in the pilot study were excluded later from the study sample. According to the pilot study results the necessary modifications were done some questions were restated in Arabic language on the study tool after pilot study.

### **Field Work:**

The researcher reviewed the recent, past, and local and international current related literature in the various aspects of medical psychiatric complications to be acquainted with in depth information in order to design the study tools. The actual field work for the process of the data collection has consumed six months started on beginning of July 2018 and was completed by the end of December 2018.

The researcher collect data from research setting three days/week Saturday, Sunday and Tuesday at morning (from 8 am-2 pm) and afternoon shift (from 2pm to 8 pm) from surgical ward and from 10am - 2pm from the obesity clinic. The researcher distributing the tools to the patients after explaining aim and objectives of the study to them. The patients filled in the sheet; it took about 20-30 minutes to complete it. The researcher collected the answered sheet from the patients.

### **Limitations of the study:**

During data collections there was renewal of the surgical ward and patients were distributed to other sections this made difficulty to find the target patients.

### **Ethical Considerations:**

In order to conduct the study, the researcher obtained official permission from the following authorities:

### **Faculty of Nursing Ain Shams University:**

The chairperson and council members of psychiatric/mental health nursing department, ethical approval was obtained from the scientific ethical committee of Faculty of Nursing - Ain Shams University before starting the study. Official letter from the dean of the faculty to the director of Ain Shams University Hospitals.

### **After securing official requirement for conducting the study:**

The researcher clarified the objective and aim of the study to the subjects of the study. They will be assured that anonymity and confidentiality would be guaranteed and will be used for scientific purpose only. They had the right to participate or withdraw from the study at any time without giving reasons.

### **Administrative Design:**

Official letter including the aim of the study was sent from the Dean of faculty of Nursing Ain Shams University to the director of Ain Shams University Hospitals seeking his approval to carry out the study. The official permission to conduct the study was obtained from director of Ain Shams University Hospitals. Oral consent obtained from every participant to conduct the study.

### **Statistical design:**

Recorded data were analyzed using the statistical package for social sciences, version 20.0 (SPSS Inc., Chicago, Illinois, USA). Quantitative data were expressed as mean  $\pm$  standard deviation (SD). Qualitative data were expressed as frequency and percentage.

## Result

**Table (1):** Number and percentage distribution of patient post bariatric surgery according to their socio-demographic data (N=160).

Socio-Demographic Data	No.	%
<b>Age (years)</b>		
16 > 29 years	32	20.0%
29 >40 years	70	43.7%
40 > 50 years	36	22.5%
50 >55 years	22	13.8%
Mean±SD	38.44±4.23	
<b>Gender</b>		
Male	42	26.3%
Female	118	73.7%
<b>Educational level</b>		
Illiterate	52	32.5%
Primary	14	8.7%
Secondary	34	21.3%
University	46	28.7%
Post graduate	14	8.8%
<b>Marital status</b>		
Single	66	41.3%
Married	88	55.0%
Divorced	6	3.7%
<b>Previous bariatric surgery</b>		
Yes	54	33.7%
No	106	66.3%
<b>type of bariatric surgery</b>		
Adjustable gastric band	0	0.0%
Sleeve Gastrostomy	84	52.5%
Gastric bypass	70	43.7%
Biliopancreatic Diversion with Duodenal Switch	6	3.8%
<b>Medical history:</b>		
Diabetes millets( DM)	50	31.3%
Hypertension (HTN)	40	25.0%
Morbid obesity	90	56.3%

**Table (1):** Showed that, 43.7% of the studied patients were in the age group 29 >40 years with a mean age of 38.44±4.23 and 73.7% of them were females. Regarding level of education, 32.5% of the studied patients were illiterate and 55.0% of those patients were married. Moreover, 66.3% of the studied patients never had previous bariatric surgery. As regard to type of bariatric surgery; 52.5% of the studied patients were performed sleeve gastrostomy and 56.3% of patient's medical history were obese.


**Table (2):** Number and percentage distribution of patient post bariatric surgery according to their physical problems (N=160).

Physical problems post bariatric surgery	Yes		No	
	No.	%	No.	%
<b>1. Dumping syndrome:</b>				
Nausea	94	58.7%	66	41.3%
Vomiting	102	63.7%	58	36.3%
Diarrhea	62	38.7%	98	61.3%
Weakness	58	36.2%	102	63.8%
No symptoms	44	27.5%	116	72.5%
<b>2. Vit B12 deficiency:</b>				
Depression	88	55.0%	72	45.0%
Neuropathy	16	10.0%	144	90.0%
Pernicious anemia	4	2.5%	156	97.5%
No symptoms	66	41.3%	94	58.7%
<b>3. Iron deficiency:</b>				
General malaise	118	73.7%	42	26.3%
Anemia	104	65.0%	56	35.0%
Pale skin and nails	88	55.0%	72	45.0%
Dizziness/lightheadedness	128	80.0%	32	20.0%
No symptoms	20	12.5%	140	87.5%
<b>4. Vit A deficiency:</b>				
Decreased vision	18	11.3%	142	88.7%
Night vision difficulty	65	40.6%	95	59.4%
Itching	8	5.0%	152	95.0%
Dry hair	114	71.2%	46	28.8%
No symptoms	34	21.2%	126	78.8%
<b>5. Vit D deficiency:</b>				
Osteopenia	4	2.5%	156	97.5%
Osteoporosis	8	5.0%	152	95.0%
Bone aches	106	66.3%	54	33.7%
No symptoms	42	26.2%	118	73.8%
<b>6. Zinc deficiency:</b>				
Nail dystrophy	14	8.8%	146	91.2%
Skin eruptions	18	11.3%	142	88.7%
Poor wound healing	66	41.3%	94	58.7%
Hair loss	116	72.5%	44	27.5%
No symptoms	44	27.5%	116	72.5%
<b>7. Copper deficiency:</b>				
Anemia	60	37.5%	100	62.5%
Unexplained bleeding under skin	4	2.5%	156	97.5%
Fatigue	4	2.5%	156	97.5%
Difficulty learning and remembering	56	35.0%	104	65.0%
No symptoms	60	37.5%	100	62.5%
<b>8. Folate deficiency:</b>				
Sore tongue and redness	44	27.5%	116	72.5%
General weakness	112	70.0%	48	30.0%
Anemia	104	65.0%	56	35.0%
No symptoms	48	30.0%	112	70.0%
<b>9. Steatorrhea.</b>	144	90.0%	16	10.0%
<b>10. Sagging skin.</b>	100	62.5%	60	37.5%

**Table (2):** Represented that, 72.5% of the studied patients had symptoms of dumping syndrome. Regarding vitamin B12 deficiency, 55.0% of the studied patients had symptoms of depression, 87.5% of them had symptoms of iron deficiency, 78.8% of them had symptoms of vitamin A deficiency, 73.8% of them had symptoms of vitamin D deficiency, 72.5% of them had symptoms of zinc deficiency, 62.5% of them had symptoms of copper deficiency and 70.0% of them had symptoms


of folate deficiency. Meanwhile, 90.0% of the studied patients had steatorrhea and 62.5% of them complained from sagging skin.

**Figure (1):** Percentage distribution of patient post bariatric surgery according to total level of depression.



**Figure (1):** displayed that, two fifth of the studied patients (40.0%) had moderate depression.


**Figure (2):** Percentage distribution of patient post bariatric surgery according to total degree of body images acceptance.



**Figure (2):** clarified that, less than three quarter of the studied patients (73.8%) not accept their body image.




**Figure (3):** Percentage distribution of patient post bariatric surgery according to total level of self-esteem.



**Figure (3):** evidenced that, less than half of the studied patients had low self-esteem.

**Figure (4):** Percentage distribution of patients post bariatric surgery according to total level of eating disorders.



**Figure (4):** illustrated that, less than three quarter of the studied patients (73.8%) weren't restraining in their eating.

---

**Discussion**

---

**Concerning the demographic characteristics of the studied patients:**

The results of the present study revealed that, less than half of the study patient's ages were 29 to less than 40 years. According to researcher point of view this may be due to selection criteria of pre-operative patients. This finding is consistent with a study done by **Contreras, Santander, Court & Bravo, (2013)** who stated that, patients younger than 45 years lose greater amount of excess BMI than older patients after bariatric surgery.

Related to gender, less than three quarter of studied patients were females. This may be as females are more interested in their shape and may be related to social issues as chance for marriage also increased obesity in females than males in Egypt 26.4% of men and 48.4% of women are obese as reported by **Elrazek, Elbanna & Bilasy (2014)**.

This result on the same line with the study done by **Hans Fuchs, Broderick, Harnsberger, Chang, Bryan, Sandler, Jacobsen & Horgan (2015)** who stated that, Women seem to be more aware of the problems obesity brings to health. They are much more willing to look at surgical weight loss earlier in life, whereas men tend to wait until they have more comorbidity (adverse health conditions). Regarding marital status, the present study clarified that, more than half of studied patient were married. This may be related to social factors as unsatisfactory relationship with partner. This finding is in congruence with a study done by **De zwaan & Mitchell (2012)** who assured that pre surgical patient report frequently unsatisfactory marital relationship and difficulties in sexual relation.

Related to Type of bariatric surgery, the present study clarified that, slightly more than half of the studied patients had performed Sleeve Gastrectomy. This finding may be due to that Sleeve gastrectomy has fewer complications than other procedure. The finding of the present study is the same line with the study done by **Angrisani, Santonicola, Iovino, Formisano, Buchwald & Scopinaro (2017)** who assured that, the most commonly performed procedure in the world was sleeve gastrectomy (SG) that reached 45.9%, followed by Roux-en-Y gastric bypass (RYGB) 39.6%, in Egypt 1500 patient has Roux-en-Y gastric bypass (RYGB) and 31 00 have sleeve gastrectomy (SG).

**Concerning assessment of the physical problems post bariatric surgery.**

The present study finding indicated that, less than three quarter of the studied patients complain of Dumping syndrome (DS). There is agreement with **Ahmed, Kornrich, Krasner, Braslow, Broggelwirth & Eckardt (2017)** who assured that, patients underwent SG complain of signs of DS. Of those patients 73.2% of LSG reported symptoms of early DS this compares with LRYGB 79.5% of patents reported early symptoms of DS.

Regarding to vitamin B12 deficiency, the present study clarified that, more than half of the studied patients complain from symptoms of B12 deficiency. This finding is on the same line with a study done by **Carvalho, Loscalzo, Freitas, Jordão, & Friano (2012)** who assured that, it was seen a reduction in vitamin B12 in 47,2% patients.

The current study revealed that, the majority of the studied patients had symptoms of iron deficiency. This finding

is contradicting with study done by **Ben-Porat, Elazary, Yuval, Wieder, Khalaileh, & Weiss (2015)** who stated that, one year post surgery, deficiencies of iron was 28% of patients. In addition, **Alexandrou, Armeni, Kouskouni, Tsoka, Diamantis & Lambrinouadaki (2014)** Concluded that, iron deficiency was 30% versus 36.4% in patients with RYGB compared with SG.

Concerning Vit A deficiency, the current study clarified that, prevalence of Vit A deficiency was slightly more than three quarter of the studied patients. This is contradicting with a study done by **Eckert, Perry, Sohn, Boden, Martin, Rush & Steele (2010)** who assured that A low vitamin A level was identified in 11.0% of RYBG patients.

Regarding Vit D deficiency the present study showed that, less than three quarter of the studied patients present with Vit D deficiency. This finding goes on the same line with a study done by **Pellitero, Martínez, Puig, Leis, Zavala, Granada, Pastor, Moreno, Tarascó, Balibrea & Puig-Domingo (2017)** who displayed that, Vitamin D deficiency is the most prevalent long-term nutritional deficiency after SG as vitamin D prevalence was 73% of the subjects in his study.

Concerning Zinc deficiency, the current study proposed that, slightly less than three quarter of studied patients had symptoms of Zinc deficiency. This is contradicting with **Belfiore, Cataldi, Minichini, Aiello, Trio, Rossetti & Guida (2015)** who reported that 36 % of patients developed zinc deficiency.

Based on the results of the current study, it was found that, Folate deficiency was found in less than three quarter of the studied patients. Conversely, **Alexandrou et al. (2014)** concluded that At 4-year

follow-up more Folate deficiency frequently detected after SG than RYGB (20.0% versus 18.4%). The current study revealed that, the majority of the studied patients complain from steatorrhea. There is an agreement with **O'Keefe, Rakitt, Ou, El Hajj, Blaney, Vipperla, Holst & Rehlfeld (2017)** who concluded that, result showed that all patients had some fat mal absorption, 80%, indicative of steatorrhea.

The present study reported that, nearly two thirds of the studied patients complain from excess skin. This finding had an agreement with **Baillet, Brais-Dussault, Bastin, Cyr, Brunet, Aimé, Romain, Langlois, Bouchard, Tchernof, Lhoret, Garneau & Bernard (2017)** who concluded that, although bariatric surgery (BS) improves health among adults with severe obesity, it results in excess skin in more than 70% of adults.

#### **Concerning the third part: Assessment of psychological problems of the patients post bariatric surgery.**

Regarding depression, the current study indicated that, two fifth of the studied patients presented with moderate depressive symptoms. This finding had agreement from **Ivezaj, Barnes & Grilo. (2016)** who assumed that, based on a BDI cut-off score of >15, 41.9% (n=52) were categorized as High-BDI and 58.1% (n=72) as Low-BDI. In addition, **Faria, Faria, Cardeal, de Novaes, Marques & Feitosa (2016)** proposed that, depression Post-surgery was 58.8%.

Regarding body image the present study finding illustrated that, less than three quarter of the studied patients not accept their body image. This finding on the same line with a study done by **Sarwer, Dilks & West-Smith (2011)**, assured that, more than 66% of post-bariatric surgery patients reported body image dissatisfaction, this

dissatisfaction considered the development of excess skin to be a negative consequence of surgery and **Björserud, Shams, Elander & Olsen (2018)**, proposed that, self-image is lower among those who experience a high degree of discomfort of excess skin after bariatric surgery.

The present study demonstrated that, less than half of studied patients present with low self-esteem. This finding is contradicting with a study done by **Aldaql & Sehlo (2013)** who stated that, Self-esteem significantly improved 1 year after surgery. In addition, **Ghanbari, Lotfi, Pazouki, Mazaheri, Soheilipour & Jesmi (2016)** clarified that, no significant difference was found between self-esteem before and after the surgery ( $P = 0.321$ ). Conclusions Weight loss after bariatric surgery did not improve self-esteem.

The current study revealed that, less than three quarter of the studied patients are not restraining in their eating. This finding goes on the same line with **Sioka, Tzouvaras Oikonomou, Katsogridaki, Zachari Papamargaritis (2013)** explained that, eating disorder and emotional patterns had the worst Excessive Weight Loss ( $42.84 \pm 29.42$  and  $34.55 \pm 19.34$ , respectively). While **Kofman, Lent & Swencionis (2010)** reported that, (49.9%) of subject experiencing times when they could not stop eating or control how much they were eating. Also **Colles, Dixon & O'brien (2008)** noted that, BED, uncontrolled eating, and NES occurred after surgery in 3.1%, 22.5%, and 7.8%, Grazing was prevalent after surgery (38.0%).

### Conclusion

Physical problems less than three quarter of the studied patients had dumping

syndrome, majority of those patients had nutritional deficiency, majority of them had steatorrhea and more than half of them complained from excess skin.

Psychological problems, it was found that, depressive symptoms presented on two fifth of the studied patients, less than three quarter of them not accept their body image, less than half of them had low self-esteem and less than three quarter of them weren't restraining their eating.

### Recommendations

- Replication of the current study on large sample and different hospitals settings to be able to generalize the results.

- Further study is recommended to evaluate psychological assessment before and after the surgery.

- Further study is recommended about lifestyle modification and cognitive behavioral therapy to patients post bariatric surgery.

### References

- Ahmed, A., Kornrich, B.D., Krasner, H, Braslow. A., Broggelwirth, B., Eckardt, S. (2017):** Prevalence of Dumping Syndrome After Laparoscopic Sleeve Gastrectomy and Comparison with Laparoscopic Roux-En-Y Gastric Bypass. Surgery for obesity and related diseases, 13(10), s7-s30. DOI: <https://doi.org/10.1016/j.soard.2017.09.524> [https://www.soard.org/article/S1550-7289\(17\)30950-4/fulltext](https://www.soard.org/article/S1550-7289(17)30950-4/fulltext)
- Angrisani, L., Santonicola, A., Iovino, P., Formisano, G., Buchwald, H., and Scopinaro, N. (2015):** Bariatric surgery worldwide 2013. Obesity surgery, 25(10), 1822-1832.

- Aldaqa, S.M., Sehlo, M.G. (2013):** Self-esteem and quality of life in adolescents with extreme obesity in Saudi Arabia: the effect of weight loss after laparoscopic sleeve gastrectomy. *Gen Hosp Psychiatry*, 35(3), 259-64. DOI: 10.1016/j.genhosppsych.2012.12.011. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3662098/>
- Alexandrou, A., Armeni, E., Kouskouni, E., Tsoka, E., Diamantis, T., and Lambrinou, I. (2014):** Cross-sectional long-term micronutrient deficiencies after sleeve gastrectomy versus Roux-en-Y gastric bypass: A pilot study. *Surg Obes Relat Dis*, 10(2), 262-8. DOI: 10.1016/j.soard.2013.07.014 <https://www.ncbi.nlm.nih.gov/pubmed/24182446>
- Baillet, A., Brais-Dussault, E., Bastin, A., Cyr, C., Brunet, J., Aimé, A., Romain, J.A., Langlois, M., Bouchard, S., Tchernof, A., Lhoret, R., Garneau, P., Bernard, P. (2017):** What Is Known About the Correlates and Impact of Excess Skin After Bariatric Surgery: a Scoping Review. *Obesity Surgery*, 27(12). DOI: 10.1007/s11695-017-2814-3
- Ben-Porat, T., Elazary, R., Yuval, J.B., Wieder, A., Khalileh, A., and Weiss, R. (2015):** Nutritional deficiencies after sleeve gastrectomy: Can they be predicted preoperatively? *Surg Obes Relat Dis*, 11(5):1029-36. DOI: 10.1016/j.soard.2015.02.018. <https://www.ncbi.nlm.nih.gov/pubmed/25857443>
- Belfiore, A., Cataldi, M., Minichini, L., Aiello, M.L., Trio, R., Rossetti, G., and Guida, B. (2015):** Short-term changes in body composition and response to micronutrient supplementation after laparoscopic sleeve gastrectomy. *Obes Surg.*; 25(12): 2344-51. DOI: 10.1007/s11695-015-1700-0 <https://www.ncbi.nlm.nih.gov/pubmed/25948283>
- Biörserud, C., Shams, K., Elander, A., and Fagevik Olsén, M. (2018):** Skin and health-related quality of life. *J Plast Surg Hand Surg*, 1-6. DOI: 10.1080/2000656X.2018.1481860. <https://www.ncbi.nlm.nih.gov/pubmed/29957078>
- Carvalho, I.R., Loscalzo, I.T., Freitas, M.F., Jordão, R.E., and Friano Tde, C. (2012):** Incidence of vitamin B12 deficiency in patients submitted to Fobi-Capella Roux-en-Y bariatric surgery. *Arq Bras Cir Dig*, 25(1), 36-40. [https://www.ncbi.nlm.nih.gov/pubmed/?term=Friano%20Tde%20C%5BAuthor%5D&cauthor=true&cauthor\\_uid=22569977](https://www.ncbi.nlm.nih.gov/pubmed/?term=Friano%20Tde%20C%5BAuthor%5D&cauthor=true&cauthor_uid=22569977)
- Contreras, J.E., Santander, C., Court, and Bravo, J. (2013):** correlation between age and weight loss after bariatric surgery. *Obes Surg.* (8), 1286-9. DOI: 10.1007/s11695-013-0905-3
- Colles, S.L., Dixon, J.B., and O'Brien, P. E. (2008):** Grazing and loss of control related to eating: two high-risk factors following bariatric surgery. *Obesity*, 16(3), 615-622.
- Courcoulas, A.P., Yanovski, S.Z., Bonds, D., Eggerman, T.L., Horlick, M., Staten, M. A., & Arterburn, D.E. (2014):** Long-term outcomes of bariatric surgery: a National Institutes of Health symposium. *JAMA surgery*, 149(12), 1323-1329.
- Dambaugh, L. A., and Ecklund, M. M. (2016):** Progressive Care of Obese Patients, *Critical care nurse*, 36(4), 58-

63. DOI: 10.4037/ccn 2016510Mandal, A.
- De zwaan, M., Mitchell, E.J. (2012):** Psychosocial problems and psychiatric disorders, psychosocial assessment and treatment of bariatric surgery. Taylor and Francis group, New York. pp. 67.
- Eckert, M.J., Perry, J.T., Sohn, V.Y., Boden, J., Martin, M.J., Rush, R.M., and Steele, S.R. (2010):** Incidence of low vitamin A levels and ocular symptoms after Roux-en-Y gastric bypass. *Surg Obes Relat Dis*, 6(6), 653-7, doi: 10.1016/j.soard.2010.02.044. <https://www.ncbi.nlm.nih.gov/pubmed/20947440>
- Elrazek, A.E., Elbanna, A.E., & Bilasy, S.E. (2014):** Medical management of patients after bariatric surgery: Principles and guidelines. *World journal of gastrointestinal surgery*, 6(11), 220-228. doi: 10.4240/wjgs.v6.i11.220.
- Faria, S., Faria, O., Cardeal, M., de Novaes, M., Marques, J., and Feitosa, I. (2016):** Association between depression and nutritional deficiencies among Roux-en-Y Gastric Bypass patients. *Surgery for obesity and related disease*, 12(7), S175-S176. DOI: <https://doi.org/10.1016/j.soard.2016.08.313> [https://www.soard.org/article/S1550-7289\(16\)30492-0/fulltext](https://www.soard.org/article/S1550-7289(16)30492-0/fulltext)
- Ghanbari, J. A., Lotfi, T., Pazouki, A., Mazaheri, M. A., Soheilipour, F., & Jesmi F. (2016):** Comparison Between Marital Satisfaction and Self-Esteem Before and After Bariatric Surgery in Patients With Obesity. *Iran J Psychiatry Behav Sci*, 10(3), e2445. DOI: 10.17795/ijpbs-2445
- Hans F. Fuchs, H.F., Broderick, R.C., Harnsberger, C.R., Chang, C. D., Bryan J. Sandler, B.J., Jacobsen, G.R., & Horgan, S. (2015):** Benefits of Bariatric Surgery Do Not Reach Obese Men. *Journal of Laparoendoscopic & Advanced Surgical Techniques*, 25(3). <https://doi.org/10.1089/lap.2014.0639>
- Ivezaj, V., Barnes, R. D., & Grilo, C. M. (2016):** Validity and clinical utility of subtyping by the beck depression inventory in women seeking gastric bypass surgery. *Obesity surgery*, 26(9), 2068-2073. DOI: 10.1007/s11695-016-2047-x <https://europepmc.org/articles/pmc5129658>
- Jumbe, S. Hamlet, C. & Meyrick, J. (2017):** Psychological Aspects of Bariatric Surgery as a Treatment for Obesity, *current obesity reports* 6 (1), 71-78.
- Kinzl, J.F., Traweger, C., Trefalt, E., & Biebl, W. (2007):** Psychosocial consequences of weight loss following gastric banding for morbid obesity. *Obesity Surgery*, 13(1), 105-110. DOI: 10.1381/096089203321136683
- Kofman, M.D., Lent, M.R., & Swencionis, C. (2010):** Maladaptive Eating Patterns, Quality of Life, and Weight Outcomes Following Gastric Bypass: Results of an Internet Survey. *Obesity*, 18(10), 1938-2043. DOI: <https://doi.org/10.1038/oby.2010.27> <https://onlinelibrary.wiley.com/doi/full/10.1038/oby.2010.27>
- Lim, R., Beekley, A., Johnson, D.C., Davis, A.K. (2018):** Early and late complications of bariatric operation.

- Trauma Surgery & Acute Care Open, 3(1), e000219. DOI: 10.1136/tsaco-2018-000219.  
<https://www.ncbi.nlm.nih.gov/pubmed/30402562>
- Mandal, A. (2014):** What is Bariatric surgery?. Retrieved from <http://www.news-educal.net/health/What-is-Bariatric-Surgery.aspx>.
- O'Keefe, S.J.D., Rakitt, T., Ou, J., El Hajj, I.I., Blaney, E., Vippera, K., Holst, J.J. & Rehlfeld, J. (2017):** Pancreatic and Intestinal Function Post Roux-en-Y Gastric Bypass Surgery for Obesity. *Clin Transl Gastroenterol*, 8(8), e112. DOI: 10.1038/ctg.2017.39  
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5587840/>
- Pellitero, S., Martínez, E., Puig, R., Leis, A., Zavala, R., Granada, M.L., Pastor, C., Moreno, P., Tarascó, J., Balibrea, J., & Puig-Domingo, M. (2017):** Evaluation of vitamin and trace element requirements after sleeve gastrectomy at long term. *Obes Surg*, 27(7), 1674-1682. DOI: 10.1007/s11695-017-2557-1.  
<https://www.ncbi.nlm.nih.gov/pubmed/28161887>
- Sarwer D.B., Dilks R.J. & West-Smith L. (2011):** Dietary intake and eating behavior after bariatric surgery: threats to weight loss maintenance and strategies for success, surgery for obesity and related diseases, 7(5), 644-651. DOI: <https://doi.org/10.1016/j.soard.2011.06.016>
- Schauer P.R., Kashyap S.R., Wolski K, Brethauer S.A., Kirwan J.P., Pothier C.E., Thomas S., Abood B., Nissen S. E., Bhatt DL. (2012):** Bariatric surgery versus intensive medical therapy in obese patients with diabetes. *N Engl J Med*. 366: 1567-1576. DOI: 10.4240/wjgs.v6.i11.220  
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4241489/#>
- Sioka E, Tzouvaras G, Oikonomou K, Katsogridaki G, Zachari E, Papamargaritis D, et al. (2013):** Influence of eating profile on the outcome of laparoscopic sleeve gastrectomy. *Obesity Surgery*, 23(4), 501-8. DOI: 10.1007/s11695-012-0831-9
- World Health Organization (WHO). (2015):** Obesity. Retrieved from: <http://www.emro.who.int/health-topics/obesity/>.