

ANTIMICROBIAL ACTIVITY FOR EXCRETION AND SECRETION OF THE GREENBOTTLE FLY LARVAE *LUCILIA SERICATA* (MEIGEN) (DIPTERA: CALLIPHORIDAE)

By

**MOSTAFA I. HASSAN*, MOUNIR S. AMER, KOTB M. HAMMAD
AND MAHMOUD M. ZIDAN**

Department of Zoology and Entomology, Faculty of Science, Al-Azhar University,
Nasr City, Cairo, Egypt (*mostafa012@gmail.com)

Abstract

Sterile larval excretion/secretion (ES) exhibited antibacterial activity against some species of bacteria. They were shown to inhibit the growth of Gram-positive bacteria *Staphylococcus aureus* and *Bacillus subtilis* Gram-negative bacteria *Pseudomonas aeruginosa*, *Escherichia coli* and *Klebsiella pneumoniae* and Fungi *Geotricum candidum* and *Aspergillus fumigatus* thus exhibited limited inhibitory effect towards Gram-positive bacteria *Streptococcus pyogenes* and *Staphylococcus epidermidis* and Gram-negative *Proteus vulgaris* and Fungi *Syncephalastrum racemosum*, *Candida albicans*, that effect was slowed down when challenged with secretion on a solid media but no zone of complete inhibition was detected. Growth inhibiting activity was determined in liquid growth media using the Gram-positive, Gram-negative bacterial and fungal strains as indicator organisms.

Keywords: Antimicrobial activity, Antifungal activity, Maggot therapy, *Lucilia sericata*, Excretion/Secretion and Greenbottle fly larvae

Introduction

Effect of maggot, *Lucilia sericata* against biofilm of wound isolates *S. aureus* was carried out by Bohova *et al.* (2014), they suggested that maggot (ES) may act selectively against different bacterial strains. Most of excretion/secretion (ES) larvae of insects exhibit antibacterial activity against both Gram-positive, Gram-negative bacteria and fungi, especially *Lucilia sericata* (Valachova *et al.*, 2013).

The maggots (ES) of *L. sericata* larvae (maggots) have effective activity for debridement and disruption of bacterial such as *Staphylococcus epidermidis*, The excretions/secretions (ES) activity was characterized according to concentration, incubation time and temperature, thermal stability, and size (Harris *et al.*, 2013). Maggot's (ES) have a bactericidal and/or bacteriostatic activity against one or more of the bacterial species; the growth of Gram-positive bacteria is more reduced by maggots and/or their ES than the growth of Gram-negative bacteria (Cazander *et al.*, 2009).

The present study aimed to the evaluation of *L. sericata* excretion and secretion on some pathogenic organisms as antimicrobial agent.

Materials and Methods

Insect used: The 3rd larval instar of *L. sericata* was collected from Tonamel Village, Aga Center, Al-Dakahia Governorate, Egypt and maintained at Medical Entomology Laboratory, Animal House, Zoology Department, Faculty of Science, Al-Azhar University, under controlled laboratory conditions of 27±2°C and 70±10% RH and 12-12 light-dark photoperiod.

Collection of larval excretion/ secretion: Secretion and excretion of *L. sericata* maggots were collected as the following; washed sterile 3rd instar larvae (6000 larvae) with 70% ethanol and sterile ultrapure water (ddH₂O) and incubated overnight (10hrs) at 30°C, after incubation, centrifuged at 20,000×g for 15 min (kerridge *et al.*, 2005).

Antimicrobial bioassay: Twelve pathogenic organisms were used for the antimicrobial assay; *Staphylococcus aureus*, *Bacillus subtilis*, *Staphylococcus epidermidis* and *Strep-*

Staphylococcus aureus as Gram-positive bacterial strains, while the *Pseudomonas aeruginosa*, *Escherichia coli*, *Proteus vulgaris* and the *Klebsiella pneumoniae* were used as Gram-negative bacterial strains. The fungi which used in this assay were *Aspergillus fumigatus*, *Syncephalastrum racemosum*, *Candida albicans* and *Geotricum candidum*.

The antimicrobial assay employed was broadly based on the standard agar diffusion assay at Regional Center for Mycology and Biotechnology Antimicrobial unit test organism, Al-Azhar University, Cairo, Egypt. A colony of the tested microorganism was picked off a stock plate and suspended in Ringer's solution (10mL). An aliquot of the microorganism suspension (100mL) was swabbed onto agar plates (10mL agar). Excretion/ secretion products (25mL) were dispensed into the wells and the plates incubated at 35°C for 24h. Radial zones of inhibition (mm) of bacterial growth around the sample wells were measured to determine the antibacterial activity. The diameter of growth-inhibition zone was standardized by inhibition zone of Ampicillin used as a positive control for Gram-positive bacterial strains, Gentamycin used as a positive control for Gram-negative bacterial strains and Amphotericin B used as a positive control for fungi occurrence (Holder and Boyce, 1994). The growth of inhibition zone was determined according Bulet *et al.*, (1991).

Results

The antimicrobial activity of maggots ES against tested organisms:

Gram positive bacteria: The highest antibacterial (growth-inhibitory) activity of the maggots ES against Gram-positive bacterial was detected in *Staphylococcus aureus*, where the mean growth-inhibition zone was 18.3±1.2 mm vs. 27.4±1.5 mm for Ampicillin antibiotic, then followed by the *Bacillus subtilis*. The mean growth-inhibition zone was 22.6±0.58mm vs. 32.4±1.2 mm for the same antibiotic (Tab. 1 & fig. 1 A&B)

No antibacterial activity was detected in *Staphylococcus epidermidis* and *Streptococcus pyogenes* (Tab. 1 & Fig. 1 C&D).

Gram-negative bacteria:

The highest antibacterial growth-inhibitory activity against the bacterial strain Gram-negative was detected in *Escherichia coli*, where, the mean growth-inhibition zone was 20.4±1.0 mm vs. 22.3±0.72 mm for Gentamicin antibiotic, and then followed by the *Pseudomonas aeruginosa*, where, the mean growth-inhibition zone was 18.3±1.5mm vs. 22.6±1.5 mm for the same antibiotic (Tab. 2 & fig. 2 A&B)

No antibacterial activity was detected in *Proteus vulgaris* and *Klebsiella pneumonia* (Tab. 2 & Fig. 2 C&D)

Fungi: The most sensitive strain of tested fungi was detected in *Geotricum candidum*, where the mean growth-inhibition zone was 21.3±1.5mm vs. 23.2±2.1mm for Amphotericin B antibiotic, and then followed by the *Aspergillus fumigatus*, where as the mean growth-inhibition zone was 18.3±2.1mm vs. 22.6±1.5mm for the same antibiotic (Tab. 3 & Fig. 3 A&B)

No antifungal activity was detected in *Candida albicans* and *Syncephalastrum racemosum* (Tab. 3 & Fig. 3 C&D).

Discussion

Lucilia sericata larvae possess antibacterial agents in their excretion/secretion, for this reason the antibacterial activity of these larvae was three times than that of *Sarcophaga carnaria*. These findings are in consistent with the observations noticed by Thomas *et al.* (1999), where they showed that, *Lucilia sericata* larvae were able to kill or decrease the total bacterial count of *Staphylococcus aureus* in vitro and to combat clinical infections in a variety of wound types including these caused by antibiotic resistance strains of bacteria, in agreement with the current study.

The results of the present study are comparable with those obtained by Jaklic *et al.*, (2008). They carried out in vitro and in vivo quantitative research to assess the effect of larval (ES) of *Lucilia sericata* on bacterial strains *Staphylococcus aureus*, most commonly colonizing the chronic wounds.

In agreement with the present study, Bexfield *et al.* (2004) showed that the excretion/secretion of *L. sericata* larvae has potent antibacterial activity against some pathogenic bacteria, including MRSA. Also, they concluded that the extract of the housefly, *Musca domestica* generally possess wide broad antibacterial activity against both Gram+ve and Gram-ve bacteria, this conclusion was in consistence with the present study.

The antibacterial properties of secretions aseptically collected from larvae of the greenbottle fly, *Lucilia sericata* (Meigen) (Diptera: Calliphoridae) were examined by Kerridge *et al.* (2005) and they observed that there is antibacterial effective against Gram-positive *Staphylococcus aureus* and *Streptococcus pyogenes*, Gram-negative *Pseudomonas aeruginosa*, the present study is similar to the author in case of Gram-positive *Staphylococcus aureus* and *Streptococcus pyogenes* but, in consistent with Gram-negative *Pseudomonas aeruginosa*.

Daeschlein *et al.*, (2007) found that the secretions of maggots, *L. sericata* are known to have antibacterial properties against *Micrococcus luteus*, *E. coli* and *S. aureus*, and this is similar with present study.

Generally, some of insect body extracts exhibit antibacterial activity against both Gram-positive and Gram-negative bacteria, for example, the silk worm, *Bombyx mori* (Hara and Yamakawa, 1995), the European bumblebee, *Bombus pascuorum* (Rees *et al.*, 1997) their conclusion was in consistence with the present study. However, some insect species showed activity against only Gram-positive bacteria; for example; *Aedes aegypti* (Low-

enberger *et al.*, 1995) and *Chironomus plumosus* (Lauth *et al.*, 1998).

Besides, Vizioli *et al.* (2001) reported that *Anopheles gambiae* and *Phormia terranova* defensins displayed antibacterial activity spectra similar to that of other insect defensins as reported by Cociancich *et al.* (1994) against most of bacterial strains tested.

Killing of bacteria in the digestive tract of the maggot of *Lucilia sericata* (Diptera: Calliphoridae) was concluded by Mumcuoglu *et al.* (2001), indicated that the feces were either sterile or contained only small numbers of bacteria, this conclusion was in consistence with the present study.

In agreement with the present study Bexfield *et al.* (2007) demonstrated in vitro antibacterial activity of native excretion/ secretions (nES) from *L. sericata* against *B. cereus*, *E. coli* and *S. aureus*.

Evaluation of maggot excretions/ secretions are differentially effective against *S. aureus* and *P. aeruginosa* (Van der Plas *et al.*, 2008), their results are similar with the present study.

Antifungal activity of maggots and their secretion: Hou *et al.* (2007) reported that the extract of the housefly larvae showed higher activity against Gram-positive bacteria than Gram-negative bacteria and did not show any antifungal activity and the present study disagreement with these results. In consistent with the present results,

The present study showed variable antifungal activity of excretion/secretion of *Lucilia sericata* against the filamentous fungus, *Asperigillus fumigatus*, the yeast, *Candida albicans*, *Synecephalastrum racemosum* and *Geotricum*. The different (ES) samples of *L. seictata* possessed the highest antifungal activities as compared with antibacterial activity, and possessed the highest antibacterial activities Gram-ve as compared with the Gram+ve, and also, showed that the bacterial strains tested were more sensitive to the dif-

ferent insect larvae used than the fungal strains tested. In agreement with these results, Meylaers *et al.* (2004) observed that, the last instar larvae of the housefly, *Musca domestica* displayed antifungal activity against, the *Saccharomyces cerevisiae* beside the antibacterial activity.

The present study declared the antibacterial activity against Gram+ve, Gram-ve bacteria and Fungi. In agreement with the present, Leem *et al.* (1999) showed that saw fly, *Acantholyda parki* extract was found to have a broad antibacterial spectrum against not only Gram-ve but also Gram+ve bacteria.

Conclusion

The outcome results showed that excretion and secretion of the green bottle fly larvae *Lucilia sericata* (Meigen) gave antimicrobial activities. Extensive work is ongoing and will be published in due time.

References

- Bexfield, A, Bond, AE, Roberts, EC, et al, 2007:** The antibacterial activity against MRSA strains and other bacteria of a < 500 kDa fraction from maggot excretions/secretions of *Lucilia sericata* (Diptera: Calliphoridae). *Microbes Infect.* 10:325-33.
- Bexfield, A, Nigam, Y, Thomas, S, et al, 2004:** Detection and partial characterization of two antibacterial factors from the excretion/secretions of the medicinal maggot *Lucilia sericata* and their activity against methicillin-resistant *Staphylococcus aureus*. *Microbes Infect.* 6:1297-304.
- Bohova, J, Majtan, J, Majtan, V, et al, 2014:** Selective antibiofilm effects of *Lucilia sericata* larvae secretions/excretions against wound pathogens. *Evid Based Complement Alternat Med.* 0.1155/2014/857360.
- Bulet, P, Hetru, C, Dimarcq, JL, et al, 1991:** Antimicrobial peptides in insects; structure and function. *DCI:*329-44.
- Cazander, G, Van Veen, KEB, Bernards, A T, et al, 2009:** Do maggots have an influence on bacterial growth? A study on the susceptibility of strains of six different bacterial species to maggots of *Lucilia sericata* and their excretions/secretions. *J. Tissue Viabil.* 18, 3: 80-7.
- Cociancich, S, Dupont, A, Hegy, G, et al, 1994:** Novel inducible antibacterial peptides from a hemipteran insect, the sap-sucking bug *Pyrrhocoris apterus*. *Biochem. J.* 300: 567- 75.
- Daeschlein, G, Mumcuoglu, KY, Assadian, O, et al, 2007:** In vitro antibacterial activity of *Lucilia sericata* maggot secretions. *Skin Pharmacol Physiol,* 20, 2 : 112-5.
- Hara, S, and Yamakawa, M, 1995:** A novel antibacterial peptide family isolated from the silkworm, *Bombyx mori*. *Biochem. J.* 310, 2: 651-5.
- Harris, LG, Nigam, Y, Sawyer, J, et al, 2013:** *Lucilia sericata* chymotrypsin disrupts protein adhesion-mediated staphylococcal biofilm formation. *Appl. Environ. Microbiol.* 79, 4:1393-5.
- Holder, IA, and Boyce, ST, 1994:** Agar well diffusion assay testing of bacterial susceptibility to various antimicrobials in concentrations non-toxic for human cells in culture. *Burns.* 20, 5:426-9.
- Hou, L, Shi, Y, Zhai, P, et al, 2007:** Antibacterial activity and in vitro anti-tumor activity of the extract of the larvae of the housefly (*Musca domestica*). *J. Ethnopharmacol.* 111:227-31.
- Jaklic, D, Lapanje, A, Zupancic, K, et al, 2008:** Selective antimicrobial activity of maggots against pathogenic bacteria. *J. Med. Microbiol.* 57, 5:617-25.
- Kerridge, A, Lappin-Scott, H, and Stevens, JR, 2005:** Antibacterial properties of larval secretions of the blowfly, *Lucilia sericata*. *Med. Vet. Entomol.* 19, 3: 333-7.
- Lauth, X, Nesin A, Briand J, et al, 1998:** Isolation, characterization and chemical synthesis of a new insect defensin from *Chironomus plumosus* (Diptera). *Insect Biochem. Mol. Biol.* 1059-66.
- Leem, JY, Jeong IJ, Prak, KT, et al, 1999:** Isolation of P-hydroxycinnamaldehyde as an antibacterial substance from the saw fly, *FEBS Lett.* 442: 53-56.
- Lowenberger, C, Bulet, P, Charlet, M, et al, 1995:** Insect immunity: Isolation of three novel inducible antibacterial defensins from the vector mosquito, *Aedes aegypti*. *Insect Biochem. Mol. Biol.* 25, 7:867-73.
- Meylaers, K, Clynen, E, Daloze, D, et al, 2004:** Identification of 1-lysophosphatidylethanolamine (C16:1) as an antimicrobial compound in the housefly, *Musca domestica*. *Insect Biochem. Mol. Biol.* 34, 1:43-9.
- Mumcuoglu, KY, Miller, J, Mumcuoglu, M, et al, 2001:** Destruction of bacteria in the digestive tract of the maggot of *Lucilia sericata* (Diptera: Calliphoridae). *J. Med. Entomol.* 38, 2:161-6.

Rees, JA, Moniatte, M, and Bulet, P, 1997: Novel antibacterial peptides isolated from a European Bumblebee *Bombus pascuorum* (Hymenoptera, Apoidea). *Insect Biochem. Mol. Biol.* 27, 5:413-22.

Thomas, S, Andrews, AM, Hay, NP, et al, 1999: The anti-microbial activity of maggot secretions: results of a preliminary study. *J. Tissue Viabil.* 9, 4:127-32.

Valachova, J, Bohova, M, Kozanek, P, et al, 2013: *Lucilia sericata* medicinal maggots: a new

source of antimicrobial compounds. *Micro. Path.* 17:45-53.

Van der Plas, MJ Jukema, GN, Wai, SW, et al, 2008: Maggot excretions/secretions are differentially effective against biofilms of *Staphylococcus aureus* and *Pseudomonas aeruginosa*. *J. Antimicrob Chemother.* 61, 1:117-22.

Vizioli, J, Bulet, P, Hoffmann, JA, et al, 2001: Gambicin: A novel immune responsive antimicrobial peptide from the malaria vector *Anopheles gambiae*. *Proc. Natl. Acad. Sci. USA.* 98, 22:12630-5.

Table 1: Antibacterial activity as indicated by growth-inhibition zone of maggots (ES) of *L. sericata*, against Gram-positive bacteria.

Gram-positive Bacteria	Inhibition Zone/mm	Mean±SD Ampicillin (control)
<i>Staphylococcus aureus</i>	18.3±1.2	27.4±1.5
<i>Bacillus subtilis</i>	22.6±0.58	32.4±1.2
<i>Staphylococcus epidermidis</i>	-	22.4±1.5
<i>Streptococcus pyogenes</i>	-	23.1±2.3

SD: Standard deviation - : No inhibition zone

Table 2: Antibacterial activity as indicated by growth-inhibition zone of maggots (ES) of *L. sericata*, against Gram-negative bacteria.

Gram-negative bacteria	Inhibition Zone/mm	Mean±SD Gentamicin (control)
<i>Pseudomonas aeruginosa</i>	18.3±1.5	22.6±1.5
<i>Escherichia coli</i>	20.4±1.0	22.3±0.72
<i>Proteus vulgaris</i>	-	24.2±1.2
<i>Klebsiella pneumonia</i>	-	23.2±2.1

SD: Standard deviation - : No inhibition zone

Table 3: Antifungal activity as indicated by growth-inhibition zone of ES of *L. sericata* maggots against strains of fungi.

Fungi	Inhibition Zone/mm	Mean±SD Amphotericin B (control)
<i>Aspergillus fumigates</i>	18.3±2.1	22.6±1.5
<i>Geotricum candidum</i>	21.3± 1.5	23.2±2.1
<i>Candida albicans</i>	-	24.2±1.2
<i>Syncephalastrum racemosum</i>	-	22.3±0.72

SD: Standard deviation

- : No inhibition zone

Explanation of figures

Fig. 1: Excretion and secretion effect of *L. sericata* larvae against Gram-positive bacteria (A) *Staphylococcus aureus*. (B) *Bacillus subtilis*. (C) *Staphylococcus epidermidis*. (D) *Streptococcus pyogenes*.

Fig. 2: Excretion and secretion effect of *L. sericata* larvae against Gram-negative bacteria (A) *Pseudomonas aeruginosa*. (B) *Escherichia coli*. (C) *Klebsiella pneumonia*. (D) *Proteus vulgaris*.

Fig. 3: Excretion and secretion effect of *L. sericata* larvae against fungal strains (A) *Geotricum candidum*. (B) *Aspergillus fumigates*. (C) *Candida albicans*. (D) *Syncephalastrum racemosum*.

