

EFFECT OF ORGANIC MANURE ENRICHED WITH MACRO AND MICRONUTRIENTS:

II. NUTRIENTS UPTAKE BY WHEAT AND THEIR AVAILABILITY IN SAND SOIL

Ali, M.E.*;S.A.Ismail**O.H.M.El-Hussieny*and A.M.Abd El-Hafeez**

* Dept. of Soils, Fac. of Agric. Benha, Univ.

** Soil, Water and Environment Res. Inst., A.R.C., Egypt.

ABSTRACT

Two field experiments were performed at Village No. 1, west of Samallote, Minia Governorate, Egypt during 2004/2005 and 2005/2006 seasons to study the effect of four levels of farmyard manure (0, 10, 20 and 30 ton/fed); NPK fertilization doses, (i.e. 0, low (60/15/24) and high (120/30/48) kg of N, P₂O₅ and K₂O/fed) and the micronutrients (Fe, Mn, Zn and Cu) which were applied, soil application or as foliar spraying on N, P, K, Fe, Mn, Zn and Cu uptake by wheat plants (Sids 1 cv) grown on a sandy soil and their availability in soil. The results could be summarized as follows;

Uptake of N, P, K, Fe, Mn, Zn and Cu by wheat plants significantly increased with increasing rate of the applied FYM and NPK, as well as the micronutrients application, where foliar spraying surpassed soil application in both seasons.

The soil availability of N, P and K responded to manuring only. However, phosphorus only significantly increased by increasing NPK fertilization.

Soil available contents of Fe, Mn, Zn and Cu were significantly affected by manuring and micronutrients as soil application, while foliar spray did not affect contents of these micronutrient in both seasons.

It could be concluded that wheat under sand soils should be fertilized with 30 ton FYM/fed; NPK at rate of 120 kg N, 30 kg P₂O₅ and 48 kg K₂O/fed. Also, the micronutrients should be applied to improve both nutrients uptake by wheat plants and their availability in soil.

Keywords: wheat, sand soils, farmyard manure, nutrients uptake, nutrients availability.

INTRODUCTION

It is well known in Egypt that the expansion in wheat cultivation in sand soils is the best solution for curtailing the gap between consumption and production of wheat. However, production of wheat in the sand soils is facing many problems, among them the low clay and organic matter contents and hence the poor soil fertility.

Improving the poor physical, chemical and nutritional properties of these soils are a compacted problem that requires enormous efforts. Using organic manure has been proved to be of vital importance in this concern. Atta Allah and Mohamed (2003), Salem (2003) and Khalil and Aly (2004) reported that N, P and K uptake significantly increased with adding organic manure to wheat plants. Also, El-Koumey (1998) and Mekail *et al.* (2006) stated that organic manure fertilization increased nutrients uptake. On the other hand, soil available nutrients were markedly affected by soil manuring (Negm *et al.*, 2002, El-Ghamry *et al.*, 2005, Ali, Maha, 2007 and Taha, 2007).

Inorganic fertilizers are considered an important source for nutrients in soil where very few soils are able to supply sufficient quantities of all the nutrients. Many workers stated that N, P and K uptake by wheat plants were significantly affected by increasing N, P and K levels (El-Leithi *et al.*, 1996 and Galal, 2007).

Micronutrients application is now a days of considered importance in soils of Egypt, especially the sandy ones where an adequate nutrients supply can be anticipated in the newly reclaimed sandy soils. Macronutrients as well as micronutrients are constantly removed with the harvest of high yielding varieties which require high doses of NPK and substantial quantities of micronutrients. On the other hand, the efficiencies of micronutrients are mainly dependent on the method of application. The alternative approaches for the application of micronutrients includes: soil and foliar application. The positive effect of micronutrients on nutrient uptake were reported by Badr *et al.* (1991), Mohamed (1994) and Galal (2007). However, the superiority of foliar spray than soil application for improving nutrients uptake was mentioned by Shams El-Din (1993).

The objective of the present investigation was to study the influence of different levels of FYM and NPK fertilization as well as methods of micronutrients application on nutrients uptake by wheat plant and nutrients availability in sandy soil.

MATERIALS AND METHODS

Two field experiments were performed at Village No. 1, west of Samallote, Minia Governorate, Egypt (newly reclaimed sand soils) during the winter seasons of 2004/2005 and 2005/2006, to study the response of wheat (*Triticum aestivum* L.), Sids 1 cv to four levels of FYM (0, 10, 20 and 30 ton/fed), three doses of NPK fertilization, i.e. without NPK, low (60/15/24) and high (120/30/48) kg doses N, P₂O₅ and K₂O/fed, respectively. Some micronutrients (Fe, Mn, Zn and Cu) were applied as soil or foliar application. The soil physical and chemical properties of the experimental field were determined as described by Klute (1982) and Page *et al.* (1982) (Table, 1).

Farmyard manure (A) was applied before planting and its chemical compositions within the first and second seasons are shown in Table (2). Nitrogen fertilizer (B) was added as ammonium nitrate (33.5% N) at six equal doses, the first was applied before the first irrigation and the others were added before every irrigation. Phosphorus and potassium were added before planting as mono calcium phosphate (15.5% P₂O₅) and potassium sulphate (48% K₂O), respectively. The micronutrients (C) were added as soil application (10 kg Fe, Mn or Zn sulphate/fed and 5 kg Cu sulphate/fed) before planting whereas foliar spraying was conducted twice, at 45 and 60 days from planting with volume of 400 L/fed, each liter contain 3 g of Fe, Mn or Zn or 1 g Cu sulphate (about 0.6, 1.0, 0.7 and 0.25 g Fe, Mn, Zn and Cu, respectively). Normal agronomic practices were conducted as usual for wheat production in studied region.

Table (1): Some physical and chemical properties of the studied soil before planting of wheat in the two growing seasons.

Soil properties	Season	
	First season	Second season
Particle size distribution%		
Clay (%)	10.94	11.84
Silt (%)	12.75	9.35
Sand (%)	76.31	78.81
Texture grade	Sandy loam	Sandy loam
Chemical properties		
pH*	8.00	7.98
EC (dS m ⁻¹)**	1.60	1.65
Organic matter (%)	0.85	0.78
Ca CO ₃ (%)	4.20	3.94
Soluble Ca ²⁺ m mol L ⁻¹ ***	5.62	5.67
Soluble Mg ²⁺ m mol L ⁻¹	5.21	5.42
Soluble Na ⁺ m mol L ⁻¹	2.57	2.90
Soluble K ⁺ m mol L ⁻¹	2.16	2.58
Soluble CO ₃ ²⁻ m mol L ⁻¹	0.00	0.00
Soluble HCO ₃ ⁻ m mol L ⁻¹	1.95	2.29
Soluble Cl ⁻ m mol L ⁻¹	6.89	7.29
Soluble SO ₄ ²⁻ m mol L ⁻¹	6.72	6.99
Available N (ug/g)	2.33	2.95
Available P (ug/g)	6.12	8.01
Available K (ug/g)	7.35	9.62
Available Fe (ug/g)	2.39	2.13
Available Mn (ug/g)	1.36	1.12
Available Zn (ug/g)	0.88	0.78
Available Cu (ug/g)	0.42	0.43

* Measured in 1:2.5 soil water suspension.

** Measured in soil paste.

*** Measured in soil paste.

Table (2): Chemical composition of F.Y.M.

Properties	1 st Season	2 nd Season
Organic matter %	28.50	25.60
Total N%	1.68	1.42
C/N ratio	1:17	1:18
Total N %	0.62	0.56
Total P %	0.19	0.18
Total K %	0.71	0.66
Available Fe (ug/g)	5.2	6.7
Available Mn (ug/g)	3.1	4.6
Available Zn (ug/g)	1.8	2.0
Available Cu (ug/g)	0.9	0.6

At harvest the plots were hand-harvested and plant samples were taken from grains and straw to determine nutrient concentrations according to Jackson (1973), then total nutrients uptake were calculated.

The available soil N, P, K, Fe, Mn, Zn and Cu were determined according to Klute (1982). The data were subjected to statistical analysis according to Sendecor and Cochran (1980). The treatment means were compared by L.S.D. at 5% level of probability.

RESULTS AND DISCUSSION

Nutrients uptake:

The results tabulated in Tables (3-9) showed the effect of manuring and NPK fertilization as well as different methods of micronutrients application on N, P, K, Fe, Mn, Zn and Cu uptake by wheat plants. It is evident that nutrients uptake significantly increased due to FYM application in both seasons. The highest values of the studied nutrients uptake were achieved when wheat plants received 30 ton/fed followed by those supplied with 20 and 10 ton/fed, respectively while the plants without manuring recorded the lowest values of nutrients uptake. This could be due to the decomposition of FYM which resulted in chelated compounds enhanced the availability and uptake of the micronutrients. In addition, FYM contains sufficient amounts of most plant nutrients which were released in available forms upon its decomposition. Similar results were obtained by El-Koumey (1998) and Mekail *et al.* (2006).

Table (3): Nitrogen uptake (kg/fed) as affected by organic manure, NPK fertilization treatments and methods of application of the micronutrients.

Rate of applied O.M (A) ton/fed	Rate of N/P ₂ O ₅ /K ₂ O application (kg/fed) (B)	Micronutrients (C)							
		First season				Second season			
		Control 0.0	Soil appli.	Foliar appli.	Mean	Control 0.0	Soil appli.	Foliar appli.	Mean
0.0	0.0	15.49	16.58	19.03	17.03	13.32	13.28	15.57	14.06
	60/15/24	21.04	22.73	23.76	22.51	18.55	18.85	19.94	19.12
	120/30/48	36.10	37.84	39.51	37.82	28.52	31.46	32.27	30.75
	Mean	24.21	25.72	27.43	25.79	20.13	21.20	22.59	21.31
10	0.0	25.67	26.57	30.15	27.46	21.51	22.09	24.13	22.57
	60/15/24	33.08	33.18	33.57	33.27	27.09	28.32	28.61	28.01
	120/30/48	47.83	48.61	49.87	48.77	34.91	35.14	37.08	35.71
	Mean	35.52	36.12	37.86	36.49	27.83	28.52	29.94	28.76
20	0.0	35.09	35.65	37.86	36.20	31.81	33.09	33.86	32.92
	60/15/24	40.39	42.62	44.45	42.49	35.87	37.19	39.08	37.38
	120/30/48	50.78	51.40	54.97	52.38	43.52	44.28	45.76	44.52
	Mean	42.08	43.22	45.76	43.69	37.07	38.19	39.57	38.28
30	0.0	34.93	36.56	37.90	36.46	41.74	43.88	43.09	42.90
	60/15/24	44.82	45.91	45.94	45.55	45.83	46.36	47.32	46.50
	120/30/48	57.29	58.66	61.35	59.10	53.77	54.80	55.43	54.67
	Mean	45.68	47.04	48.40	47.04	47.11	48.35	48.61	48.02
Mean of NPK	0.0	28.00	28.84	31.23	29.36	27.10	28.09	29.16	28.12
	60/15/24	34.83	36.11	36.93	35.96	31.83	32.68	33.73	32.75
	120/30/48	48.00	48.13	51.42	49.52	40.18	41.42	42.63	41.41
Mean of micronutrients		36.87	38.02	39.86		33.04	34.07	35.18	

L.S.D at 0.05	First season							Second season						
	A	B	C	AB	AC	BC	ABC	A	B	C	AB	AC	BC	ABC
	0.57	0.46	0.30	0.91	N.S	N.S	1.28	0.80	0.50	0.34	1.02	N.S	N.S	1.35

Table (4): Phosphorus uptake (kg/fed) as affected by organic manure, NPK fertilization treatments and methods of application of the micronutrients.

Rate of applied O.M (A) ton/fed	Rate of N/P ₂ O ₅ /K ₂ O application (kg/fed) (B)	Micronutrients (C)							
		First season				Second season			
		Control 0.0	Soil appli.	Foliar appli.	Mean	Control 0.0	Soil appli.	Foliar appli.	Mean
0.0	0.0	5.64	6.13	6.50	6.09	4.55	4.60	5.76	4.97
	60/15/24	7.02	7.43	7.49	7.31	6.36	6.41	7.06	6.61
	120/30/48	11.01	11.28	11.53	11.27	8.51	9.77	9.97	9.42
	Mean	7.89	8.28	8.50	8.22	6.47	6.96	7.59	7.00
10	0.0	9.04	9.07	10.34	9.48	7.12	7.50	7.99	7.54
	60/15/24	10.50	11.20	11.11	10.93	8.33	8.90	9.38	8.87
	120/30/48	13.95	15.00	15.79	14.91	10.29	10.38	10.59	10.42
	Mean	11.16	11.76	12.41	11.78	8.58	8.89	9.32	8.94
20	0.0	12.06	12.13	12.63	12.27	10.20	11.07	11.66	10.98
	60/15/24	12.71	13.93	14.06	13.56	11.06	11.79	12.12	11.65
	120/30/48	15.00	15.47	16.87	15.78	12.06	12.74	13.26	12.69
	Mean	13.26	13.84	14.52	13.87	11.10	11.87	12.35	11.77
30	0.0	13.47	14.22	14.29	13.99	12.98	13.35	13.60	13.31
	60/15/24	13.92	14.64	15.06	14.54	13.36	13.84	14.83	14.02
	120/30/48	16.33	17.38	17.74	17.15	14.30	14.74	14.62	14.55
	Mean	14.57	15.41	15.70	15.22	13.55	13.98	14.35	13.96
Mean of NPK	0.0	10.05	10.39	10.94	10.46	8.71	9.13	9.75	9.20
	60/15/24	11.04	11.80	11.93	11.59	9.78	10.23	10.85	10.28
	120/30/48	14.07	14.78	15.48	14.78	11.29	11.91	12.11	11.77
Mean of micronutrients		11.72	12.32	12.78		9.99	10.42	10.87	

L.S.D at	First season							Second season						
	A	B	C	AB	AC	BC	ABC	A	B	C	AB	AC	BC	ABC
0.05	0.08	0.15	0.12	0.37	N.S	0.27	0.54	0.24	0.17	0.16	0.37	N.S	0.30	0.63

As for NPK fertilization, the data at the same Tables revealed that N, P, K, Fe, Mn, Zn and Cu uptake by wheat plants were significantly affected by the level of N, P and K fertilization. The maximum nutrients uptake were obtained with the highest doses of NPK fertilization, i.e. 120/30/48 kg N, P₂O₅ and K₂O/fed, while the wheat plants without NPK application recorded the lowest uptake ones. Such increases in uptake values of all the nutritive elements, is mainly due to N, P and K are considered the most important nutrients for plants. Consequently, their application increased both grains and straw yields, which in turn increased nutrients uptake since nutrient uptake is calculated as multiplying nutrient concentration by grain and straw yields. The results are in line with those obtained by El-Leithi *et al.* (1996) and Galal (2007).

Table (5): Potassium uptake (kg/fed) as affected by organic manure, NPK fertilization treatments and methods of application of the micronutrients.

Rate of applied O.M (A) ton/fed	Rate of N/P ₂ O ₅ /K ₂ O application (kg/fed) (B)	Micronutrients (C)							
		First season				Second season			
		Control 0.0	Soil appli.	Foliar appli.	Mean	Control 0.0	Soil appli.	Foliar appli.	Mean
0.0	0.0	27.92	30.48	33.09	30.50	24.93	25.86	27.12	25.97
	60/15/24	41.55	45.71	47.40	44.89	35.87	37.45	38.77	37.36
	120/30/48	65.33	72.46	73.71	70.50	45.37	47.98	48.39	47.25
	Mean	44.93	49.55	51.40	48.62	35.39	37.10	38.10	36.86
10	0.0	45.38	46.25	48.51	46.72	38.39	38.93	40.22	39.18
	60/15/24	52.94	54.89	58.80	55.04	46.09	47.49	47.74	47.11
	120/30/48	78.53	79.91	84.05	80.83	54.61	55.83	57.10	55.85
	Mean	58.95	60.35	63.79	61.03	46.36	47.42	48.35	47.38
20	0.0	44.28	49.65	56.57	50.17	48.39	50.71	51.67	50.26
	60/15/24	63.29	67.03	68.19	66.17	56.09	57.84	58.15	57.36
	120/30/48	76.09	80.89	85.36	80.78	62.18	64.54	71.89	66.21
	Mean	61.22	65.85	70.04	65.70	55.55	57.70	60.57	57.94
30	0.0	54.76	60.65	62.23	59.21	56.99	59.39	59.94	58.77
	60/15/24	63.43	68.98	70.39	67.00	66.33	67.11	68.39	67.27
	120/30/48	78.17	88.38	89.39	85.31	73.35	73.66	74.37	73.79
	Mean	65.32	72.67	74.00	70.66	65.55	66.72	67.57	66.61
Mean of NPK	0.0	43.08	46.75	50.10	46.64	42.18	43.72	43.99	43.29
	60/15/24	55.30	59.15	61.19	58.54	51.09	52.47	53.26	52.28
	120/30/48	74.53	79.91	82.63	79.02	58.88	60.50	62.94	60.77
Mean of micronutrients		57.61	61.94	64.64		50.71	52.24	53.42	

L.S.D at 0.05	First season							Second season						
	A	B	C	AB	AC	BC	ABC	A	B	C	AB	AC	BC	ABC
	2.10	1.77	1.20	3.60	N.S	N.S	4.75	0.69	1.19	0.60	2.43	N.S	N.S	2.58

As regard to micronutrients, the data obtained clearly showed that nutrients uptake values were significantly affected by micronutrients application. It is worthy to notice that micronutrients application as foliar spray resulted in nutrient uptake values higher than the soil application method. Mohamed (1994) mentioned that foliar application of micronutrients was more effective specially under the Egyptian condition, where these nutrients are converted to unavailable forms if they applied to the soil. These results are in good agreement with those obtained by Shams El-Din (1993).

Concerning the interaction, data, in general showed that the highest values of N, P, K, Fe, Mu, Zn and Cu uptake by wheat plants were recorded for the plants sprayed with micronutrients under 30 ton FYM/fed and 120/30/48 kg N, P₂O₅ and K₂O/fed. The plants without manuring or micronutrients on the other hand, recorded the lowest values of the studied nutrient uptake.

Table (6): Iron uptake (g/fed) as affected by organic manure, NPK fertilization treatments and methods of application of the micronutrients.

Rate of applied O.M (A) ton/fed	Rate of N/P ₂ O ₅ /K ₂ O application (kg/fed) (B)	Micronutrients (C)							
		First season				Second season			
		Control 0.0	Soil appli.	Foliar appli.	Mean	Control 0.0	Soil appli.	Foliar appli.	Mean
0.0	0.0 60/15/24 120/30/48	1774.33	2264.15	2747.85	2262.11	1458.25	1707.76	2153.43	1773.15
		2403.22	3137.08	3574.23	3038.17	1871.45	2173.38	2616.57	2220.47
		3631.20	4699.77	4700.30	4343.75	2371.66	2844.54	3275.57	2830.59
		Mean	2602.91	3367.00	3674.13	3259.12	1900.45	2241.89	2681.85
10	0.0 60/15/24 120/30/48	2734.93	3194.13	3738.10	3222.39	2351.78	2702.30	3178.94	2744.34
		3168.55	3751.62	4179.85	3700.09	2431.82	2996.64	3403.79	2944.08
		3582.74	5382.73	4762.80	4276.09	2704.27	3463.02	3719.68	3295.65
		Mean	3162.07	4109.49	4226.91	3855.04	2495.95	3053.98	3434.13
20	0.0 60/15/24 120/30/48	3082.05	3247.29	3793.31	3374.22	2894.20	3461.85	3722.95	3359.67
		3252.53	3781.85	4462.08	3832.15	3028.23	3605.47	4024.55	3552.75
		3633.40	5471.55	5923.83	5009.59	3052.10	3825.59	4351.08	3742.92
		Mean	3322.66	4166.90	4726.41	3905.32	2991.51	3630.97	4032.86
30	0.0 60/15/24 120/30/48	3679.29	4202.49	4754.93	4212.24	3298.26	4026.83	4457.13	3927.41
		3869.02	4218.73	5074.31	4387.36	3500.47	4064.08	4562.04	4042.19
		4243.65	5499.53	6017.29	5253.49	3606.06	4213.64	4723.06	4180.90
		Mean	3930.65	4640.25	5282.18	4617.70	3468.26	4101.52	4580.74
Mean of NPK	0.0 60/15/24 120/30/48	2817.65	3352.01	3758.33	3309.33	2500.65	2974.68	3428.11	2967.81
		3173.33	3722.32	4322.62	3739.42	2707.99	3209.89	3653.73	3190.54
		3822.74	5263.39	6369.56	5151.89	2933.52	3586.57	4017.34	3512.48
Mean of micronutrients		3271.24	4070.91	4477.41		2714.05	3257.09	3699.06	

L.S.D at 0.05	First season							Second season						
	A	B	C	AB	AC	BC	ABC	A	B	C	AB	AC	BC	ABC
	49.55	49.79	43.93	99.87	88.50	88.29	175.20	33.80	22.73	28.70	46.71	57.97	58.71	110.53

Table (7): Manganese uptake (g/fed) as affected by organic manure, NPK fertilization treatments and methods of application of the micronutrients.

Rate of applied O.M (A) ton/fed	Rate of N/P ₂ O ₅ /K ₂ O application (kg/fed) (B)	Micronutrients (C)							
		First season				Second season			
		Control 0.0	Soil appli.	Foliar appli.	Mean	Control 0.0	Soil appli.	Foliar appli.	Mean
0.0	0.0 60/15/24 120/30/48	604.92	777.49	966.41	782.94	500.12	628.47	823.03	650.54
		764.20	991.89	1159.88	971.99	665.29	842.96	974.83	827.69
		1196.96	1502.36	1687.78	1462.37	882.60	1192.55	1262.12	1112.42
		Mean	855.36	1090.58	1271.35	1072.43	682.67	887.99	1019.99
10	0.0 60/15/24 120/30/48	919.99	1201.69	1510.54	1210.74	801.63	945.03	1153.06	966.57
		1043.56	1360.86	1522.07	1308.83	937.26	1158.46	1327.47	1141.06
		1283.60	1866.81	2186.16	1778.86	1094.01	1231.54	1493.76	1273.10
		Mean	1082.38	1476.45	1739.59	1432.80	944.30	1111.69	1324.76
20	0.0 60/15/24 120/30/48	1232.89	1432.34	1690.35	1451.86	1078.61	1306.24	1584.09	1322.98
		1288.98	1569.19	1821.19	1559.79	1134.67	1483.74	1598.31	1405.57
		1468.60	1886.55	2257.66	1870.94	1216.91	1499.00	1698.97	1471.62
		Mean	1330.15	1629.36	1923.06	1627.52	1143.39	1429.66	1627.12
30	0.0 60/15/24 120/30/48	1439.00	1739.88	2001.05	1726.64	1347.50	1593.03	1755.70	1565.41
		1450.46	1867.08	2057.64	1791.73	1415.83	1756.90	1891.50	1688.08
		1667.87	2261.95	2644.70	2191.50	1552.16	1930.79	2046.83	1843.26
		Mean	1519.11	1956.30	2234.46	1903.29	1438.49	1760.24	1898.01
Mean of NPK	0.0 60/15/24 120/30/48	1049.20	1287.85	1542.08	1293.04	931.96	1118.19	1328.97	1126.37
		1136.80	1447.26	1640.95	1408.34	1038.26	1310.52	1448.03	1265.60
		1404.07	1879.42	2194.07	1825.85	1186.42	1463.45	1625.42	1425.10
Mean of micronutrients		1196.92	1538.17	1792.19		1052.21	1297.40	1467.47	

L.S.D at 0.05	First season							Second season						
	A	B	C	AB	AC	BC	ABC	A	B	C	AB	AC	BC	ABC
	36.16	37.39	27.31	76.39	57.71	57.71	N.S	52.10	34.02	12.97	69.19	37.89	37.89	N.S

Table (8): Zinc uptake (g/fed) as affected by organic manure, NPK fertilization treatments and methods of application of the micronutrients.

Rate of applied O.M(A) ton/fed	Rate of N/P ₂ O ₅ /K ₂ O application (kg/fed)(B)	Micronutrients (C)											
		First season				Second season							
		Control 0.0	Soil appli.	Foliar appli.	Mean	Control 0.0	Soil appli.	Foliar appli.	Mean				
0.0	0.0 60/15/24 120/30/48	800.63	941.53	1173.79	972.00	639.23	729.44	926.67	765.11				
		951.89	1145.98	1444.91	1180.93	863.30	908.98	1147.73	973.34				
		1353.12	1713.74	1909.70	1658.85	1153.44	1164.83	1555.70	1291.33				
		Mean	1035.21	1267.08	1509.47	1270.59	885.32	934.41	1210.03	1009.92			
10	0.0 60/15/24 120/30/48	1266.30	1458.71	1613.93	1446.31	996.51	1152.82	1408.46	1185.93				
		1457.33	1608.33	1865.25	1643.63	1114.52	1264.81	1541.17	1306.83				
		1637.67	1964.60	2575.62	2059.29	1123.92	1331.79	1792.17	1415.96				
		Mean	1453.76	1677.21	2018.27	1716.41	1078.31	1249.80	1580.60	1302.90			
20	0.0 60/15/24 120/30/48	1632.05	1806.43	2029.71	1822.73	1254.33	1486.48	1875.87	1538.89				
		1647.40	1868.90	2257.34	1924.55	1291.55	1494.16	2054.31	1613.40				
		1693.06	2113.02	2671.30	2159.13	1297.41	1628.83	2088.57	1671.60				
		Mean	1657.50	1929.45	2319.45	1968.80	1281.09	1536.49	2006.24	1607.94			
30	0.0 60/15/24 120/30/48	1650.74	1914.06	2154.39	1906.40	1582.85	1746.13	2292.88	1873.95				
		1754.62	1923.37	2215.94	1964.64	1628.10	1844.85	2311.24	1928.06				
		1996.16	2191.79	2823.48	2337.14	1711.44	1890.06	2468.59	2023.36				
		Mean	1800.50	2009.74	2397.93	2069.39	1640.81	1827.01	2357.57	1941.80			
Mean of NPK	0.0 60/15/24 120/30/48	1337.43	1530.18	1742.95	1536.85	1118.23	1303.39	1625.97	1349.20				
		1452.81	1636.64	1945.86	1678.44	1224.36	1378.20	1763.61	1455.39				
		1670.00	1995.79	2495.03	2053.61	1321.55	1503.87	1976.25	1600.56				
Mean of micronutrients		1486.74	1720.87	2061.28		1228.88	1386.95	1788.61					

L.S.D at	First season							Second season						
	A	B	C	AB	AC	BC	ABC	A	B	C	AB	AC	BC	ABC
0.05	86.99	52.89	49.30	105.71	N.S	99.53	N.S	33.24	43.43	40.30	79.39	81.31	81.31	160.21

Table (9): Copper uptake (g/fed) as affected by organic manure, NPK fertilization treatments and methods of application of the micronutrients.

Rate of applied O.M(A) ton/fed	Rate of N/P ₂ O ₅ /K ₂ O application (kg/fed)(B)	Micronutrients (C)											
		First season				Second season							
		Control 0.0	Soil appli.	Foliar appli.	Mean	Control 0.0	Soil appli.	Foliar appli.	Mean				
0.0	0.0 60/15/24 120/30/48	361.97	448.39	532.83	447.73	264.04	335.75	407.42	335.73				
		437.45	574.25	634.73	548.81	342.68	442.10	521.31	435.36				
		647.45	793.89	994.65	811.09	447.12	593.77	619.20	553.36				
		Mean	482.29	605.51	720.73	602.84	351.28	457.20	515.97	441.48			
10	0.0 60/15/24 120/30/48	464.49	578.07	773.42	605.32	396.14	429.38	645.11	490.21				
		588.71	719.30	861.21	723.07	431.88	538.15	704.06	558.04				
		714.90	846.19	1098.55	886.55	454.79	597.65	784.95	610.80				
		Mean	589.36	714.52	911.05	738.31	427.60	520.06	711.37	553.02			
20	0.0 60/15/24 120/30/48	533.45	611.50	879.46	674.80	418.31	498.48	736.79	551.18				
		595.82	739.50	965.33	766.88	439.32	539.59	776.75	582.22				
		727.51	894.70	1116.90	889.70	505.41	627.87	908.60	680.63				
		Mean	618.92	715.23	997.23	777.13	454.35	552.31	807.38	604.68			
30	0.0 60/15/24 120/30/48	549.11	682.37	952.95	728.14	425.33	593.38	933.59	650.76				
		635.48	797.25	1033.37	822.03	480.74	612.57	981.09	691.47				
		751.14	899.24	1120.32	923.56	515.39	629.34	1019.47	721.40				
		Mean	645.24	792.95	1035.54	824.57	473.82	611.76	978.05	687.88			
Mean of NPK	0.0 60/15/24 120/30/48	477.25	580.08	784.66	613.99	375.95	464.24	680.72	506.97				
		564.36	707.57	853.55	708.49	423.65	533.10	745.80	567.52				
		710.25	858.50	1082.60	803.78	480.67	612.15	833.05	641.96				
Mean of micronutrients		583.95	703.07	916.13		422.61	538.66	753.19					

L.S.D at 0.05	First season							Second season						
	A	B	C	AB	AC	BC	ABC	A	B	C	AB	AC	BC	ABC
	63.67	33.35	27.51	66.97	56.08	N.S	N.S	67.35	26.45	23.87	52.98	47.79	N.S	N.S

Macro nutrients availability:

Data presented in Tables (10-12) show the effect of manuring, N, P and K fertilization and micronutrients application on nitrogen, phosphorus and potassium availability in the investigated sand soil. The obtained data clearly reveal that farmyard manure increased soil contents of available N, P and K as compared with the treatment without manuring. These increases were proportional to the increase in application rate up to 30 ton FYM/fed. The significant increase in available studied macro nutrients caused by manuring could be due to producing N, P and K through organic materials decomposition, fixation of atmospheric N and their reflection on soil fertility. Also, such materials might increase the cation exchange capacity of sand soil and consequently reduce losses of nutrients by leaching. These results are in agreement with those obtained by El-Ghamry *et al.* (2005), Ali, Maha (2007) and Taha (2007).

Table (10): Soil available nitrogen (mg kg⁻¹) as affected by organic manure, NPK fertilization treatments and methods of application of the micronutrients.

Rate of applied O.M (A) ton/fed	Rate of N/P ₂ O ₅ /K ₂ O application (kg/fed) (B)	Micronutrients (C)							
		First season				Second season			
		Control 0.0	Soil appli.	Foliar appli.	Mean	Control 0.0	Soil appli.	Foliar appli.	Mean
0.0	0.0	2.25	2.42	2.35	2.34	2.60	2.82	2.75	2.72
	60/15/24	2.30	2.60	2.65	2.53	2.70	2.85	2.82	2.79
	120/30/48	2.60	2.75	2.70	2.68	2.73	2.90	2.85	2.83
	Mean	2.38	2.59	2.56	2.51	2.68	2.86	2.81	2.78
10	0.0	3.92	4.15	4.10	4.07	4.10	4.35	4.17	4.21
	60/15/24	4.15	4.27	4.22	4.21	4.12	4.37	4.22	4.24
	120/30/48	4.25	4.52	4.35	4.71	4.22	4.47	4.25	4.15
	Mean	4.10	4.31	4.22	4.21	4.15	4.40	4.21	4.25
20	0.0	4.25	4.72	4.60	4.53	5.50	5.60	5.55	5.45
	60/15/24	5.15	5.38	5.25	5.26	5.53	5.65	5.60	5.59
	120/30/48	5.45	5.80	5.75	5.66	5.55	5.90	5.80	5.75
	Mean	4.95	5.30	5.20	5.15	5.53	5.72	5.65	5.63
30	0.0	6.12	6.37	6.30	6.26	6.50	6.60	6.55	6.55
	60/15/24	6.27	7.52	6.32	6.70	6.70	6.80	6.70	6.73
	120/30/48	6.52	7.72	7.60	7.28	6.75	6.85	6.80	6.80
	Mean	6.30	7.20	6.74	6.75	6.65	6.75	6.69	6.69
Mean of NPK	0.0	4.13	4.41	4.34	4.29	4.68	4.84	4.75	4.76
	60/15/24	4.47	4.94	4.61	4.67	4.76	4.89	4.83	4.83
	120/30/48	4.71	5.20	5.10	5.00	4.81	5.03	4.92	4.92
Mean of micronutrients		4.43	4.85	4.68		4.75	4.98	4.84	

L.S.D at 0.05	First season							Second season						
	A	B	C	AB	AC	BC	ABC	A	B	C	AB	AC	BC	ABC
	0.11	N.S	N.S	N.S	N.S	N.S	N.S	N.S	0.21	N.S	N.S	N.S	N.S	N.S

As for macro and micronutrients application, the data showed that N, P and K availability was not affected by either macro or micronutrients fertilization treatments, except for phosphorus in both seasons where phosphorus availability significantly increased with increasing NPK levels up to the highest doses, i.e. 120/30/48 kg N, P₂O₅ and K₂O/fed, respectively. The effect of NPK fertilization on phosphorus availability is mainly due to increasing plant growth and consequently increasing values of increasing root growth system, which in turn enhanced nutrients uptake, consequently increase available B after it is decomposition.

Table (11): Soil available phosphorus (mg kg⁻¹) as affected by organic manure, NPK fertilization treatments and methods of application of the micronutrients.

Rate of applied O.M (A) ton/fed	Rate of N/P ₂ O ₅ /K ₂ O application (kg/fed) (B)	Micronutrients (C)							
		First season				Second season			
		Control 0.0	Soil appli.	Foliar appli.	Mean	Control 0.0	Soil appli.	Foliar appli.	Mean
0.0	0.0	5.80	6.18	6.35	6.11	7.12	7.23	7.20	7.18
	60/15/24	6.15	7.90	6.22	6.76	8.45	8.59	8.55	8.53
	120/30/48	8.00	8.10	7.87	7.99	9.65	9.87	9.75	9.76
	Mean	6.65	7.39	6.81	6.95	8.41	8.56	8.50	8.49
10	0.0	6.25	6.50	6.40	6.38	8.14	8.35	8.25	8.25
	60/15/24	7.87	8.00	8.12	7.99	9.40	9.65	9.57	9.54
	120/30/48	8.35	8.87	8.70	8.64	10.60	10.75	10.65	10.67
	Mean	7.49	7.79	7.74	7.67	9.38	9.58	9.49	9.48
20	0.0	7.35	7.65	7.45	7.48	8.50	8.72	8.57	8.60
	60/15/24	8.35	8.57	8.50	8.49	9.63	9.67	9.67	9.66
	120/30/48	9.10	9.72	9.65	9.49	10.70	10.85	10.75	10.77
	Mean	8.26	8.65	8.53	8.48	9.61	9.75	9.66	9.67
30	0.0	8.12	8.35	8.55	8.34	9.57	9.75	9.67	9.66
	60/15/24	9.12	9.50	9.35	9.32	10.80	10.95	10.90	10.88
	120/30/48	10.20	10.90	10.29	10.46	11.10	11.32	11.22	11.21
	Mean	9.14	9.58	9.40	9.37	10.49	10.67	10.60	10.59
Mean of NPK	0.0	6.88	7.17	7.19	7.08	8.33	8.51	8.42	8.42
	60/15/24	7.87	8.49	8.05	8.14	9.57	9.71	9.67	9.65
	120/30/48	8.91	9.40	9.13	9.15	10.51	10.70	10.59	10.60
	Mean of micronutrients	7.88	8.35	8.12		9.43	9.65	9.54	

L.S.D at 0.05	First season							Second season						
	A	B	C	AB	AC	BC	ABC	A	B	C	AB	AC	BC	ABC
	0.11	0.07	N.S	0.14	N.S	N.S	N.S	0.19	0.15	N.S	0.31	N.S	N.S	N.S

Table (12): Soil available potassium (mg kg⁻¹) as affected by organic manure, NPK fertilization treatments and methods of application of the micronutrients.

Rate of applied O.M(A) ton/fed	Rate of N/P ₂ O ₅ /K ₂ O application (kg/fed)(B)	Micronutrients (C)							
		First season				Second season			
		Control 0.0	Soil appli.	Foliar appli.	Mean	Control 0.0	Soil appli.	Foliar appli.	Mean
0.0	0.0 60/15/24 120/30/48	6.05	7.55	7.25	6.95	9.33	9.45	9.42	9.40
		7.62	7.95	7.65	7.74	9.45	9.75	9.00	9.40
		7.65	7.97	7.77	7.80	9.83	9.93	9.87	9.88
	Mean	7.10	7.82	7.56	7.49	9.54	9.71	9.43	9.56
10	0.0 60/15/24 120/30/48	7.30	7.87	7.55	7.57	10.95	11.28	11.25	11.16
		8.60	8.79	8.63	8.67	11.15	11.39	11.26	11.27
		8.85	9.03	8.97	8.95	11.25	11.45	11.30	11.33
	Mean	8.25	8.56	8.38	8.40	11.12	11.37	11.27	11.25
20	0.0 60/15/24 120/30/48	9.15	9.30	9.20	9.22	12.37	12.70	12.58	12.75
		9.25	9.50	9.40	9.38	12.60	12.87	12.78	12.52
		9.70	9.80	9.75	9.75	12.70	12.90	12.73	12.79
	Mean	9.37	9.53	9.45	9.45	12.56	12.82	12.71	12.69
30	0.0 60/15/24 120/30/48	10.00	10.50	10.40	10.30	13.00	13.20	13.18	13.13
		10.80	10.90	10.85	10.86	13.20	13.50	13.27	13.32
		11.50	11.70	11.62	11.61	13.22	13.93	13.70	13.62
	Mean	10.77	11.03	10.95	10.92	13.14	13.54	13.38	13.35
Mean of NPK	0.0 60/15/24 120/30/48	8.12	8.80	8.60	8.51	11.41	11.66	11.61	11.56
		9.07	9.28	9.13	9.16	11.60	11.87	11.58	11.68
		9.42	9.62	9.53	9.52	11.75	12.05	11.90	11.90
Mean of micronutrients	8.87	9.23	9.13		11.59	11.86	11.70		

L.S.D at 0.05	First season							Second season						
	A	B	C	AB	AC	BC	ABC	A	B	C	AB	AC	BC	ABC
	0.13	N.S	N.S	N.S	N.S	N.S	N.S	0.15	N.S	N.S	N.S	N.S	N.S	N.S

Micronutrients availability:

Values of available micronutrients as affected by organic manure, NPK and micronutrients fertilization are presented in Tables (13-16). Contents of available Fe, Mn, Zn and Cu significantly increased by manure application. The increase was progressive with increasing rate of the applied manure. The increment contents of available micronutrients are due to the contents of such nutrients applied to the soil within the manure it self. In addition, the organic acids produced from FYM decomposition caused a reduction in soil pH, hence increase nutrients availability. These results are in harmony with the findings of Negm *et al.* (2002) who reported that organic manure improved the availability of nutrients.

Concerning macro and micronutrients, data in the above mentioned Tables revealed that soil available micronutrients, i.e. Fe, Mn, Zn and Cu were not affected by NPK fertilization, while addition of micronutrients as soil application significantly increased available studied micronutrients as compared with foliar application or with no micronutrients application. It is evident to notice that addition of micronutrients as foliar spray not affected the micronutrients availability when compared with no micronutrients application. The effect of soil application method on increasing Fe, Mn, Zn and Cu availability is mainly due to application of these nutrients in the sulphate form which are not easily leached from soil.

Table (13): Soil available iron (mg kg⁻¹) as affected by organic manure, NPK fertilization treatments and methods of application of the micronutrients.

Rate of applied O.M (A) ton/fed	Rate of N/P ₂ O ₅ /K ₂ O application (kg/fed) (B)	Micronutrients (C)							
		First season				Second season			
		Control 0.0	Soil appli.	Foliar appli.	Mean	Control 0.0	Soil appli.	Foliar appli.	Mean
0.0	0.0	1.70	3.50	1.80	2.33	1.40	3.15	1.60	2.05
	60/15/24	1.78	3.32	1.87	2.32	1.42	3.22	1.65	2.10
	120/30/48	1.85	3.80	1.90	2.51	1.50	3.24	1.98	2.24
	Mean	1.78	3.54	1.86	2.39	1.44	3.20	1.74	2.13
10	0.0	1.50	3.72	2.52	2.58	1.80	3.45	1.85	2.37
	60/15/24	2.60	3.80	2.53	2.98	1.90	3.60	1.95	2.48
	120/30/48	2.65	3.85	2.62	3.04	1.95	3.75	1.99	2.56
	Mean	2.22	3.65	2.56	2.81	1.88	3.60	1.93	2.47
20	0.0	2.80	4.12	2.85	3.25	2.12	4.03	2.22	2.79
	60/15/24	2.92	4.25	2.95	3.37	2.15	4.10	2.23	2.83
	120/30/48	2.95	4.45	2.99	3.46	2.20	4.15	2.34	2.90
	Mean	2.89	4.28	2.94	3.37	2.16	4.09	2.26	2.84
30	0.0	3.05	4.27	3.12	3.48	2.35	4.45	2.45	3.08
	60/15/24	3.12	4.35	3.15	3.54	2.50	4.52	2.57	3.19
	120/30/48	3.32	4.60	3.37	3.76	2.57	4.62	2.65	3.28
	Mean	3.16	4.41	3.21	3.59	2.47	4.53	2.56	3.19
Mean of NPK	0.0	2.26	3.90	2.57	2.91	1.92	3.77	2.03	2.57
	60/15/24	2.60	3.93	2.63	3.05	1.99	3.86	2.10	2.65
	120/30/48	2.69	4.18	2.72	3.20	2.06	3.94	2.24	2.75
Mean of micronutrients		2.51	3.97	2.66		1.98	3.86	2.12	

L.S.D at 0.05	First season							Second season						
	A	B	C	AB	AC	BC	ABC	A	B	C	AB	AC	BC	ABC
	0.08	N.S	0.08	N.S	N.S	N.S	N.S	0.14	N.S	0.08	N.S	N.S	N.S	N.S

Regarding the interaction effect, data in general, reveal that the micronutrients availability not significantly responded to the interaction between manuring and the methods of micronutrients application. The highest values of available Fe, Mn, Zn and Cu were recorded for the sand soil received 30 ton FYM/fed and supplied with micronutrients as sulphate salts. At the same time the soil without application of both FYM and micronutrients showed the lowest available contents of Fe, Mn, Zn and Cu.

Table (14): Soil available manganese (mg kg⁻¹) as affected by organic manure, NPK fertilization treatments and methods of application of the micronutrients.

Rate of applied O.M(A) ton/fed	Rate of N/P ₂ O ₅ /K ₂ O application (kg/fed)(B)	Micronutrients (C)							
		First season				Second season			
		Control 0.0	Soil appli.	Foliar appli.	Mean	Control 0.0	Soil appli.	Foliar appli.	Mean
0.0	0.0	1.12	1.80	1.20	1.37	0.80	1.30	0.90	1.00
	60/15/24	1.20	2.02	1.42	1.55	0.90	1.40	0.95	1.08
	120/30/48	1.73	2.32	2.22	2.09	0.92	1.42	0.99	1.11
	Mean	1.35	2.05	1.61	1.67	0.87	1.36	0.95	1.06
10	0.0	1.50	2.60	1.60	1.90	1.20	1.50	1.32	1.34
	60/15/24	1.60	2.70	1.68	1.99	1.25	1.60	1.38	1.41
	120/30/48	1.70	2.72	1.79	2.07	1.30	1.70	1.42	1.47
	Mean	1.60	2.72	1.69	2.00	1.25	1.60	1.37	1.41
20	0.0	1.72	3.22	1.80	2.25	1.60	1.90	1.63	1.70
	60/15/24	2.03	3.52	2.15	2.57	1.70	1.93	1.76	1.79
	120/30/48	2.22	3.55	2.92	2.89	1.80	1.99	1.88	1.89
	Mean	1.99	3.43	2.29	2.57	1.70	1.94	1.76	1.80
30	0.0	2.23	4.17	2.30	2.90	1.03	2.20	2.00	1.74
	60/15/24	2.33	4.30	2.35	2.99	1.90	2.32	2.15	2.12
	120/30/48	2.52	4.50	2.65	3.22	2.13	2.40	2.20	2.24
	Mean	2.36	4.33	2.43	3.04	1.69	2.31	2.12	2.04
Mean of NPK	0.0	1.64	2.95	1.92	2.17	1.16	1.72	1.46	1.45
	60/15/24	1.79	3.13	1.90	2.27	1.44	1.81	1.56	1.60
	120/30/48	2.04	3.27	2.39	2.57	1.53	1.63	1.62	1.59
Mean of micronutrients		1.82	3.13	2.00		1.38	2.40	1.55	

L.S.D at	First season							Second season						
	A	B	C	AB	AC	BC	ABC	A	B	C	AB	AC	BC	ABC
	0.05	0.11	N.S	0.06	N.S	N.S	N.S	N.S	0.12	N.S	0.09	N.S	N.S	N.S

Table (15): Soil available zinc (mg kg⁻¹) as affected by organic manure, NPK fertilization treatments and methods of application of the micronutrients.

Rate of applied O.M(A) ton/fed	Rate of N/P ₂ O ₅ /K ₂ O application (kg/fed)(B)	Micronutrients (C)							
		First season				Second season			
		Control 0.0	Soil appli.	Foliar appli.	Mean	Control 0.0	Soil appli.	Foliar appli.	Mean
0.0	0.0	0.65	1.12	0.70	0.82	0.60	0.92	0.70	0.74
	60/15/24	0.82	1.20	0.85	0.96	0.70	1.00	0.75	0.82
	120/30/48	0.95	1.30	0.99	1.08	0.80	1.22	0.85	0.96
	Mean	0.81	1.21	0.85	0.96	0.70	1.05	0.77	0.84
10	0.0	1.25	1.60	1.31	1.39	0.80	1.20	0.90	0.97
	60/15/24	1.30	1.77	1.31	1.46	0.85	1.22	0.98	1.02
	120/30/48	1.32	1.92	1.83	1.69	0.92	1.32	1.30	1.18
	Mean	1.29	1.76	1.48	1.51	0.86	1.25	1.06	1.06
20	0.0	1.52	1.83	1.60	1.65	1.20	1.52	1.22	1.31
	60/15/24	1.62	1.93	1.70	1.75	1.22	1.55	1.25	1.34
	120/30/48	1.65	1.99	1.88	1.84	1.27	1.62	1.31	1.40
	Mean	1.59	1.92	1.73	1.75	1.23	1.56	1.26	1.35
30	0.0	1.60	2.12	1.70	1.81	1.42	1.60	1.50	1.51
	60/15/24	1.70	2.13	1.92	1.92	1.47	1.70	1.55	1.57
	120/30/48	1.80	2.33	1.98	2.04	1.50	1.80	1.65	1.65
	Mean	1.70	2.19	1.87	1.92	1.47	1.70	1.57	1.58
Mean of NPK	0.0	1.25	1.67	1.33	1.42	1.00	1.31	1.08	1.13
	60/15/24	1.36	1.73	1.44	1.51	1.06	1.37	1.13	1.19
	120/30/48	1.43	1.88	1.67	1.66	1.12	1.28	1.28	1.23
Mean of micronutrients		1.35	1.77	1.48		1.06	1.39	1.17	

L.S.D at 0.05	First season							Second season						
	A	B	C	AB	AC	BC	ABC	A	B	C	AB	AC	BC	ABC
	0.11	N.S	0.06	N.S	N.S	N.S	N.S	0.12	N.S	0.06	N.S	N.S	N.S	N.S

Table (16): Soil available copper (mg kg⁻¹) as affected by organic manure, NPK fertilization treatments and methods of application of the micronutrients.

Rate of applied O.M (A) ton/fed	Rate of N/P ₂ O ₅ /K ₂ O application (kg/fed) (B)	Micronutrients (C)							
		First season				Second season			
		Control 0.0	Soil appli.	Foliar appli.	Mean	Control 0.0	Soil appli.	Foliar appli.	Mean
0.0	0.0 60/15/24 120/30/48	0.40	0.70	0.42	0.51	0.32	0.60	0.35	0.42
		0.50	0.80	0.52	0.61	0.35	0.70	0.37	0.47
		0.55	0.92	0.57	0.68	0.40	0.75	0.42	0.52
		Mean	0.48	0.81	0.50	0.60	0.36	0.68	0.38
10	0.0 60/15/24 120/30/48	0.70	0.80	0.72	0.74	0.50	0.70	0.52	0.57
		0.75	0.85	0.82	0.81	0.60	0.85	0.62	0.69
		0.82	0.92	0.87	0.87	0.62	0.92	0.69	0.74
		Mean	0.76	0.86	0.80	0.81	0.57	0.82	0.61
20	0.0 60/15/24 120/30/48	0.87	1.20	0.89	0.99	0.85	1.00	0.87	0.91
		0.98	1.30	1.00	1.09	0.92	1.12	0.94	0.99
		1.00	1.35	1.12	1.16	0.98	1.20	0.99	1.06
		Mean	0.95	1.28	1.00	1.08	0.92	1.11	0.93
30	0.0 60/15/24 120/30/48	1.30	1.80	1.40	1.50	1.10	1.40	1.17	1.22
		1.40	2.03	1.50	1.64	1.22	1.45	1.24	1.30
		1.50	2.92	1.60	2.00	1.32	1.55	1.35	1.41
		Mean	1.40	2.25	1.50	1.72	1.21	1.47	1.25
Mean of NPK	0.0 60/15/24 120/30/48	0.82	1.12	0.86	0.93	0.69	0.77	0.73	0.74
		0.91	1.24	0.96	1.04	0.77	0.84	0.79	0.80
		0.97	1.53	1.04	1.18	0.83	1.10	0.86	0.93
Mean of micronutrients		0.90	1.30	0.95		0.77	1.02	0.79	

L.S.D at 0.05	First season							Second season						
	A	B	C	AB	AC	BC	ABC	A	B	C	AB	AC	BC	ABC
	0.10	N.S	0.05	N.S	N.S	N.S	N.S	0.11	N.S	0.05	N.S	N.S	N.S	N.S

REFERENCES

- Ali, Maha, M.E. (2007). Growth of wheat plant on a light textured soil as influenced by water stress and some soil conditioners. M.Sc. Thesis, Fac. of Agric., Benha Univ., Egypt.
- Atta Allah, S.A. and Mohamed, G.A. (2003). Response of wheat grown in newly reclaimed sandy soil to poultry manure and nitrogen fertilization. J. Agric. Sci., Mansoura Univ., (28): 7531-7538.
- Badr, M.M.A.; Hassan, M.A.M. and Monged, Nadia, O.L. (1991). Effect of nitrogen application and micronutrients on the yield and chemical contents of wheat plant. Zagazig J. Agric. Res., 18(5): 1661-1668.
- El-Ghamry, A.; Elsirafy, Z.M. and El-Dissoky, R.A. (2005). Response of potato grown on clay loam soil to sulfur and compost application. J. Agric. Sci., Mansoura Univ., 30(7): 4337-4353.
- El-Koumey, B.Y. (1998). Influence of Zn, Cu and farmyard manure on wheat plants. Zagazig J. Agric. Res. 25(4): 687-697.

- El-Lethi, A.A.; Sayed, K.M. and El-Yamani, M.S. (1996). Influence of different levels of N, K and Zn fertilization on wheat yield and chemical composition in salt affected soil. *J. Agric. Sci. Mansoura Univ.*, 21(10): 3735-3741.
- Galal, O.A.M. (2007). Studies on balanced fertilization of wheat plant. Ph.D. Thesis, Fac. of Agric., Benha Univ., Egypt.
- Jackson, M.L. (1973). *Soil Chemical Analysis*, Printice Hall, New Delhi, India.
- Khalil, F.A. and Aly, S.A. (2004). Effect of organic fertilizers as substitutions of mineral nitrogen fertilizer applied at planting on yield and quality of wheat. *Minufiya J. Agric. Res.*, (2): 435-449.
- Klute (1982). *Methods of Analysis. Part 1: Physical and Mineralogical Methods (2nd ed)*. American Society of Agronomy, Madison, Wisconsin, USA.
- Mekail, M.M.; Hassan, H.A.; Mohamed, W.S.; Telep, A.M. and Abd El-Azeim, M.M. (2006). Integrated supply system of nitrogen for wheat grown in the newly reclaimed sandy soils of West El-Minia: Efficiency and economics of the system. *Minia J. Agric. Res. and Dev.*, 26(1): 101-103.
- Mohamed, K.A. (1994). The effect of foliar spray of wheat with Zn, Cu, Fe and urea on yield, water use efficiency and nutrients uptake at different levels of soil salinity. *Assiut Journal of Agricultural Sciences*, 25(3): 179-189.
- Negm, M.A.; El-Sayed M.H.; Ahmed, A.S. and Adel-Ghani, M.M. (2002). a: Wheat and sorghum response to biocomposite compost and sulphur added to a calcareous soil on II. Cereal productivity and nutrient uptake. *Minufiya J. Agric. Res.*, 27(2): 381-390.
- Page, A.L.; Miller, R.H. and Keeny, D.R. (1982). *Methods of Soil Analysis. Part 2: Chemical and Microbiological Properties. (2nd ed)*. American Society of Agronomy, Madisons, Wisconsin, USA.
- Salem, F.S. (2003). Effect of some soil amendments on the clayey soil properties and some crops production. *Minufiya J. Agric. Res.*, 28(2): 1705-1715.
- Sendecor, G.W. and Cochran W.G. (1980). "Statistical Methods" 7th Edin. Iowa State Univ., Press, Iowa, USA.
- Shams El-Din, H.A.I. (1993). Application methods and rates of some micronutrients on wheat plant. Ph.D. Thesis, Fac. of Agric., El-Mansoura Univ., Egypt.
- Taha, M.B. (2007). Recycling of organic wastes for using as soil amendments. Ph.D. Thesis, Fac. of Agric. Minia Univ., Egypt.

تأثير الأسمدة العضوية المثراه بالعناصر الكبرى والصغرى على القمح فى الأراضى الرملية

٢- امتصاص العناصر وصلاحتها

محمد السيد على* ، صفوت أحمد إسماعيل** ، عمر حسيني محمد الحسينى* و أحمد محمد عبد الحفيظ**

* قسم الأراضى - كلية الزراعة - جامعة بنها

** معهد بحوث الأراضى والمياه والبيئة - مركز البحوث الزراعية

أقيمت تجربتان حقليتان بالقربية الأولى غرب سمالوط بمحافظة المنيا، ممثلة للأراضى الرملية الحديثة الاستصلاح لدراسة تأثير أربع مستويات من التسميد العضوى (صفر، ١٠، ٢٠، ٣٠ طن سماد بلدى للفدان) ثلاث مستويات من السماد المخلوط من النتروجين والفوسفور والبوتاسيوم (بدون تسميد، مستوى منخفض (٢٤/١٥/٦٠) ومستوى مرتفع (٤٨/٣٠/١٢٠) كجم نتروجين، فوسفور، بوزا للفدان على الترتيب) وكذلك إضافة العناصر الصغرى (حديد + منجنيز + زنك + نحاس) بطرق مختلفة وهى بدون إضافة وإضافة أرضية قبل الزراعة والإضافة رشاً على النبات، على امتصاص العناصر الصغرى والكبرى لنبات القمح وكذلك صلاحيتها فى التربة. ويمكن تلخيص أهم النتائج فيما يلى:

أزداد امتصاص عناصر النتروجين والفوسفور والبوتاسيوم والحديد والمنجنيز الزنك والنحاس بنبات القمح بزيادة كلاً من التسميد بالأسمدة العضوية والعناصر الكبرى وكذلك إضافة العناصر الصغرى مع أفضلية لأضافتها رشاً عن الإضافة الأرضية.

كما استجابت صلاحية عناصر النتروجين والفوسفور والبوتاسيوم بالتربة للتسميد العضوى فقط، بينما أزداد صلاحية الفوسفور بزيادة التسميد بالعناصر الكبرى.

وتأثرت صلاحية الحديد والمنجنيز والزنك والنحاس بإضافة كلاً من الأسمدة العضوية والعناصر الصغرى أرضاً قبل الزراعة، بينما لم تتأثر صلاحية هذه العناصر بإضافة أسمدة العناصر الصغرى رشاً على النبات فى موسمى الزراعة.

ويمكن من هذه الدراسة التوصية بإضافة الأسمدة العضوية حتى ٣٠ طن/فدان وأسمدة النتروجين والفوسفور والبوتاسيوم بمعدل ١٢٠، ٣٠، ٤٨ كجم ن، فوسفور، بوزا على الترتيب مع التوصية بإضافة العناصر الصغرى لتحسين امتصاص العناصر بواسطة نبات القمح وكذلك صلاحيتها فى التربة.