

The Frequency of Palmaris Longus Absence among Female Students in King Faisal University in Al-Ahsa, Saudi Arabia

Alabbad, Aqilah A' Alkhamis, Marwah H, Alsultan, Marwah S, Alahmad, Sarah A

College of Medicine, King Faisal University

Corresponding author: Aqilah Ali Alabbad, E-mail: aana-213@hotmail.com

ABSTRACT

Background: Palmaris longus (PL) is one of the forearm muscles that lie between the flexor carpi ulnaris and the flexor carpi radialis muscles. PL action is flexion of the hand at the wrist and making the palmar aponeurosis tense. Plastic surgeons utilize the Palmaris longus in restoration of lip and chin defects.

Objectives: We sought to determine the frequency of the absence of the palmaris longus in Saudi Arabia among female students in King Faisal University, AL-Ahsa. **Materials and Methods:** Two hundred normal subjects were chosen randomly from King Faisal University female students. Subjects who had gone through a surgical procedure or have any deformities in the forearm were excluded. We have examined the presence or absence of palmaris longus using three tests. Subjects were asked to do standard test for the assessment of PL tendon. If PL cannot be detected by the standard test, two more tests were performed to confirm the absence. **Results:** The overall prevalence of absence both unilaterally and bilaterally is 40.5 %. Unilateral absence was 20.5%. The bilateral absence was 20%. The distribution on the right and left was 29% and 31.5% respectively.

Conclusions: The present study found palmaris longus to be absent equally bilateral and unilateral in more than one third of the sample and significantly more common in the left side.

Keywords: Palmaris longus, agenesis, forearm, Saudi Arabia.

INTRODUCTION

The palmaris longus (PL) is a slender, spindle shaped weak flexor of the wrist and it is located medial to the flexor carpi radialis (FCR). The number and form of PL muscle is extremely variable^[1]. The PL muscle may be absent, double, split, tendinous, digastric and may have various insertions. It may be inserted on the flexor retinaculum, the fascia of the forearm, the fascia and the muscles of the hypothenar, the short abductor of the thumb, near the metacarpophalangeal joints, the tendon of the flexor carpi ulnaris muscle, the pisiform bone or the scaphoid bone^[2].

The presence of the PL can be determined through noninvasive and standard physical examination of the volar wrist^[3].

Considering its dispensability, surgeons agree that PL is the best choice for tendon grafts in tendon reconstruction, helping the function of paralyzed muscles used for repairing ptosis, treatment of facial paralysis, and urinary incontinence. Plastic surgeons also utilize the PL in lip augmentation and restoration of lip and chin defects. As its absence is immaterial to the function of the wrist, PL has the greatest variation in the human body and its most common variation is agenesis (PLA)^[4]. However, this muscle variation can lead to median and/or ulnar nerve compression syndromes in some rare cases^[5].

The rate of congenital absence of the PL is estimated to be 15% among individuals worldwide and its incidence is population-dependent ranging from

0.6% (in the Korean population) to 63.9% (in the Turkish population)^[6, 7].

A lot of studies have been made regarding the frequency of palmaris longus absence in Korea^[8], Iran^[9], Egypt^[10], and East Africa^[11] and others. However, the frequency of palmaris longus absence have not been studied yet in Alahsa.

Taking into account the large variability of palmaris longus presence, the aim of this study is to investigate the frequency of Palmaris longus absence among female students in king Faisal University and to compare the results to the other studies worldwide. The results of our study are going to be useful for the surgeons working in our population.

MATERIALS AND METHODS

A cross sectional study was designed to investigate the absence of Palmaris longus in both sides in a sample of normal population which was taken from female students of king Faisal University included 200 students randomly. Their ages range between 18 to 25 years old. We excluded the subjects who have done any surgical procedure in the forearms or have any deformities in the forearms.

The study was done after approval of ethical board of King Faisal university.

We have examined the presence of Palmaris Longus muscle using three tests. The subject was

initially asked to do the standard test for the assessment of the PL tendon. If the tendon was not visualized, 2 additional tests were done to confirm the absence. The standard test is Schaeffer's test and the other tests are Thompson's test and Pushpakumar's test.

	Number of cases	percentage
total	200	100%
Overall absence	81	40.5%
Bilateral absence	40	20%
Unilateral absence	41	20.5%
Right side absence	58	29%
Left side absence	63	31.5%

Table (1): The frequency of palmaris longus absence

In Schaeffer's test (figure 1) the subject is asked to oppose the thumb to the little finger and then flex the wrist [12].

Thompson's test (figure 2) involves flexion of the fingers to form a fist followed by flexion of the wrist and finally the thumb is opposed and flexed over the fingers [13]. Pushpakumar's test (figure 3) involves extension of the index and middle finger with flexion of the other fingers and the wrist followed by opposition and flexion of the thumb [14].

The presence or absence of the palmaris longus tendon was recorded for both sides.



Figure 1 [15]
Schaeffer's test



Figure 2 [15]
Thompson's test



Figure 3 [15]
Pushpakumar's test

Using the SPSS statistics 17.0, we've worked out the frequency of palmaris longus absence in the right and left forearms and watched the differences between the two sides. We used the Chi-square test to see if there is a significant difference between them. A value of $p < 0.05$ was considered to indicate a significant difference between two sides [16].

RESULTS

We examined 200 female students in King Faisal University in Al-Ahsa. The overall frequency of palmaris longus absence, either bilaterally or unilaterally, was found to be 40.5 % (81 subjects). Palmaris longus was absent unilaterally in 20.5% (41 subjects) and bilaterally in 20% (40 subjects). The overall difference between right and left in term of absence of palmaris longus was statistically significant ($P < 0.05$). Palmaris longus was absent in 29% (58 subjects) in the right side and in 31.5% (63 subjects) in the left side.




Figure 4: Absence of Palmaris longus among female students in KFU.

DISCUSSION

Palmaris longus (PL) is one of the forearm muscles that lie between the flexor carpi ulnaris and the flexor carpi radialis muscles [17]. The absence of the PL has been shown to vary based on body side, gender, and ethnicity in prior studies [18]. Most studies point out that the absence of PLM is most commonly encountered in women and in left arms [5].

In Our study, it was found that the overall prevalence of palmaris longus absence among female students of KFU to be 40.5%. This value is comparable to the high prevalence of PL absence

among females in the Andhra population of India (40.2%)^[19], Bahraini population (36.8%)^[20], and Serbian population (37.5%)^[3]. On the other hand, the overall absence is found to be very high in relation to the prevalence of PL absence reported in Korean population (4.1%)^[8] and Yoruba population (6.4%)^[21]. Higher frequencies was found in Pakistani population (47.12%)^[22] and Egyptian population (50.8%)^[10]. This reaffirms that the prevalence of absence of the palmaris longus muscle shows a marked variation in different populations.

The high prevalence of Palmaris longus absence in our study coincides with high prevalence reported in Middle East as in Jordanian^[23] and Bahraini population^[20]. Also, the fact that our sample included only female subjects explains the high prevalence founded in the present study as many studies revealed the same result before^[19, 10, 24].

On average, the prevalence of unilateral absence of palmaris longus is higher than that of the bilateral absence accounting for 16% and 9%, respectively, of the population all over the world^[25]. The present study revealed almost equal frequencies of unilateral (20.5%) and bilateral (20%) absence of PL. Similar results regarding the equality of the unilateral and bilateral absence was found in South Africa giving unilateral absence frequency of 6% and bilateral absence frequency of 5.5%^[26]. Some studies that reported huge differences between the frequencies of unilateral (43.2%) and bilateral (20.6%) absence is the study in Indian population^[25].

Our study found that the difference in the prevalence of PL absence between the right and the left side to be significant which is the same result founded in Indian population^[27] and Serbian population^[3], whereas no significant difference was found in Bahraini population^[20] and Korean population^[8].

It has been documented that the absence of PLM is hereditary and HOX is the gene responsible for the regulation of its morphological development. Also, it has been proposed that a possible dominant expression of genes is responsible for PLM variations in family members^[5]. However, we were not able to demonstrate that in our study since our subjects included college students who were randomly selected.

One of the weaknesses of the present work is that our study included only female subjects and we weren't able to compare the results based on gender. Another weakness is that the presence of palmaris longus was determined by clinical exams which depend on the examiners to a large extent and is not

considered as a definite conclusion. We tried to overcome this weakness by performing three tests to detect the presence of PL.

CONCLUSION

In conclusion, this study demonstrates absence of palmaris longus (PL) muscle in more than one-third of the female Saudi population with almost equal bilateral and unilateral absence. Also, PL was absent on left side more than right one. As palmaris longus is the first choice for tendon grafts and reconstructive surgeries, by the virtue of its structure and function, more researches should be done regarding its anatomy and variations. We suggest that more studies need to be done regarding the use of palmaris longus muscle on reconstructive and plastic surgeries worldwide and here in Saudi Arabia and to compare the use of palmaris longus versus other muscles and their advantages and disadvantages.

ACKNOWLEDGEMENTS

The authors of this work wish to thank all the staff in college of Medicine, King Faisal University for their help in collecting the sample and doing the research.

REFERENCES

1. **Yildiz M, Sener M, Aynaci O (2000):** Three-headed reversed palmaris longus muscle: a case report and review of the literature. *Surg Radiol Anat.*, 22:217-9.
2. **Lalit M, Singla RK, Piplani S Bifi d (2014):** inverted palmaris longus muscle – a case report. *Eur J Anat.*, 18(3):341-3.
3. **Erić M, Koprivčić I, Vučinić N (2011):** Prevalence of the palmaris longus in relation to the hand dominance. *Surgical and Radiologic Anatomy*, 33 (6):481–484.
4. **Thompson NW, Mockford BJ, Cran GW (2001):** Absence of the palmaris longus muscle: a population study. *The Ulster Med J.*, 70: 22 – 24.
5. **Georgiev GP, Iliev AA, Dimitrova IN, Kotov GN, Malinova LG, Landzhov BV (2017):** Palmaris Longus Muscle Variations: Clinical Significance and Proposal of New Classifications. *Folia Med (Plovdiv)*, 59(3):289-297.
6. **Sebastin SJ, Lim AY, Bee WH, Wong TC, Methil BV (2005):** Does the absence of the palmaris longus affect grip and pinch strength?. *Hand Surg Br.*, 30:406-8.
7. **Hiz Ö, Ediz L, Ceylan MF, Gezici E, Gülcü E, Erden M (2011):** Prevalence of the absence of palmaris longus muscle assessed by a new examination test (Hiz-Ediz Test) in the population residing in the area of Van, Turkey. *J Clin Exp Invest.*, 2:254-9.
8. **Kyung DS, Lee JH, Choi IJ, Kim DK (2012):** Different frequency of the absence of the palmaris longus according to assessment methods in a Korean population.. *Anat cell biol.*, 45(10):53-56.

9. **Lahiji FA , Ashoori K, Dahmardehei M (2013):** Prevalence of Palmaris Longus Agenesis in a Hospital in Iran. *Archives of Iranian medicine*, 16(3):187-8.
10. **Raouf HA, Kader GA, Jaradat A, Dharap A, Fadel R, Salem A (2013):** Frequency of palmaris longus absence and its association with other anatomical variations in the Egyptian population. *Clin. Anat.*, 26(5):572-7.
11. **Kigera JWM, Mukwaya S (2011):** Frequency of Agenesis Palmaris Longus through Clinical Examination – An East African study. *PLoS ONE*, 6(12): e28997.
12. **Schaeffer JP (1909):** On the variations of the palmaris longus muscle. *Anat Rec.*, 3:275-8.
13. **Thomson JW, McBatts J, Danforth CH (1921):** Hereditary and racial variations in the muscles Palmaris Longus. *Am J Phys Anthropol.*, 4:205-20.
14. **Pushpakumar SB, Hanson RP, Carroll S (2004):** Clinical examination of Palmaris Longus (PL) tendon. *Br J Plast surg.*, 57:184-5.
15. **Enye LA, Saalu LC, Osinubi AA (2010):** The Prevalence of Agenesis of Palmaris Longus Muscle amongst Students in Two Lagos-Based Medical Schools. *Int. J. Morphol.*, 28(3):849-854.
16. **Duncan, D. B (1957):** Multiple range tests for correlated and heteroscedastic means. *Biometrics*, 13:164-76.
17. **Williams PL, Bannister LH, Berry MM, Collins P, Dyson M, Dussek JE, Ferguson MWJ (1999):** Cardiovascular system. In: Gabella G, editor. *Gray's Anatomy*. 38th Ed. Edinburgh, London: Churchill Livingstone, 1542–1544.
18. **Singh D, Kumar KA, M. Dinesh C, Raj R (2012):** Chronic triceps insufficiency managed with extensor carpi radialis longus and palmaris longus tendon grafts. *Indian Journal of Orthopaedics*, 46(2):236-8.
19. **Sankar KD, Bhanu PS, John SP (2011):** Incidence of agenesis of palmaris longus in the Andhra population of India. *Indian J Plast Surg.*, 44(1): 134–138.
20. **Sater MS, Dharap AS, Abu-Hijleh MF (2010):** The prevalence of absence of the Palmaris longus muscle in the Bahraini population. *Clin. Anat.*, 23:956-961.
21. **Mbaka GO, Ejiwunmi AB (2009):** Prevalence of Palmaris longus absence – a study in the Yoruba population. *Ulster Med J.*, 78(2):90-93.
22. **Mahmood T, Butt KA, Ahmed N, Shahid R, Hussain FN (2011):** Prevalence of congenital absence of Palmaris longus tendon in healthy volunteers- a cross sectional study. *Pakistan Journal of Medical and Health Sciences*, 5(1):144-146.
23. **Abu Hassan F, Jabaiti S (2008):** Absence of the Palmaris longus tendon in Mid Eastern population. *Journal of the Bahraini Medical Society*, 20:70-73.
24. **Kose O, Adanir O, Cripar M, Kurklu M, Komurcu M (2009):** The prevalence of Palmaris longus: a study in Turkish population. *Arch Orthop Trauma Surg.*, 129(5):609-11.
25. **Agarwal P (2010):** Absence of Palmaris longus tendon in Indian population. *India J Orthop.*, 44:212-5.
26. **Ndou R, Gangata H, Mitchell B, Ngongo T, Louw G (2010):** The frequency of absence of Palmaris longus in a south African population of mixed race. *Clin Anat.*, 23(4):437-42.
27. **Kapoor SK, Tiwari A, Kumar A, Bhatia R, Tantuway V, Kapoor S (2008):** Clinical relevance of palmaris longus agenesis: common anatomical aberration. *Anat Sci Int.*, 83(1):45-8.